

# COMPARISON OF PRECIPITATION TRENDS AND ITS EFFECT ON LAKE WATER LEVELS USING SATELLITE ALTIMETRY:A CASE STUDY OF LAKES KAINJI AND SHIRORO



BY

HAWA ABDULAI , RAKIYA BABAMAAJI, AWE-PETER HELEN, DANLAMI MUHAMMED  
NATIONAL SPACE RESEARCH AND DEVELOPMENT AGENCY,NIGERIA

## INTRODUCTION

- In all hydrological systems, precipitation (in different forms) has been found to be a major driver/component; In humid regions, water losses are usually through direct precipitation (Cretaux et al.,(2011).
- Precipitation trends help to reveal recurring patterns as well as differences that can aid in the understanding of water fluctuations, to enable proper planning and forecasting, and to ultimately prevent extreme events such as droughts and floods as a result of climate change.
- Lakes on the other hand are viewed as perfect proxies for climate change and their monitoring has aided in observing trends that has helped in the understanding of climatic parameters (Cretaux et al.,(2006). This monitoring has been made possible since 1993 by the use of satellite altimetry with the introduction of the Topex/Poseidon, Jason 1,2 and 3 missions where pulses are sent to lake water surfaces from radar altimeters and the height difference is measured.
- The Kainji dam is the largest of the dams on the River Niger, the Shiroro Lake was created from the River Kaduna, which is a tributary of the River Niger. The Lakes Kainji and Shiroro were created for the purpose of hydro-electric power generation, flood control and fishing, these purposes have not been fully achieved due to water fluctuations in the lakes, hence this study.
- The lake water levels should not be confused with the depth as the Lake water level is measured in relation to a reference point; usually the mean sea level which is taken as 0
- The choice of the two lakes is due to their similarities based on their usage for hydropower generation. Their differences based on their sources and their sizes were also reasons for comparison to determine if there are unique climatic controls on water level fluctuations in the two lakes

## THE STUDY AREA




Fig 1: Lakes Kainji and Shiroro, North Central Nigeria

## METHODOLOGY

- Rainfall data for Lakes Kainji and Shiroro were obtained from Tropical Applications of Meteorology using SATellite and ground-based observations (TAMSAT).A trend analysis (Mann Kendall) was performed using the Excel Stat software 2023 to determine if there were positive trends in the rainfall data.
- The Mann Kendall is a non-parametric test which involves accepting an alternative hypothesis ( $H_a$ ) which indicates the presence of a monotonic trend and rejecting the null hypothesis, ( $H_0$ ) which indicates that no monotonic trend was present.
- The processed data for Lake Water Levels for the 14 year period was derived from hydroweb, the satellite tracks where the data was obtained for Lake Kainji are :Topex Poseidon, Jason 2,3 and sentinel 6A on track 135 respectively.Envisat and saral altika on track 874 respectively. (see fig 3)For Lake Shiroro, it was obtained from Jason 2,Jason 3 and sentinel 6A on track no 20 respectively. The water levels data collected were averaged into yearly data for the 14year period.




Fig 2:Work flow for the study




Fig 3:Radar altimetry tracks for Lake kainji




Fig 4:Radar altimetry tracks for Lake Shiroro




Fig 5:Sen's slope graph for rainfall at Lake Kainji (Note positive trend)




Fig 6 :Sen's slope graph for rainfall at Lake Shiroro (Note negative trend)

## RESULTS

- From the trend analysis, the null hypothesis was accepted albeit a slight positive trend for Lake Kainji was noticed from the Sen's slope estimator with a positive Z value for Kendall's tau ( see fig 8a ).The same trend was noticed for the Lake water levels, as the null hypothesis was also accepted, with a slight positive trend noticed from the Sen's slope estimator. For Lake Shiroro, the Null hypothesis was also accepted with no positive trend noticed and a negative Z value (see fig 8b) Correlation analysis between rainfall and Lake water level showed a very weak correlation for Lake Kainji and a weaker relationship was seen for lake Shiroro.
- The slight positive trend for Lake Kainji might be as a result of the Lake being an endorheic Lake. Endorheic lakes are usually shallow as in this case and they are largely affected by inflow parameters such as rainfall, temperature, hence they are strong indicators for climate change.




Fig 7:Correlating Rainfall and LWL for Lake Kainji




Fig 10:Correlating Rainfall and LWL for Lake Shiroro




Fig 12:Sen's slope graph for LWL at Lake Kainji (note positive trend)

Kendall's tau	0.155
S	14
Var(S)	332.667
p-value (Two-tailed)	0.476
alpha	0.05

Fig 8a

Kendall's tau	-0.121
S	-11
Var(S)	333.667
p-value (Two-tailed)	0.584
alpha	0.05

Fig 8b

Kendall's tau	0.133
S	12
Var(S)	332.667
p-value (Two-tailed)	0.546
alpha	0.05

Fig 8c

Kendall's tau	-0.209
S	-19
Var(S)	333.667
p-value (Two-tailed)	0.324
alpha	0.05

Fig 8d

Figs 8a and 8b :Summary statistics (Mann Kendall) for rainfall in Lakes Kainji and Shiroro respectively

8c and 8d:Summary statistics (Mann Kendall) for LWL in Lakes Kainji and Shiroro respectively




Fig 9: Rainfall and LWL plot for Lake Kainji




Fig11: Rainfall and LWL plot for Lake Shiroro




Fig 13:Sen's slope graph for LWL at Lake Shiroro (Note negative trend)

## CONCLUSION

The effects of precipitation trends on Lake water levels for Lakes Kainji and Shiroro are similar albeit with a slight difference for Lake Kainji where there is a slight positive trend. Validation with insitu data can provide more insights into the dominant climatic and hydrological factors that affect the lake water levels. This further investigation is more recommended for Lake Kainji which is a shallow lake and a slight positive trend was noticed for the slope of rainfall and Lake water level during the trend analysis conducted.

## REFERENCES

- Crétau, J. F., & Birkett, C. (2006). Lake studies from satellite radar altimetry. *Comptes Rendus Geoscience*, 338 (14-15), 1098-1112
- Crétau, J. F., Calmant, S., Del Rio, R. A., Kouraev, A., Bergé-Nguyen, M., & Maisongrande, P. (2011). Lakes studies from satellite altimetry. *Coastal altimetry*, 509-533.
- Cretaux J-F, Arsen A., Calmant S., et al., (2011). A lake database to monitor in the Near Real Time water level and storage variations from remote sensing data, *Advances in space Research*(Vol. 47, 1497-1507)
- Dai, L., Liu, H., Li, W., & Tang, Z. (2020, March). Trend and abrupt analysis of Dongting Lake water level and its driving forces using Mann-Kendall approach. In *IOP Conference Series: Earth and Environmental Science* (Vol. 467, No. 1, p. 012140). IOP Publishing.
- Salami, A. W., Mohammed, A. A., Adeyemo, J. A., & Olanlokun, O. K. (2015). Assessment of impact of climate change on runoff in the Kainji lake basin using statistical methods. *International Journal of Water Resources and Environmental Engineering*, 7(2), 7-16.

Acknowledgement: The Authors will like to acknowledge hydroweb and TAMSAT for providing data for this study