

# On the variability of the semidiurnal solar and lunar tides of the equatorial electrojet during sudden stratospheric warmings

Tarique Adnan Siddiqui<sup>1</sup>

<sup>1</sup>GFZ Potsdam

November 24, 2022

## Abstract


The variabilities of the semidiurnal solar and lunar tide of the equatorial electrojet (EEJ) are investigated during the 2003, 2006, 2009 and 2013 major sudden stratospheric warming (SSW) events in this study. For this purpose, the ground-magnetometer recordings at the equatorial observatories in Huancayo and Fuquene are utilized. Results show a major enhancement in the amplitude of the EEJ semidiurnal lunar tide in each of the four warming events. The EEJ semidiurnal solar tidal amplitude shows an amplification prior to the onset of warmings, a reduction during the deceleration of the zonal mean zonal wind at 60°N and 10hPa and a second enhancement a few days after the peak reversal of the zonal mean zonal wind during all the four SSWs. Results also reveal that the amplitude of the EEJ semidiurnal lunar tide becomes comparable or even greater than the amplitude of the EEJ semidiurnal solar tide during all these warming events. The present study also compares the EEJ semidiurnal solar and lunar tidal changes with numerical simulations of the variability of the migrating semidiurnal solar (SW2) and lunar (M2) tide in neutral temperature at ~120km altitude. A better agreement between the enhancements of the EEJ semidiurnal lunar tide and the M2 tide in neutral temperature is observed in comparison with the enhancements of the EEJ semidiurnal solar tide and the SW2 tide in neutral temperature.

## Introduction


### Sudden Stratospheric Warmings (SSWs)

- SSWs are large-scale meteorological events usually occurring during the northern hemisphere winters. SSW was first observed by Richard Scherhag at the Free University of Berlin in 1952.
- SSWs are characterized by a weakening or sometimes even a reversal of the westerly winds in the northern stratosphere that leads to a sudden rise in polar stratospheric temperature by several tens of degrees.
- The underlying mechanism behind SSWs is understood to be the nonlinear interaction of the vertically propagating planetary waves with the zonal mean flow (Matsuno, 1971).

### The 2018 SSW event




### SSW related impacts in the atmosphere



### The Equatorial Electrojet (EEJ)

- EEJ is a narrow band of an intense electric current flowing during daytime above the dip equator in the ionospheric E-region.




### Motivation and goals of this study

- The EEJ variability due to SSWs is believed to be due to the SSW-induced modulation of the atmospheric tides. In particular, the changes in the semidiurnal solar and lunar tides have been found to be the major source of ionospheric variabilities during SSWs. The main purpose of this work is to investigate the variability of EEJ semidiurnal solar and lunar tidal enhancements with respect to the occurrence of SSWs and also to study the relative enhancements of the EEJ semidiurnal solar and lunar tides during SSWs.

## Results

### Semidiurnal solar and lunar tides of the EEJ from ground-magnetometer recordings




- The EEJ amplitudes presented in these figures are normalized to a solar flux level of 150 s.f.u.
- The semidiurnal perturbation pattern in the EEJ that increasingly shifts in local time on succeeding days can be seen near the new moon or full moon days during all the four SSW events


Case studies of four major SSWs have been performed and the semidiurnal solar and lunar tides of the EEJ have been determined. The subplots in the figures denote the following:

- Day-to-day variations of the EEJ (in nT) obtained from Huancayo and Fuquene observations
  - The amplitude of the semidiurnal solar (red line) and lunar (blue line) tide of the EEJ
  - Zonal mean zonal wind at 60° N and 10 hPa (red line) and the North Pole temperature at 10 hPa (black line)
  - Phase of the semidiurnal solar tide
  - Daily solar flux values
  - Phase of the semidiurnal lunar tide
- The open and filled circles at the bottom of the subplots denote the full moon and new moon, respectively.


### Comparison with numerical simulation results



### 2008-2009 SSW: WACCMX+DART



### 2012-2013 SSW: TIME-GCM nudged with WACCMX-L116/GEOS-5



## Summary

- The EEJ semidiurnal lunar and solar tides both show enhancements during the SSW events but the variability of the EEJ semidiurnal solar tide during the SSWs is more complex with enhancements occurring prior to the onset of SSWs followed by a reduction and another enhancement following the peak reversal of the zonal mean zonal wind at 60° N and 10 hPa.
- The relative amplification of the EEJ semidiurnal lunar tide is seen to be larger than the EEJ semidiurnal solar tide during all the four analyzed SSWs.
- The timing of global M2 enhancements in zonal wind at ~120 km altitude and the EEJ semidiurnal lunar tidal enhancements show a good agreement with each other. In case of a similar comparison between the SW2 and the EEJ semidiurnal solar tidal enhancements, the degree of agreement varies for each of the SSW events.

### Acknowledgments

We would like to thank the Instituto Geofísico del Peru and Instituto Geográfico Agustín Codazzi, Colombia for supporting geomagnetic observatory operations at Huancayo and Fuquene, respectively. The F10.7 data are obtained from NASA GSFC/SPDF OMNIWeb. The NCEP-NCAR reanalysis data are available at the NOAA/OAR/ESRL website. C.S and H.L are partly supported by SPP 1788 "Dynamic Earth" of the Deutsche Forschungsgemeinschaft (DFG). Y. Y was supported by the Humboldt Research Fellowship for Experienced Researchers from the Alexander von Humboldt Foundation. The National Center for Atmospheric Research is sponsored by National Science Foundation. T.A.S and A.M are supported by NASA grant X13AF77G.

### References

Pedatella, N., Chau, J., Schmidt, H., Goncharenko, L., Stolle, C., Hocke, K., Harvey, V., Funke, B., and Siddiqui, T.: How Sudden Stratospheric Warming Affects the Whole Atmosphere, *Eos*, 2018a.  
 Pedatella, N., Liu, H.-L., Marsh, D., Raeder, K., Anderson, J., Chau, J., Goncharenko, L., and Siddiqui, T.: Analysis and Hindcast Experiments of the 2009 Sudden Stratospheric Warming in WACCMX+ DART, *Journal of Geophysical Research: Space Physics*, 123, 30 3131–3153, 2018b.  
 Matsuno, T.: A dynamical model of the stratospheric sudden warming, *Journal of the Atmospheric Sciences*, 28, 1479–1494, 1971.  
 Maute, A., Hagan, M. E., Yudin, V., Liu, H.-L., and Yizengaw, E.: Causes of the longitudinal differences in the equatorial vertical ExB drift during the 2013 SSW period as simulated by the TIME-GCM, *Journal of Geophysical Research: Space Physics*, 120, 5117–5136, 2015.