

Spike development inhibition in the *ftin* mutant is associated with multiple phenotypic characteristics and regulated multiple biological pathways

Yong-sheng Zheng¹, Jinpeng Zhang¹, Cheng Liu², Rongzhi Zhang², Xiajie Ji¹, Han Zhang², Ruyu Li², and Weihua Liu¹

¹Chinese Academy of Agricultural Sciences

²Shandong Academy of Agricultural Sciences

April 28, 2020

Abstract

Spike development of wheat is closely associated with the ability of response to cold stress and inhibited under cold stress in spring. Morphological investigation showed that the *ftin* gene in 3558M is associated with multiple phenotypic characteristics, including fewer tillers, delayed floral transition, and the death of shoot apical meristem. In this study, we systematically researched the genetic nature of spike development using ITRAQ, transcriptome sequencing, western blot and RNAi technologies. The results showed that the *ftin* mutant is cold sensitive and activates the cold acclimation pathway. Multiple defence responses, including the ROS-mediated hypersensitive response, SA-mediated systemic acquired resistance and FLS2-induced pathogen-associated defence response, were activated to respond to normal cold stress and led to the apex death. Meanwhile, the continuous cold acclimation pathway inhibited the SVP-SCO1-LFY flowering pathway and leads to inhibition of spike development. Two TaPIN proteins were significantly downregulated, and multiple auxin signalling genes were also differentially expressed after cold stress. We used RNAi technology to knock down the two TaPIN genes and the tiller number was significantly reduced in TaPINa-RNAi and TaPINb-RNAi wheat. Taken together, the results revealed that the *ftin* gene might directly or indirectly regulate multiple biological pathways to affect multiple phenotypic characteristics.

Hosted file

Spike development inhibition in the *ftin* mutant is associated with multiple phenotypic characteristics available at <https://authorea.com/users/313917/articles/444324-spike-development-inhibition-in-the-ftin-mutant-is-associated-with-multiple-phenotypic-characteristics-and-regulated-multiple-biological-pathways>