

# Medial Thigh Fasciocutaneous Flaps for Reconstruction of the Scrotum following Fournier Gangrene: A case report

Rodrigue Mwenibamba<sup>1</sup>, Daniel Safari Nteranya<sup>2</sup>, Augustin Mukakala<sup>1</sup>, Christian Wabene<sup>1</sup>, Angelique Rugendabanga<sup>1</sup>, Fabrice Kibukila<sup>1</sup>, Nathan Mitima<sup>1</sup>, and Jean Claude Niyondiko<sup>3</sup>

<sup>1</sup>Official University of Bukavu

<sup>2</sup>Universite Officielle de Bukavu

<sup>3</sup>University of Burundi

February 5, 2022

## Abstract

Fournier's gangrene, a rapidly progressive necrotizing fasciitis of the perineum and external genitalia remains a medical-surgical emergency. The authors present a case of coverage of a scrotal loss of substance with testicular exposure by a medial fasciocutaneous flap in a fifty years male patient.

## Hosted file

Medial Thigh Fasciocutaneous Flaps for Reconstruction of the Scrotum following Fournier Gangrene.docx available at <https://authorea.com/users/459042/articles/555442-medial-thigh-fasciocutaneous-flaps-for-reconstruction-of-the-scrotum-following-fournier-gangrene-a-case-report>