

# A Case of Suspected Lobular Endocervical Glandular Hyperplasia in a Cervical Cystic Lesion During Pregnancy

Sayaka Suzuki<sup>1</sup>, Yoshinobu Sugo<sup>2</sup>, Kazuya Hiiragi<sup>1</sup>, Soichiro Obata<sup>3</sup>, Akiko Iwata<sup>4</sup>, Yuichi Imai<sup>5</sup>, Shigeru Aoki<sup>1</sup>, Kentaro Kurasawa<sup>4</sup>, and Etsuko Miyagi<sup>2</sup>

<sup>1</sup>Yokohama City University Medical Center

<sup>2</sup>Yokohama City University Hospital

<sup>3</sup>Yokohama Shiritsu Daigaku Fuzoku Shimin Sogo Iryo Center

<sup>4</sup>Yokohama City University Urafane Hospital

<sup>5</sup>Yokohama City University Graduate School of Medicine

May 24, 2022

## Abstract

Lobular endocervical glandular hyperplasia (LEGH) is characterized by clinically profuse and watery vaginal discharge. In pregnancy with LEGH, with watery fluid leakage persisting throughout pregnancy, it is often difficult to visually diagnose PROM. Adding to this difficulty, auxiliary diagnostic tests might also show positive results, complicating treatment and management.

## Hosted file

Sayaka\_Suzuki.docx available at <https://authorea.com/users/484653/articles/570338-a-case-of-suspected-lobular-endocervical-glandular-hyperplasia-in-a-cervical-cystic-lesion-during-pregnancy>



