

# A case of hemorrhagic shock in a patient with neurofibromatosis type 1

Yutaro Sakaguchi<sup>1</sup>, Hiromu Okano<sup>1</sup>, Ryosuke Furuya<sup>1</sup>, Hiroshi Miyazaki<sup>1</sup>, and Tsuyoshi Otsuka<sup>1</sup>

<sup>1</sup>National Hospital Organisation Yokohama Medical Center

July 20, 2022

## Abstract

Neurofibromatosis type 1 (NF1) is not known to cause serious vascular complications. This report describes a 49-year-old man with NF1 who presented to our hospital in hemorrhagic shock due to vascular leakage within a neurofibroma tumor. Physicians should be aware that NF1 can cause a wide variety of vascular lesions.

## Hosted file

Figure 1 Clinical image.pptx available at <https://authorea.com/users/496312/articles/577756-a-case-of-hemorrhagic-shock-in-a-patient-with-neurofibromatosis-type-1>

1 A case of hemorrhagic shock in a patient with neurofibromatosis type 1

2 Shock in neurofibromatosis type 1

3  
4 Yutaro Sakaguchi<sup>1,2</sup>, Hiromu Okano<sup>1,3</sup>, Ryosuke Furuya<sup>1,2</sup>, Tsuyoshi Otsuka<sup>1,2</sup>,  
5 Hiroshi Miyazaki<sup>1</sup>,

6  
7 <sup>1</sup>Department of Emergency and Critical Care Medicine, National Hospital Organization  
8 Yokohama Medical Center, 3-60-2 Harajuku, Totsuka-ku, Yokohama, Kanagawa 245-  
9 8575, Japan

10 <sup>2</sup>Department of Emergency Medicine, Graduate School of Medicine, Yokohama City  
11 University, 3-9 Fukuura, Kanazawa-ku, Yokohama, Kanagawa 236-0004 Japan.

12 <sup>3</sup>International University of Health and Welfare Graduate School of Public Health,  
13 4-1-26 Akasaka, Minato City, Tokyo 107-8402, Japan

14  
15 **Corresponding author**

16 Hiromu Okano,  
17 Department of Critical Care and Emergency Medicine,  
18 National Hospital Organization Yokohama Medical Center,  
19 3-60-2 Harajuku, Totsuka-ku, Yokohama-shi, Kanagawa 245-8575, Japan  
20 E-mail: okanohiromu0121@gmail.com

## **Abstract**

Neurofibromatosis type 1 (NF1) is not known to cause serious vascular complications. This report describes a 49-year-old man with NF1 who presented to our hospital in hemorrhagic shock due to vascular leakage within a neurofibroma tumor. Physicians should be aware that NF1 can cause a wide variety of vascular lesions.

## **Case presentation**

A 49-year-old man with neurofibromatosis type 1 (NF1) was brought to the emergency room for swelling and pain in the left side of his abdomen. He was in shock on arrival, and a large mass was observed in his lateral abdomen. Contrast-enhanced computed tomography (CT) was performed to identify the source of bleeding and search for other vascular lesions. It revealed vascular leakage within a neurofibroma tumor (Fig. 1A), leading to the observed hemorrhagic shock. An aneurysm of the left 4th lumbar artery was observed, but no cerebral aneurysms or other vascular abnormalities were found. Emergency interventional radiology (IVR) and vascular embolization of the left 1st, 3rd, and 4th lumbar arteries were performed (Fig. 1B). The aneurysm of the left 4th lumbar artery was treated conservatively.

NF1 is an autosomal-dominant inherited disease with an incidence of 1:3000.<sup>(1)</sup> The incidence of vascular lesions in patients with NF1 is reportedly 3.6%.<sup>2</sup> In this case, an aneurysm was found in the left 4th lumbar artery. Therefore, when emergency room physicians encounter shock patients with NF1, they should perform whole body contrast-enhanced CT to detect vascular lesions as well as other vascular malformations.

## **ACKNOWLEDGEMENT**

We would like to thank Editage (<http://www.editage.jp>) for English language editing.

## **CONFLICT OF INTEREST**

Authors declare no Conflict of Interests for this article.

## **AUTHOR CONTRIBUTIONS**

YS wrote the first draft of the manuscript. HO, TO, HM, and RF critically revised the manuscript. All authors read and approved the final version of the manuscript.

## **ETHICAL APPROVAL**

This article does not contain any studies involving human participants or animals.

## **INFORMED CONSENT**

Written consent was obtained from the patient to disclose the case details and images pertaining to diagnosis.

72

73 **References**

74 1. Evans DG, Howard E, Giblin C, et al. Birth incidence and prevalence of tumor-  
75 prone syndromes: estimates from a UK family genetic register service. *Am J Med Genet*  
76 *A*. 2010;152A(2):327-32.

77 2. Miura H, Taira O, Uchida O, Usuda J, Hirai S, Kato H. Spontaneous  
78 haemothorax associated with von Recklinghausen's disease: review of occurrence in  
79 Japan. *Thorax*. 1997;52(6):577-8; discussion 575-6.

80