

# Risk Factors of Sepsis and Prevalance of Multi-Drug Resistant Organisms in Pediatric Cardiac Surgery in Tertiary Care Teaching Rural Hospital In India: Retrospective, Observational Study Short running title: Risk Factors of Sepsis and Prevalance of Multi- Drug Resistant Organisms

Dr. Vishal V. Bhende<sup>1</sup>, Vishal V Bhende<sup>2</sup>, Sohilkhan R Pathan<sup>3</sup>, Tanishq S Sharma<sup>2</sup>, Amit Kumar<sup>4</sup>, Hardil P Majmudar<sup>2</sup>, and Vishal A Patel<sup>3</sup>

<sup>1</sup>Affiliation not available

<sup>2</sup>Department of Pediatric Cardiac Surgery, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University

<sup>3</sup>Department of Pediatric Cardiac Intensive Care, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University

<sup>4</sup>Clinical Research Services, Clinical Research Co-ordinator, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University

February 3, 2023

# Risk Factors of Sepsis and Prevalance of Multi-Drug Resistant Organisms in Pediatric Cardiac Surgery in Tertiary Care Teaching Rural Hospital In India: Retrospective, Observational Study

Short running title: Risk Factors of Sepsis and Prevelance of Multi-Drug Resistant Organisms

## Authors:

Vishal V. Bhende<sup>1</sup>, Sohilkhan R. Pathan<sup>2</sup>, Tanishq S. Sharma<sup>1</sup>, Amit Kumar<sup>3</sup>, Hardil P. Majmudar<sup>1</sup>, Vishal A. Patel<sup>2</sup>

<sup>1</sup>Department of Pediatric Cardiac Surgery, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University, Gokal Nagar, Karamsad, Anand, Gujarat-388 325, India.

<sup>2</sup>Department of Pediatric Cardiac Intensive Care, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University, Gokal Nagar, Karamsad, Anand, Gujarat - 388 325, India.

<sup>3</sup>Clinical Research Co-ordinator, Clinical Research Services, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University, Karamsad, Anand, Gujarat - 388 325, India.

**Corresponding Author:** Vishal V. Bhende Department of Pediatric Cardiac Surgery, Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Bhaikaka University, Karamsad, Anand, Gujarat - 388 325, India

Email: drvishalbhende@gmail.com; Phone: +91 98453 83405

**Author Contributions:**

**Data Curation:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar, Sohilkhan R. Pathan, Hardil P. Majmudar, Vishal A. Patel.

**Formal Analysis:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar, Sohilkhan R. Pathan, Hardil P. Majmudar, Vishal A. Patel.

**Funding Acquisition:** Vishal V. Bhende, Tanishq S. Sharma

**Investigation:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar, Sohilkhan R. Pathan, Hardil P. Majmudar

**Methodology:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar,  
Sohilkhan R. Pathan, Hardil P. Majmudar

**Project Administration:** Vishal V. Bhende, Tanishq S. Sharma, Amit  
Kumar, Sohilkhan R. Pathan, Hardil P.  
Majmudar, Vishal A. Patel.

**Resources:** Vishal V. Bhende, Tanishq S. Sharma, Sohilkhan  
R. Pathan, Hardil P. Majmudar, Vishal A. Patel.

**Software:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar,  
Sohilkhan R. Pathan, Hardil P. Majmudar

**Supervision:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar,  
Sohilkhan R. Pathan, Hardil P. Majmudar, Vishal A.  
Patel.

**Validation:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar,  
Sohilkhan R. Pathan, Hardil P. Majmudar, Vishal A.  
Patel.

**Visualization:** Vishal V. Bhende, Tanishq S. Sharma, Amit Kumar,  
Sohilkhan R. Pathan, Hardil P. Majmudar

**Writing-Original Draft Preparation:** Vishal V. Bhende,  
Tanishq S. Sharma

**Writing - Review & Editing:** Vishal V. Bhende, Tanishq S. Sharma

### **Acknowledgments**

We are thankful to -

- Mr. Mayur KS (Bio-statistician, Central Research Services, Bhaikaka University, Karamsad, India),

### **Funding statement:**

All authors have declared that no financial support was received from any organization for the submitted work.

### **Conflict of interest statement:**

All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. All authors have

declared that there are no other relationships or activities that could appear to have influenced the submitted work.

**Transparency statement:**

The corresponding author confirms that manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

**Data availability statement :**

The data that support the findings of this study are openly available in [Authorea] at  
, reference number [reference number].

**Informed consent statement :**

Waived off by IEC as it was a retrospective observational study.

## **Abstract**

### **Background and Aims**

Cardiac surgery and cardiopulmonary bypass lead to an immunoparalyzed state in children which makes them susceptible to sepsis and other hospital acquired infections. Identification of risk factors of sepsis would lead to appropriate management

### **Objective**

The present study was conducted to assess the prevalence of sepsis and risk factors associated with sepsis in pediatric cardiac surgical patients and the subsequent prevalence of multidrug resistant organisms.

### **Methods**

A retrospective, single-center observational study was conducted on 100 pediatric patients admitted to the PICU after cardiac surgery between January, 2017 and February, 2018. All the data of the patients were collected from the medical record department of the hospital. Patient case report form consisted of demography, surgery details, Pre-operative & post-operative haematological reports and clinical details. After collecting the data, chi-square test and logistic regression analysis were used to find the risk factors associated with sepsis.

## Results

The prevalence of sepsis in our population was 27 % and mortality rate due to sepsis was 1 %. The only statistically significant risk factor for sepsis we found in this study was prolonged ICU (intensive care unit) stay for more than 5 days. Total 8 patients had blood culture positive for bacterial infection. The alarming finding was that all 8 were multi-drug resistant organisms, requiring the last line of antibacterials.

## Conclusion

Our study implies that special clinical care is required when ICU stay is prolonged to mitigate the risk of sepsis. These new and upcoming infections not only lead to high mortality and morbidity rates but also contribute to increased cost of care due to the use of newer broad-spectrum antibiotics and longer hospital stay. The high prevalence of multi-drug resistant organisms is unacceptable in the current scenario and hospital infection and prevention control play a critical role in minimizing such infections.

*This abstract was presented at European Academy of Paediatrics, EAP 2021 Congress and MasterCourse 22-25 April 2021 (frontiers Event Abstracts)*

**Key Words:** sepsis, pediatric cardiac surgery



## Introduction

The survival rate for neonates, infants, and children with congenital heart disease (CHD) has increased significantly with improved surgical techniques, cardiopulmonary bypass (CPB) and post-operative care. However, CPB elicits a complex, systemic inflammatory response that is characterized by activation of the complement cascade, release of endotoxin, activation of leukocytes and release of pro-inflammatory cytokines. This complex humoral and cellular-mediated immune response results in a transient and relative state of immune suppression, often referred to as “immunoparalysis”.<sup>1,2</sup> This state of immunoparalysis may result in an increased risk of sepsis in children undergoing cardiac surgery.<sup>3,4</sup> In addition, chronic hypoxia, low cardiac output syndrome and other co-morbid conditions associated with CHD,<sup>3-5</sup> as well as invasive devices may also increase the risk of sepsis in this population.

Despite wide spread adoption of healthcare guidelines & sepsis prevention bundles, the frequency of healthcare acquired infection remains high, at 6 % to 30.8 % in pediatric cardiac surgical patients. Sepsis is a significant and independent risk factor for increased duration of mechanical ventilation, cardiac intensive care unit length of stay, healthcare costs, and mortality in children with CHD.

Very few case series of sepsis prevalence in pediatric cardiac surgical patients, in a rural tertiary care hospital from India has been published. None of them has highlighted the lurking threat of increasing prevalence of multi-drug resistant organisms.

## **Materials & Methods**

### **Study Design and Setting:**

This was a retrospective, single-center observational study conducted at Cardiac Pediatric Intensive Care Unit(CPICU) of Bhanubhai and Madhuben Patel Cardiac Centre, Shree Krishna Hospital, Karamsad, India after getting approval of the research project from the Institutional Ethics Committee (IEC-2) of the Bhanubhai and Madhuben Patel Centre for Medical Care and Education, Anand, Gujarat, India, vide approval No. (IEC/BU/2021/Ex.03/44/2023 dated 24.01.2023

### **Inclusion Criteria -**

- Consecutive Pediatric Patients less than or equal to 14 years old admitted to the CPICU after pediatric cardiac surgery between January, 2017 and February, 2018.

### **Exclusion Criteria -**

- Older patient & patient undergoing any major non-cardiac surgical intervention was excluded from surgery.
- Also patients discharged against medical advice, having pre-admission infection or patient transferred from other hospital department or other center with infection were excluded from the study.

## **Data collection**

The study was initiated after receiving approval from the Institutional Ethics Committee of the hospital. It was a retrospective observational study so in need of Informed Consent form was waived off by Institutional Ethics Committee.

An observational retrospective study was conducted on one hundred pediatric patients (less than 14 years) who had undergone cardiac surgery between January, 2017 & February, 2018. All the data of the patients were received from the medical record department of the hospital. Case report form was prepared in English.

To look for the associated risk factors for sepsis patients clinical and laboratory analyses were recorded in case report form. Such data are patient's demographic data including sex, age, weight, height, cardiac and non-cardiac abnormalities, major medical history. Preoperative sepsis screen & risk factors of immune-suppression like severe malnutrition, chromosomal disorders, HIV infection and use of pre-operative antibiotics & hospital stay was recorded.

Surgery data recorded were - surgery type, Risk adjustment for congenital heart surgery (RACHS) score for evaluating surgical complexity, CPB duration, hypothermia or total circulatory arrest, delayed sternal closure, intubation & ventilation duration, need of re-intubation, central catheter lines & its duration, length of ICU (intensive care unit) stay, blood transfusion, inotropic support, renal support, tracheostomy.

Haematology data included were preoperative & postoperative haemoglobin level, leukocytes count, c-reactive protein, procalcitonin & culture report were recorded from the medical records of the

patients. In case of positive culture report, antibiotics sensitivity pattern & their minimum inhibitory concentration were recorded. Mortality with its cause were also noted.

Definition of sepsis, septic shock was confirmed as per definition mentioned in Sepsis-3. Suspected sepsis was labelled when a patient was managed as sepsis, based on clinical & laboratory parameters.

## **Statistical Analysis**

Variables were evaluated using the chi-square test. Logistic regression analysis were used to find the risk factors associated with sepsis.

## **Results and Discussion**

One hundred and three pediatric patients with age less than 14 years, had undergone cardiac surgery during January, 2017 to February, 2018. Three patients who was discharged against medical advice or transferred to other centers were excluded from study

One hundred (100) pediatric cardiac surgical patients, aged less than 14 years old, were included in study and were observed retrospectively for the incidence of sepsis after cardiac surgery.

Out of 100 pediatric patients 55 patients were male and 45 patients were female.

**Table 1 - Demographic and Surgical Characteristics**

Characteristics	Mean(SD)	SD	Median	Minimum	Maximum
Age (year)	2.39	3.29	0.9	1 (day)	14 (year)
Weight (kgs.)	8.14	5.72	5.93	1	28
Height (cms.)	78.02	25.34	70	43	144
RACHS	1.85	0.74	2	1	4
Surgery duration (minute)	89.43	39.69	83	28	230
Duration of intubation(hours)	40.62	51.07	24	1	336
Pre-operative length of stay (days)	1.93	3.10	1	0	28
Total no. of hospital stay (days)	9.58	5.61	8	3	35
Length of ICU stay (days)	3.91	4.46	2	0	26

Out of 100 patients weight of 58 patients was more than 5 kgs and 36 patients were in-between 2.5 to 5 kgs while only 6 patients had weight less than 2.5 kgs.

8 patients were in the age group of < 30 days (neonates), 44 patients were in the age group of 30 days-1 year (infants), 30 patients were in the age group of 1 year-5 years while 18 patients were in the age group of 5 years-14 years.

In the study group 8 patients had blood culture proven sepsis. Another 8 patients had suspected sepsis which developed after surgery, while

11 had suspected sepsis developed pre-operatively during hospital stay.

**Table - 2 Infection and Mortality**

Infection	N=100	%
Proven Sepsis	8	8
Suspected Sepsis	19	19
No Infection	73	73
Mortality	-	2

**Table - 3 Mortality in the Study Group**

Mortality	Sepsis	RACHS	Cause of death
Patient 1	Yes	4	Acute respiratory distress syndrome with Sepsis
Patient 2	No	3	Cardiogenic shock

In our study group there was no case of mediastinitis or endocarditis. Although 6 patients had sternal wound gaping. Swab culture from these patients was sterile. They improved on oral antibiotics.

Among type of surgery - VSD (ventricular septal defect) closure surgery had higher prevalence. While total 14 patients had VSD closure, 4 (28%) of them had sepsis.

**Table - 4 Clinical Characteristics of the Patient Population and Risk Factors for Sepsis in Bivariate Analysis (Chi-Square Statistics)**

Variable	Level	Total (N=100)	Overall %	Number with sepsis (n=27)	% with sepsis	Number with no sepsis (n=73)	% with no sepsis	p Value
Pre-operative assessment								
RACHS category	1	34	34	5	18.52	29	39.73	0.236
	2	49	49	16	59.26	33	45.21	
	3	15	15	5	18.52	10	13.70	
	4	2	2	1	3.70	1	1.37	
	5	0	0	0	0	0	0	
	6	0	0	0	0	0	0	
Demographics								
Age	<30 days	8	8	5	18.52	3	4.11	*0.005
	30 days-1 year	44	44	16	59.26	28	38.26	
	1 year- 5 year	30	30	5	18.52	25	34.25	
	5 year- 14 year	18	18	1	3.70	17	23.29	
Weight	<2.5 kg	6	6	4	14.81	2	2.74	*0.004
	2.5-5 kg	36	36	14	51.85	22	30.14	
	>5 kg	58	58	9	33.33	49	67.12	
Sex	Male	55	55	15	55.56	40	54.79	0.946
	Female	45	45	12	44.44	33	45.21	
Risk factors								
Preoperati ve length of stay (days)	0	9	9	5	18.52	4	5.48	0.156
	1	60	60	13	48.15	47	64.38	
	2	14	14	5	18.52	9	12.33	
	>3	17	17	4	14.81	13	17.81	
Surgery	<30	1	1	1	3.70	0	0	



duration (minute)	30-90	62	62	8	29.63	54	73.97	<b>*0.000</b>
	90-200	35	35	18	66.67	17	23.29	
	>200	2	2	0	0	2	2.74	
Length of ICU stay (days)	1	34	34	1	3.70	33	45.83	<b>*0.000</b>
	2-4	37	37	9	33.33	28	38.89	
	5-9	19	19	10	37.04	9	12.50	
	10-30	9	9	7	25.93	2	2.78	
Duration of intubation (hours)	3	21	21	1	3.70	20	27.40	<b>*0.001</b>
	8	9	9	1	3.70	8	10.96	
	9-120	64	64	20	74.07	44	60.27	
	>120	6	6	5	18.52	1	1.37	
Duration of central venous catheteriz ation (hours)	3	21	21	1	3.70	20	27.40	<b>*0.001</b>
	8	9	9	1	3.70	8	10.96	
	9-120	64	64	20	74.07	44	60.27	
	>120	6	6	5	18.52	1	1.37	
Re- intubation	Yes	10	10	4	14.21	6	8.22	0.329
	No	90	90	23	85.19	67	91.78	
Inotropic Support	Yes	63	63	21	77.78	42	58.33	0.073
	No	36	36	6	22.22	30	41.67	
Delayed sternal closure	Yes	7	7	6	22.22	1	1.37	<b>*0.001</b>
	No	93	93	21	77.78	72	98.63	
Hypother mia	Yes	47	47	17	62.96	30	41.10	0.052
	NO	53	53	10	37.04	43	58.90	

As we inserted risk factors data in chi-square statistics we got 7 significant p values which are < 0.05. Such as age (0.005), weight (0.004), surgery duration (0.000), length of ICU stay (0.000), duration of intubation (0.001), duration of central venous catheterization (0.001) and delayed sternal closure (0.001) respectively.

**Table - 5 Risk Factors for Sepsis in Multivariable Analysis  
(Logistic regression analysis)**

Variable	Level	Number with sepsis (n=27)	% with sepsis	Number with no sepsis (n=73)	% with no sepsis	OR	(95%CI)	P-value
Age	<30days- 1 year	16	59.26	28	38.26	1.18	[0.14,9.70]	0.87
	1 year- 5 year	5	18.52	25	34.25	0.82	[0.05,12.5]	0.89
	5 year- 14 year	1	3.70	17	23.29	0.40	[0.01,11.3]	0.59
Weight	2.5-5 kg	14	51.85	22	30.14	0.66	[0.07,5.5]	0.70
	>5 kg	9	33.33	49	67.12	0.68	[0.06,7.3]	0.75
Length of ICU stay (days)	2-4	9	33.33	28	38.89	9.03	[0.82,98.8]	0.07
	5-9	10	37.04	9	12.50	19.0	[1.5,226.8]	<b>*0.02</b>
	10-30	7	25.93	2	2.78	50.4	[2.56,994.]	<b>*0.01</b>
Duration of intubation (hours)	8	1	3.70	8	10.96	1.24	[0.05,28.9]	0.89
	9-120	20	74.07	44	60.27	4.45	[0.46,42.9]	0.19
	>120	5	18.52	1	1.37	9.52	[0.29,311.]	0.20
Delayed sterna closure	Yes	17	62.96	30	41.10	0.36	[0.003,43.]	0.68
	No	10	37.04	43	58.90	0.3	[0.001,1.0]	0.054

Multivariate logistic regression analysis was performed to analyze the risk factors of sepsis. The analysis concluded that the risk factors were associated only with the length of ICU stay. Prolonged Duration of ICU (intensive care unit) stay more than 5 days (<0.05) which is moderately significant.

Total 8 patients had blood culture positive for bacterial infection - mostly were nosocomial gram negative bacteria. One was *Enterococcus faecium*. Alarming finding is that except one, all other are multi-drug

resistant organism requiring last line of anti-bacterials as shown in chart.

## Discussion

Infections in children are frequent (incidence, 13% to 31%) after cardiac surgery <sup>3,6</sup>. Many are surgical site infections (incidence, 2.3 % to 8 %) <sup>7</sup>. Some are more serious, such as septicemia (incidence, 6.3 % to 15 %) <sup>8,9,10</sup>, mediastinitis (incidence, 0.2 % to 3.3 %) <sup>6-14</sup>, and endocarditis (incidence, 0.2 %) <sup>11</sup>.

Pollock and associates <sup>4</sup> reported blood-stream infections (BSIs) in 21 (6.8 %) of 310 children after cardiac surgery. In our study, prevalence of sepsis was 27 % with 1 % mortality. There was no incidence of mediastinitis or endocarditis in the study group. 6% patients had surgical site infection.

Dicky Fakhri et al. <sup>12</sup> reported in their study that duration of cardiopulmonary bypass  $\geq 90$  minutes was associated with 5.538 increased risk of post-surgical sepsis in comparison to those  $\leq 90$  minutes. Our study was insufficiently powered to prove it, although trends suggested it.

Rosanova et al. <sup>13</sup> concluded that longer hospitalization and inotropic support were risk factors for infections in their study.

Takeshi Hatachi et al. <sup>14</sup> concluded that mechanical ventilation greater than or equal to

3 days, dopamine use, genetic abnormality, and delayed sternal closure were associated with healthcare-associated infections after pediatric cardiac surgery.

Risk factor which was significant in our study was longer duration of ICU (intensive care unit) stay (more than 5 days). Multi-variate analysis of our study group did not find other independent risk factors that may be because of inadequate numbers to have statistical significance.

However, the alarming finding of our study is very high prevalence of multi-drug resistant nosocomial bacterial infection (7 out of 8 culture positive sepsis). They required last line of anti-bacterials. In spite of this, mortality rate due to sepsis was only 1 %. However, this may worsen if we lose sensitivity to last line of anti-bacterials. This scenario is unacceptable in pediatric cardiac patients who may encounter precarious hemodynamics due to cardiac surgery.

Other studies have also highlighted prevalence of resistant organism in pediatric cardiac septic patients. This reflects deficiencies in anti-sepsis measures in many pediatric cardiac department.

## **Conclusion**

The prevalence of sepsis in the study population was 27 % which is comparable with other studies. With implementation of proper asepsis guideline the prevalence of sepsis can be reduced further to 5% as in recent study.

There is high incidence of multi drug resistance organisms in culture positive septic patients. If urgent steps are not taken to contain it, we may lose many patients because of sepsis with resistant bacteria. Misuse & overuse of antibiotics may further aggravate the problem

Prolonged ICU (intensive care unit) stay was statistically significant as a risk factor for sepsis, in current study. It implies that special clinical

care is required when ICU stay is prolonged to mitigate the risk of sepsis.

### **Limitations of the Study**

As it is a retrospective study some data discrepancy cannot be ruled out. Although care has been taken to minimize it. Many other risk factors were not statistically significant as the study group was not sufficiently powered in this study population. As it is a single center study, its result cannot be generalized for entire pediatric cardiac surgery patients in the region or country.

I am the primary author, submitting author and corresponding author. I have provided ORCID IDs of all the co-authors including me.

- Vishal V. Bhende <https://orcid.org/0000-0003-3014-4315>
- Sohilkhan R. Pathan <https://orcid.org/0000-0002-5381-9621>
- Tanishq S. Sharma <https://orcid.org/0000-0002-7135-8160>
- Amit Kumar <https://orcid.org/0000-0001-8950-4448>
- Hardil P. Majmudar <https://orcid.org/0000-0003-2414-5602>
- Vishal A. Patel <https://orcid.org/0000-0002-6740-6764>

## References

## References

1. Tarnok A, Schneider P. Pediatric cardiac surgery with cardiopulmonary bypass: pathways contributing to transient systemic immune suppression. *Shock*. 2001;16 (Suppl 1):24-32. doi: 10.1097/00024382-200116001-00006. PMID: 11770029.
2. Allen ML, Peters MJ, Goldman A, et al. Early postoperative monocyte deactivation predicts systemic inflammation and prolonged stay in pediatric cardiac intensive care. *Crit Care Med*. 2002;30:1140-1145. doi: 10.1097/00003246-200205000-00031. PMID: 12006816.
3. Levy I, Ovadia B, Erez E, et al. Nosocomial infections after cardiac surgery in infants and children: incidence and risk factors. *J Hosp Infect*. 2003;53:111-116. doi: 10.1053/jhin.2002.1359. PMID: 12586569.
4. Pollock EM, Ford-Jones EL, Rebeyka I, et al. Early noscomial infections in pediatric cardiovascular surgery patients. *Crit Care Med*. 1990;18:378-384. doi: 10.1097/00003246-199004000-00006. PMID: 2318048.
5. Wheeler DS, Dent CL, Manning PB, et al. Factors prolonging length of stay in the cardiac intensive care unit following the arterial switch operation. *Cardiol Young*. 2008;18:41-50. doi: 10.1017/S1047951107001746. Epub 2007 Dec 20. PMID: 18093360; PMCID: PMC2757101.
6. Guardia Cami MT, Jordan Garcia I, Urrea Ayala M. Nosocomial infections in pediatric patients following cardiac surgery. *An*

- Pediatr (Barc) 2008;69(1):34-8. doi: 10.1157/13124216. PMID: 18620674.
7. Napolitano LM., Sepsis 2018: Definitions and Guideline Changes. Surg Infect (Larchmt). 2018 Feb/Mar;19(2):117-125. doi: 10.1089/sur.2017.278. PMID: 29447109.
  8. Urrea M, Pons M, Serra M, Prospective incidence study of nosocomial infections in a pediatric intensive care unit. Pediatr Infect Dis J. 2003 Jun;22(6):490-4. doi: 10.1097/01.inf.0000069758.00079.d3. PMID: 12799503.
  9. Mrowczynski W, Wojtalik M, Zawadzka D, . Infection risk factors in pediatric cardiac surgery. Asian Cardiovasc Thorac Ann. 2002 Dec;10(4):329-33. doi: 10.1177/021849230201000411. PMID: 12538279.
  10. Shah SS, Kagen J, Lautenbach E, . Bloodstream infections after median sternotomy at a children's hospital. J Thorac Cardiovasc Surg. 2007 Feb;133(2):435-40. doi: 10.1016/j.jtcvs.2006.09.026. PMID: 17258580.
  11. Hatachi T, Tachibana K, Takeuchi M. Risk Factors for Healthcare-Associated Infections After Pediatric Cardiac Surgery. Pediatr Crit Care Med. 2018 Mar;19(3):237-244. doi: 10.1097/PCC.0000000000001445. PMID: 29319633; PMCID: PMC5841862.
  12. Fakhri, Dicky, Pribadi Wiranda Busro, Budi Rahmat, "Risk factors of sepsis after open congenital cardiac surgery in infants: a pilot study." *Medical Journal of Indonesia* 25 (2016): 182-189.DOI: <https://doi.org/10.13181/mji.v25i3.1450>

13. Rosanova MT, Allaria A, Santillan A,.Risk factors for infection after cardiovascular surgery in children in Argentina. Braz J Infect Dis. 2009 Dec;13(6):414-6. doi: 10.1590/s1413-86702009000600005. PMID: 20464331.
14. [Takeshi Hatachi](#), Kazuya Tachibana,[Yu Inata](#) et al. Risk Factors for Healthcare-Associated Infections After Pediatric Cardiac Surgery, [Pediatr Crit Care Med](#). 2018 Mar; 19(3): 237-244. doi: [10.1097/PCC.0000000000001445](#)



0. Napolitano LM., Sepsis 2018: Definitions and Guideline Changes. *Surg Infect (Larchmt)*. 2018 Feb/Mar;19(2):117-125. doi: 10.1089/sur.2017.278. PMID: 29447109.
1. Fakhri, Dicky, Pribadi Wiranda Busro, Budi Rahmat, “Risk factors of sepsis after open congenital cardiac surgery in infants: a pilot study.” *Medical Journal of Indonesia* 25 (2016): 182-189. DOI: <https://doi.org/10.13181/mji.v25i3.1450>
  2. Guardia Camí MT, Jordan García I, Urrea Ayala M. Infección nosocomial en postoperados de cirugía cardíaca [Nosocomial infections in pediatric patients following cardiac surgery]. *An Pediatr (Barc)*. 2008 Jul;69(1):34-8. Spanish. doi: 10.1157/13124216. PMID: 18620674.
  3. Urrea M, Pons M, Serra M, Prospective incidence study of nosocomial infections in a pediatric intensive care unit. *Pediatr Infect Dis J*. 2003 Jun;22(6):490-4. doi: 10.1097/01.inf.0000069758.00079.d3. PMID: 12799503.
  4. Valera M, Scolfaro C, Cappello N, Nosocomial infections in pediatric cardiac surgery, Italy. *Infect Control Hosp Epidemiol*. 2001 Dec;22(12):771-5. doi: 10.1086/501861. PMID: 11876456.
  5. Pollock EM, Ford-Jones EL, Rebeyka I, et al. Early nosocomial infections in pediatric cardiovascular surgery patients. *Crit Care Med*. 1990 Apr;18(4):378-84. doi: 10.1097/00003246-199004000-00006. PMID: 2318048.
  6. Mehta PA, Cunningham CK, Colella CB, Risk factors for sternal wound and other infections in pediatric cardiac surgery

- patients. *Pediatr Infect Dis J*. 2000 Oct;19(10):1000-4. doi: 10.1097/00006454-200010000-00012. PMID: 11055604.
7. Mrowczynski W, Wojtalik M, Zawadzka D, . Infection risk factors in pediatric cardiac surgery. *Asian Cardiovasc Thorac Ann*. 2002 Dec;10(4):329-33. doi: 10.1177/021849230201000411. PMID: 12538279.
  8. Shah SS, Kagen J, Lautenbach E, . Bloodstream infections after median sternotomy at a children's hospital. *J Thorac Cardiovasc Surg*. 2007 Feb;133(2):435-40. doi: 10.1016/j.jtcvs.2006.09.026. PMID: 17258580.
  9. Allpress AL, Rosenthal GL, Goodrich KM,.Risk factors for surgical site infections after pediatric cardiovascular surgery. *Pediatr Infect Dis J*. 2004 Mar;23(3):231-4. doi: 10.1097/[01.inf.0000114904.21616.ba](https://doi.org/10.1097/01.inf.0000114904.21616.ba). PMID: 15014298.
  10. Di Filippo S, Delahaye F, Semiond B, .Current patterns of infective endocarditis in congenital heart disease. *Heart*. 2006 Oct;92(10):1490-5. doi: 10.1136/hrt.2005.085332. Epub 2006 Jul 3. PMID: 16818488; PMCID: PMC1861050.
  11. Sen AC, Morrow DF, Balachandran R,. Postoperative Infection in Developing World Congenital Heart Surgery Programs: Data From the International Quality Improvement Collaborative. *Circ Cardiovasc Qual Outcomes*. 2017 Apr;10(4):e002935. doi: 10.1161/CIRCOUTCOMES.116.002935. PMID: 28408715.
  12. Rosanova MT, Allaria A, Santillan A,.Risk factors for infection after cardiovascular surgery in children in Argentina. *Braz J*

- Infect Dis. 2009 Dec;13(6):414-6. doi: 10.1590/s1413-86702009000600005. PMID: 20464331.
13. Wheeler DS, Jeffries HE, Zimmerman JJ, . Sepsis in the pediatric cardiac intensive care unit. *World J Pediatr Congenit Heart Surg.* 2011 Jul 1;2(3):393-9. doi: 10.1177/2150135111403781. PMID: 22337571; PMCID: PMC3277844.
  14. Hatachi T, Tachibana K, Takeuchi M. Risk Factors for Healthcare-Associated Infections After Pediatric Cardiac Surgery. *Pediatr Crit Care Med.* 2018 Mar;19(3):237-244. doi: 10.1097/PCC.0000000000001445. PMID: 29319633; PMCID: PMC5841862.
  15. Willyard C. The drug-resistant bacteria that pose the greatest health threats. *Nature.* 2017 Feb 28;543(7643):15. doi: 10.1038/nature.2017.21550. PMID: 28252092.