

Monoallelic deleterious MUTYH mutations generate colorectal cancer:a case report

Bei Zhao¹, Wenqi Sun¹, Yunrong Wang¹, Xinrong Wu¹, Yifan Li¹, Weiwei Wang¹, Muhan Ni¹, Peng Yan¹, Xiaotan Dou¹, Lei Wang¹, and Min Chen¹

¹Nanjing Drum Tower Hospital, Affiliated Hospital of Medical School, Nanjing University, Nanjing, China

July 11, 2023

Monoallelic deleterious MUTYH mutations generate colorectal cancer:a case report

Abstract

Background: Here we reported a particular case of MUTYH-associated polyposis (MAP) that had only one rare heterozygous mutation, but some particular clinical manifestations contributed to occur in this male patient by only one defective MUTYH allele were worth of further investigation.

Case presentation: We reported a case of MAP. It is about a 33-year-old man with chief complaints of hematochezia who had multiple polyps which were found in his colon via colonoscopy. He followed his doctor's advice and performed a genetic analysis examination. Germline test was positive for a major heterozygous mutation: chr1:45800165 on the MUTYH gene. MUTYH gene sequence analysis confirmed the following heterozygous mutation: c.55CT (p.R19X) in exon 2 (ClinVar NM.001128425). Unfortunately, his mother and daughter have the ilk mutation according to genetic analysis (Figure 1). However, this mutation at the site was not detected in his father. Various types of polyps were found on repeated colonoscopy, which tended to become latent cancerous in the futre.

Conclusion: This case indicated that awareness of the risk of carcinogenesis of polyps in carriers of monoallelic mutations might accordingly increase and understanding of the type of genetic-related disease will be enhanced by us.

keywords: MUTYH; Colorectal cancer; Hereditary colorectal cancer; Polyposis

Background

Familial adenomatous polyposis (FAP), a rare autosomal dominant inherited disease, is generally considered to be related to the APC gene^[1]. However, there is a special type of APC-negative but MUTYH-positive called MUTYH-associated polyposis (MAP). MAP includes both monoallelic mutations and biallelic mutations^[2]. Of course, biallelic mutations lead to an increased risk of colorectal cancer. But patients with biallelic mutations are a minority, and heterozygous variants are more common. It is still unclear whether only one MUTYH mutation is of clinical importance or not. Recently, more and more research has focused on the risk of colorectal cancer caused by monoallelic mutations^[3]. Previously, a number of studies showed that monoallelic MUTYH had little to do with CRC. Whether heterozygous variants increase the risk of colorectal cancer is still under debate^[4]. Interestingly, what happens in the gut deserves further investigation if there is another mutation associated with colorectal polyps. In this report, we described the rare genetic variant, which were unusual genetic mutations that improve our awareness and understanding of the disease.

Case description

A 33-year-old man was taken for counseling after discovering genetic mutations. Firstly, on March 21st, 2022, the patient went to Anhui Provincial Hospital for treatment with chief complaint of hematochezia for several months. He then underwent a colonoscopy which was showed that a 3.5 cm long pedunculated polyp could be seen in the descending colon at a distance of 55 cm from the edge of the anus. And then a 0.8 cm size polyp was detected near to it. In addition, another larger polyp, about 6 cm in size, was found at the junction of the sigmoid and descending colon. It was worth noting that this distinct sessile polyp with a lobulated surface and varicose veins of the surrounding mucosa almost blocked the intestinal lumen. Several polyps ranging in size from 0.5 cm to 1 cm also were found in the rest of the colon. Pathologic findings after biopsy confirmed that the maximal polyp is a tubulovillous adenoma with moderate dysplasia. A computed tomography of the abdomen, performed half a month later, documented multiple lesions in the splenic flexion of the colon and in the sigmoid, considering the larger one, which is local with intussusception proximal to the sigmoid. Genetic analysis revealed a deleterious single allele heterozygous germline mutation. MUTYH gene sequence analysis confirmed the following heterozygous mutation: c.55CT (p.R19X) in exon 2 (ClinVar NM.001128425). In addition, the LHX4 mutation: c.256G>A (p.G86S) in exon 3 (ClinVar NM.033343) also was notable. The electrophoretic sequence of MUTYH exon 2 revealing mutations is shown in Figure 2. This genetic analysis confirmed that the MUTYH mutation was a single allele heterozygous mutation. A diagnosis of MAP was then confirmed. His mother and daughter have the same MUTYH gene mutation, however, while no particular genetic variations occurred in his father. Interestingly, this MUTYH mutation carrier had a family history consistent with dominant inheritance. When he was admitted to another Class A tertiary hospital on May 2nd, 2022, he underwent endoscopic mucosal resection surgery. Pathological analysis showed that cystic fibrosis located 25 cm from the margin of anus was a microvillous tubular adenoma with high grade intraepithelial neoplasia and local carcinogenesis. Specifically, cancerous tissue at the site has infiltrated into the 3mm submucosal layer, but the vertical and lateral margins were both negative. The surgeon recommended abdominal surgery for the patient, but failed. Almost about 22 days later, the patient sought further endoscopic treatment and came to our hospital. An abdominal enhancement CT was later performed, documenting multiple mass lesions of the colonic splenic curvature and sigmoid colon. The next day the patient underwent colorectal EMR and ESD (Figure 3). Lesions between 5 mm and 6 cm are found during surgery. The largest tissue sample with dissociative tissue pieces was a villous tubular adenoma consisting of large low-grade and rarely high-grade intraepithelial neoplasia. In addition, a lesion grows nearby, the pathology of which was a villous tubular adenoma with low-grade intraepithelial neoplasia. The splenic flexure and sigmoid also have several similar polyps. Some hyperplastic polyps were detected in the rectum. All pathologically confirmed lesion strains were negative. Immunohistochemistry analysis was added showing that the junction of the descending colon with the sigmoid, sigmoid and splenic flexure showed similar results: all polyps were sent for investigation manifested as Desmin (mucosal muscle +), CEA (+), P53 (+), including wild-type and partially mutant and various grades of Ki67(+).

Three months ago, he came to our hospital for colonoscopy surveillance. And the positron emission tomography (PET-CT) scan was unremarkably normal. When he was hospitalized this time, his bowel preparation was not fairly satisfactory as a large amount of feces remained in the bowel(Figure 4). Many polyps were found again after the colonoscopy. Tubular adenomas and hyperplastic polyps were removed with the technique of a cold snare. Considering the poor bowel preparation, he was instructed to repeat the colonoscopy after 3 months. At present, he went to our hospital again for a colonoscopy. During this process, several newly emerged polyps were found in the colon and were removed via conventional EMR. These polyps remain concentrated in the distal colon, including the descending colon, sigmoid colon, and rectum(Figure 5). In order to avoid unnecessary risks, we also arranged an additional CT examination for him, and this time there were no signs of intussusception and other potential malignant changes.

Discussion

According to relevant research, the particular genetic mutation has been reported only in Asian regions, in contrast to MUTYH-related common gene mutations in Caucasians [5-8]. MUTYH-associated familial intestinal polyposis caused by this gene also is attenuated compared to regular MAP. MAP occurred mainly in the proximal colon, and polyps are mainly tubular adenomas, some tubular villous adenomas, and occa-

sionally hyperplastic polyps^[9]. Interestingly, the vast majority of polyps in this patient were located in the left colon, in contrast to MAP, which occurred in the right colon. However, recent literature has shown that patients associated with MYH deficiency resulting in hyperplastic polyps are not uncommon and present in a condition where hyperplastic polyps coexisted with adenomatous polyps^[9-10]. This is consistent with the pathologic features of polyps in this patient, with proliferative polyps, tubular adenomas, and choriomatous adenomas growing together.

He has a heterozygous mutation in the MUTYH gene, which is a monoallelic mutation. The clinical severity is low, however, caused by this monoallelic heterozygous mutation was not yet reported in any relevant case. Unlike biallelic mutations, which could accelerate the development of adenomas in subsequent years, according to previous literature and ESGE guidelines, patients with uniallelic mutations rarely developed adenomatous polyposis or had only a slightly increased risk of developing CRC^[4, 11]. As mild as the description in the literature is, his immunohistochemical results predicted that the tumor had a strong invasive and metastatic capacity. This might indicate the need for closer follow-up care. As understanding increases, there also is a growing body of literature showing that even patients with single allelic mutations have a higher risk of cancer than other normal people^[12, 13, 15]. Tumorigenesis by a mechanism of functional heterozygous somatic deletion of the MUTYH allele in tumors due to monoallelic pathogenic MUTYH germline variants. And the carriers of the monoallelic pathogenic germline variant MUTYH have a higher risk of developing tumors, particularly those with frequent events of loss of heterozygosity^[13].

According to the family study, just like the patient in our report, the Clinvar database categorizes his gene as causative, and his mother and daughter have the same gene mutation site, and their colorectal cancer risk was more than approximately three-fold^[14-15, 16]. And depending on the genetic make-up of mother and daughter, it is more like an autosomal dominant inheritance. In addition, family genealogy analyzes can be used to better evaluate the common mutations in Asian populations in order to better assess and intervene in colorectal cancer hazard in monoallelic carriers^[15].

It was worth mentioning that he has another mutation in LHX4, but the genetic report does not explicitly mention the association with colorectal cancer. At present, the expression pattern of LHX gene in colorectal cancer is still unclear. The literature posits that LHX4 upregulates β -catenin levels in colorectal cancer cell lines, and LHX4 associated mutations or deletion disrupted the direct LHX4- β -catenin interaction, as well as significantly reduced the capability of LHX4 to both combine and trans-increase target gene promoters. Importantly, deletion or mutation of LHX4 also abolished its tumor-promoting function, suggesting that its mediated LHX4-catenin interaction was crucial for LHX4's tumor suppressor functions^[17]. It is unknown whether the progression of colon polyps is balanced when the two genetic mutations are present at the same time. In clinical work, we must pay more attention to the results of patients' genetic examinations to prevent rare reported gene mutations from being overlooked, and carry out individual follow-up protocols to study in detail the clinical manifestations caused by polygenic mutations.

Malignant tumors accounted for up to 90% of adult intussusception, so if adult intussusception is present, the possibility of tumorigenesis cannot be overlooked and endoscopy should be considered for surveillance screening and regular follow-up^[18].

In our case report, patients should appropriately lengthen and shorten the endoscopic follow-up time according to the bowel preparation and whether the pathology is prone to cancer, and individualize the endoscopic polypectomy plan so that the patient's lesions do not develop as much as possible.

Conclusions

At present, colon cancer triggered by homozygous mutations in the MUTYH gene was no longer surprising, and serious lesions caused by heterozygous variants have rarely been reported. Our results underscored the need for closer follow-up when patients with this gene mutation are present, and we recommended that family members of patients should be included in the follow-up population.

Ethics

Written informed consent was obtained from the patient for inclusion in this case report. The case was approved by the institutional review board of the affiliated Drum Tower Hospital of Nanjing University, Medical School (2023-CR002-01).

Statement

No funding was obtained for this study.

Competing interests

All the authors declared that they had no conflicts of interest or financial ties to disclose.

References

1. Galiatsatos P, Foulkes WD. Familial adenomatous polyposis. *Am J Gastroenterol*. 2006 Feb;101(2):385-98. doi: 10.1111/j.1572-0241.2006.00375.x. PMID: 16454848.
2. Al-Tassan N, Chmiel NH, Maynard J, Fleming N, Livingston AL, Williams GT, Hodges AK, Davies DR, David SS, Sampson JR, Cheadle JP. Inherited variants of MYH associated with somatic G:C->T:A mutations in colorectal tumors. *Nat Genet*. 2002 Feb;30(2):227-32. doi: 10.1038/ng828. Epub 2002 Jan 30. PMID: 11818965.
3. Croitoru ME, Cleary SP, Di Nicola N, Manno M, Selander T, Aronson M, Redston M, Cotterchio M, Knight J, Gryfe R, Gallinger S. Association between biallelic and monoallelic germline MYH gene mutations and colorectal cancer risk. *J Natl Cancer Inst*. 2004 Nov 3;96(21):1631-4. doi: 10.1093/jnci/djh288. PMID: 15523092.
4. Win AK, Hopper JL, Jenkins MA. Association between monoallelic MUTYH mutation and colorectal cancer risk: a meta-regression analysis. *Fam Cancer*. 2011 Mar;10(1):1-9. doi: 10.1007/s10689-010-9399-5. PMID: 21061173; PMCID: PMC3228836.
5. Takao M, Yamaguchi T, Eguchi H, Yamada T, Okazaki Y, Tomita N, Nomizu T, Momma T, Takayama T, Tanakaya K, Akagi K, Ishida H. APC germline variant analysis in the adenomatous polyposis phenotype in Japanese patients. *Int J Clin Oncol*. 2021 Sep;26(9):1661-1670. doi: 10.1007/s10147-021-01946-4. Epub 2021 Jun 9. Erratum in: *Int J Clin Oncol*. 2022 Feb;27(2):456. PMID: 34106356.
6. Li N, Kang Q, Yang L, Zhao XJ, Xue LJ, Wang X, Li AQ, Li CG, Sheng JQ. Clinical characterization and mutation spectrum in patients with familial adenomatous polyposis in China. *J Gastroenterol Hepatol*. 2019 Sep;34(9):1497-1503. doi: 10.1111/jgh.14704. Epub 2019 Jun 13. PMID: 31062380.
7. Li CG, Jin P, Yang L, Zang WC, Kang Q, Li N, He Y, Xu J, Zhang C, Wang X, Sheng JQ. Germline mutations in patients with multiple colorectal polyps in China. *J Gastroenterol Hepatol*. 2017 Oct;32(10):1723-1729. doi: 10.1111/jgh.13776. PMID: 28251689.
8. Theodoratou E, Campbell H, Tenesa A, Houlston R, Webb E, Lubbe S, Broderick P, Gallinger S, Croitoru EM, Jenkins MA, Win AK, Cleary SP, Koessler T, Pharoah PD, Küry S, Bézieau S, Buecher B, Ellis NA, Peterlongo P, Offit K, Aaltonen LA, Enholm S, Lindblom A, Zhou XL, Tomlinson IP, Moreno V, Blanco I, Capellà G, Barnetson R, Porteous ME, Dunlop MG, Farrington SM. A large-scale meta-analysis to refine colorectal cancer risk estimates associated with MUTYH variants. *Br J Cancer*. 2010 Dec 7;103(12):1875-84. doi: 10.1038/sj.bjc.6605966. Epub 2010 Nov 9. PMID: 21063410; PMCID: PMC3008602.
9. Boparai KS, Dekker E, Van Eeden S, Polak MM, Bartelsman JF, Mathus-Vliegen EM, Keller JJ, van Noesel CJ. Hyperplastic polyps and sessile serrated adenomas as a phenotypic expression of MYH-associated polyposis. *Gastroenterology*. 2008 Dec;135(6):2014-8. doi: 10.1053/j.gastro.2008.09.020. Epub 2008 Sep 20. PMID: 19013464.
10. Sieber OM, Lipton L, Crabtree M, Heinimann K, Fidalgo P, Phillips RK, Bisgaard ML, Orntoft TF, Aaltonen LA, Hodgson SV, Thomas HJ, Tomlinson IP. Multiple colorectal adenomas, classic adenomatous polyposis, and germ-line mutations in MYH. *N Engl J Med*. 2003 Feb 27;348(9):791-9. doi: 10.1056/NEJMoa025283. PMID: 12606733.
11. Claes K, Dahan K, Tejpar S, De Paepe A, Bonduelle M, Abramowicz M, Verellen C, Franchimont D, Van Cutsem E, Kartheuser A. The genetics of familial adenomatous polyposis (FAP) and MutYH-

- associated polyposis (MAP). *Acta Gastroenterol Belg.* 2011 Sep;74(3):421-6. PMID: 22103048.
12. Rosner G, Bercovich D, Daniel YE, Strul H, Fliss-Isakov N, Ben-Yehoiada M, Santo E, Halpern Z, Kariv R. Increased risk for colorectal adenomas and cancer in mono-allelic MUTYH mutation carriers: results from a cohort of North-African Jews. *Fam Cancer.* 2015 Sep;14(3):427-36. doi: 10.1007/s10689-015-9799-7. PMID: 25822476.
13. Barreiro RAS, Sabbaga J, Rossi BM, Achatz MIW, Bettoni F, Camargo AA, Asprino PF, A F Galante P. Monoallelic deleterious MUTYH germline variants as a driver for tumorigenesis. *J Pathol.* 2022 Feb;256(2):214-222. doi: 10.1002/path.5829. Epub 2021 Nov 23. PMID: 34816434.
14. Jones N, Vogt S, Nielsen M, Christian D, Wark PA, Eccles D, Edwards E, Evans DG, Maher ER, Vasen HF, Hes FJ, Aretz S, Sampson JR. Increased colorectal cancer incidence in obligate carriers of heterozygous mutations in MUTYH. *Gastroenterology.* 2009 Aug;137(2):489-94, 494.e1; quiz 725-6. doi: 10.1053/j.gastro.2009.04.047. Epub 2009 Apr 23. PMID: 19394335.
15. Jenkins MA, Croitoru ME, Monga N, Cleary SP, Cotterchio M, Hopper JL, Gallinger S. Risk of colorectal cancer in monoallelic and biallelic carriers of MYH mutations: a population-based case-family study. *Cancer Epidemiol Biomarkers Prev.* 2006 Feb;15(2):312-4. doi: 10.1158/1055-9965.EPI-05-0793. PMID: 16492921.
16. Win AK, Cleary SP, Dowty JG, Baron JA, Young JP, Buchanan DD, Southey MC, Burnett T, Parfrey PS, Green RC, Le Marchand L, Newcomb PA, Haile RW, Lindor NM, Hopper JL, Gallinger S, Jenkins MA. Cancer risks for monoallelic MUTYH mutation carriers with a family history of colorectal cancer. *Int J Cancer.* 2011 Nov 1;129(9):2256-62. doi: 10.1002/ijc.25870. Epub 2011 Apr 8. PMID: 21171015; PMCID: PMC3291738.
17. Cha N, Liu W, Yang N, Xie S, Gao Y, Chen X, Wang X, Ren J. Oncogenicity of LHX4 in colorectal cancer through Wnt/ β -catenin/TCF4 cascade. *Tumour Biol.* 2014 Oct;35(10):10319-24. doi: 10.1007/s13277-014-2210-8. Epub 2014 Jul 19. PMID: 25034524.
18. de Mesquita GHA, Carvalho BJ, de Almeida Medeiros KA, Nii F, Martines DR, Pipek LZ, Jardim YJ, Waisberg DR, Obara MT, Sitnik R, Meyer A, Manguiera CLP. Intussusception reveals MUTYH-associated polyposis syndrome and colorectal cancer: a case report. *BMC Cancer.* 2019 Apr 5;19(1):324. doi: 10.1186/s12885-019-5505-8. PMID: 30953464; PMCID: PMC6451307.




Figure 1. Family tree of patient The proband's age at onset of the polyposis (33 years). The patient's father was unaffected, but his mother and daughter had a heterozygous mutation(without colonoscopy).




Figure 2. Partial chromatogram of the junction sequencing MUTYH gene sequence analysis confirmed the following heterozygous mutation: c.55CT (p.R19X) in exon 2 (ClinVar NM.001128425). Only the patient's father does not have the mutation in this gene.





Figure 3. First colonoscopy in our hospital A:A huge mass can be seen at the junction of the descending colon and the sigmoid colon.



B: This is the specimen of the lesion that was removed. The pathology indicates that it is a villous tubular adenoma, most of which are accompanied by low-grade intraepithelial neoplasia, and a small number of high-grade intraepithelial neoplasia on the surface of the adenoma.




Figure 4. Second colonoscopy in our hospital The intestinal mucosa was not completely exposed because the intestinal preparation was not good and only a few polyps were seen this time. As shown in the image, there are polyp-like growths in the descending colon.




Figure 5. Third colonoscopy in our hospital This time colonoscopy found many flat (0-IIa and 0- IIb)polyps. It is worth noting that larger lesions missed at the last colonoscopy were also seen in the rectum and at the sigmoid junction. Not surprisingly, Polyps remain concentrated in the left colon.