

Quantitative and Qualitative Analysis of a Quantum Multisingular Problem by Computational Method and Heatmaps

Reny George¹, Fahad Sameer Alshammari¹, Mehran Ghaderi², and Shahram Rezapour³

¹Prince Sattam bin Abdulaziz University


²Dai Hoc Duy Tan


³China Medical University Hospital

September 11, 2023

Abstract

The physical phenomena with uncontrollable singularities pose challenges in solving related differential equations. In this work, we intend to investigate the quantitative and qualitative aspects of a multi-singular integro-differential equation with the help of quantum fractional operators by presenting numerical algorithms. Quantum calculus enables us to use numerical algorithms and software. The α - ψ -contraction, a new technique of fixed point theory, plays a significant role in proving the existence of the solution. To interpret tables with quantum values quickly and easily, we use heatmaps. We also presented three numerical examples to illustrate the accuracy and efficiency of our main results.





Quantitative and Qualitative Analysis of a Quantum Multisingular Problem by Computational Method and Heatmaps

Reny George^{1*}, Fahad Sameer Alshammari¹, Mehran Ghaderi², Shahram Rezapour³

¹ Department of Mathematics, College of Science and Humanities in Al-Kharj, Prince Sattam Bin Abdulaziz University, Al-Kharj 11942, Saudi Arabia, e-mail:r.kunnelchacko@psau.edu.sa,

² Institute of Research and Development, Duy Tan University, Da Nang 550000, Vietnam, e-mail:ghaderimehr@gmail.com

³ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan, email:Rezapourshahram@yahoo.ca

Abstract

The physical phenomena with uncontrollable singularities pose challenges in solving related differential equations. In this work, we intend to investigate the quantitative and qualitative aspects of a multi-singular integro-differential equation with the help of quantum fractional operators by presenting numerical algorithms. Quantum calculus enables us to use numerical algorithms and software. The α - ψ -contraction, a new technique of fixed point theory, plays a significant role in proving the existence of the solution. To interpret tables with quantum values quickly and easily, we use heatmaps. We also presented three numerical examples to illustrate the accuracy and efficiency of our main results.

(2010) Mathematics Subject Classifications: 34A08; 34A12.

Keywords: Fixed point theory; α - ψ -contraction; Quantum calculus; Multi-singular equation; Fractional calculus.

Acknowledgments: The authors extend their appreciation to the Deputyship for Research & Innovation, Ministry of Education in Saudi Arabia for funding this research work through the project number (IF2/PSAU/2022/01/22923).

1 Introduction

Scientists in natural and mathematical fields constantly seek new approaches to model complex phenomena. Fractional operators are a prime example of this. Researchers in biological, physical, and engineering sciences are giving special attention to fractional differential equations ([1, 2, 3, 4, 5, 6, 7, 8]), as new laboratory data confirm their greater efficiency [9]. Over time, as fractional calculus has advanced, various new operators have been introduced and

*Corresponding author

generalized. For example, we can mention Caputo, ψ -Caputo, Caputo-Fabrizio, Hadamard, Hilfer, Riemann-Liouville and Atangana-Baleanu. Some of the contributions made with the mentioned operators are: ([10, 11, 12, 13, 14, 15, 16, 17, 18]). Ongoing research on these operators presents new findings daily. George et. al. [19] recently demonstrated that the choice of the ψ function affects the boundedness of the solution set of the pantograph equation with the ψ -Caputo fractional operator.

Modeling natural phenomena often requires solving differential equations with singular points, which can be complicated and challenging. Therefore, the study of such equations to find analytical and numerical solutions has recently gained attention. In 2010, Agarwal et. al. [20] investigated positive solution for the following singular dirichlet problem

$$\begin{cases} \mathfrak{D}^\gamma \mathbf{v}(\kappa) + \mathbf{h}(\kappa, \mathbf{v}(\kappa), \mathfrak{D}^\tau \mathbf{v}(\kappa)) = 0, & \kappa \in (0, 1), \\ \mathbf{v}(0) = \mathbf{v}(1) = 0, \end{cases}$$

where $\gamma \in (1, 2)$, $0 < \tau \leq \gamma - 1$, \mathfrak{D}^γ denotes Riemann-Liouville derivative of fractional order γ and $\mathbf{h}(\kappa, u, v)$ is singular at $t = 0$. In 2011, Feng et. al [21] studied the existence of solution for the following singular system using the krasnoselskii's and Leray-Schauder theorem

$$\begin{cases} \mathfrak{D}^\gamma \mathbf{v}(\kappa) + \mathbf{h}(\kappa, \mathbf{u}(\kappa)) = 0, & \kappa \in (0, 1), \\ \mathfrak{D}^\mathcal{L} \mathbf{u}(\kappa) + \mathbf{t}(\kappa, \mathbf{v}(\kappa)) = 0, & \kappa \in (0, 1), \\ \mathbf{v}(0) = \mathbf{v}'(0) = \mathbf{v}(1) = 0, \\ \mathbf{u}(0) = \mathbf{u}'(0) = \mathbf{u}(1) = 0, \end{cases}$$

such that $\gamma, \mathcal{L} \in (2, 3]$, $\mathfrak{D}^\gamma, \mathfrak{D}^\mathcal{L}$ are fractional Riemann-Liouville derivatives and \mathbf{h}, \mathbf{t} are realvalued continouos functions on $[0, \infty) \times \mathbb{R}$. In 2013, Vong [22] examined the posetive solution for the following singular problem:

$$\begin{cases} {}^c\mathfrak{D}^\gamma \mathbf{v}(\kappa) + \mathbf{h}(\kappa, \mathbf{v}(\kappa)) = 0, & \kappa \in [0, 1], \\ \mathbf{v}(1) = \int_0^1 \mathbf{v}(p) dp, \\ \mathbf{v}'(0) = \dots = \mathbf{v}^{(n-1)}(0) = 0, \\ \int_0^1 dp < 1, \end{cases}$$

using upper and lower solutions method with Schauder fixed point, where $n \geq 2$, $n-1 < \gamma < n$ and \mathbf{h} have singularity at $\kappa = 1$. In 2014, Nyamoradi et. al. [23] reviewed the existence and uniqueness for the following singular BVP:

$$\begin{cases} -\mathfrak{D}^\gamma \mathbf{v}(\kappa) + \mathcal{M}\mathbf{h}(\kappa, \mathbf{v}(\kappa)) + \sum_{i=1}^m \mathcal{N}_i \mathfrak{I}^{\sigma_i} \mathbf{t}_i(\kappa, \mathbf{v}(\kappa)), & \kappa \in (0, 1), \\ \mathfrak{D}^\tau \mathbf{v}(0) = 0, \\ \mathfrak{D}^\tau \mathbf{v}(1) = b \mathfrak{D}^{\frac{\gamma-\tau-1}{2}} (\mathfrak{D}^\tau \mathbf{v}(\kappa))|_{\kappa=\ell}, & \ell \in (0, \frac{1}{2}], \end{cases}$$

where $1 < \gamma \leq 2$, $\tau_i \in (0, 1)$, $b \in [0, \infty)$, further $\mathcal{M}, \mathcal{N}_i$ are real constant and \mathfrak{D}^γ is the fractional Riemann-Liouville derivative. In 2022, Malekpour et. al. [24] studied a new class of singular equation namely multi-singular pointwised defined system as follows:

$$\begin{cases} {}^c\mathfrak{D}^{\gamma_1}\mathbf{v}(\kappa) + \mathbf{h}_1(\kappa, \mathbf{v}(\kappa), \mathbf{u}(\kappa), \mathbf{v}'(\kappa), \mathbf{u}'(\kappa), {}^c\mathfrak{D}^{\tau_1}\mathbf{v}(\kappa), \\ {}^c\mathfrak{D}^{\tau_2}\mathbf{u}(\kappa), \int_0^\kappa g_1(\lambda)\mathbf{v}(\lambda) d\lambda, \int_0^\kappa g_2(\lambda)\mathbf{v}(\lambda) d\lambda) = 0, \\ {}^c\mathfrak{D}^{\gamma_2}\mathbf{v}(\kappa) + \mathbf{h}_2(\kappa, \mathbf{v}(\kappa), \mathbf{u}(\kappa), \mathbf{v}'(\kappa), \mathbf{u}'(\kappa), {}^c\mathfrak{D}^{\tau_1}\mathbf{v}(\kappa), \\ {}^c\mathfrak{D}^{\tau_2}\mathbf{u}(\kappa), \int_0^\kappa g_1(\lambda)\mathbf{v}(\lambda) d\lambda, \int_0^\kappa g_2(\lambda)\mathbf{v}(\lambda) d\lambda) = 0, \end{cases}$$

under boundary conditions:

$$\begin{cases} {}^c\mathfrak{D}^{\sigma_1}\mathbf{v}(\omega_1) = \ell_1, & {}^c\mathfrak{D}^{\sigma_2}\mathbf{v}(\omega_2) = \ell_2, \\ \mathbf{v}(1) = \mathbf{v}^{(j)}(0) = 0, & \mathbf{u}(1) = \mathbf{u}^{(j)}(0) = 0, \quad j \geq 2 \end{cases}$$

where $\gamma_i \geq 2$, $\omega_i, \sigma_i \in (0, 1)$, $\ell_i \geq 0$, $\mathbf{h}_i, g_i \in \mathbf{L}^1$ such that \mathbf{h}_i is a singular at some points in $[0, 1]$ for $i = 1, 2$. The reader can refer to [25, 26, 27] to learn about the rest of the research done in this field.

Computers have always struggled with finding numerical solutions and computational algorithms for singular equations. In this case, we are seeking a numerical method to solve the multi-singular integrodifferential equation. We are using quantum operators to provide the necessary space for software packages. In 1910, Frank Hilton Jackson introduced the quantum derivative operator [28, 29]. In his definition of derivative, namely $(\mathcal{D}_q\mathbf{v})(\kappa) = \mathbf{v}(\kappa) - \mathbf{v}(q\kappa)/(1 - q)\kappa$, he removed the concept of limit, and this caused a discrete space to be prepared for the analysis of problems. In his quantum calculus for every real number κ defined the q -analogue of κ as: $[\kappa]_q = \frac{1-q\kappa}{1-q} = 1 + q + \dots + q^{\kappa-1}$. Moreover, q -analogue of the power function $(\kappa - p)^n$ for $n \geq 1$ defined as: $(\kappa - p)_q^{(n)} = \prod_{j=0}^{n-1}(\kappa - pq^j)$ and $(\kappa - p)_q^{(0)} = 1$, where $\kappa, p \in \mathbb{R}$. In addition, Let $\kappa \in \mathbb{R} - \{0, -1, -2, \dots\}$, then the quantum gamma function formulated as: $\Gamma_q(\kappa) = \frac{(1-q)^{(\kappa-1)}}{(1-q)^{\kappa-1}}$. Also, it is worth mentioning that $\Gamma_q(\kappa + 1) = [\kappa]_q\Gamma_q(\kappa)$ holds true. In the preliminaries section, we present an algorithm for calculating the quantum gamma function. The characteristics of quantum operators have been investigated in detail by Kac and Cheung in [30]. Later in 2007, the properties of fractional operator in q -calculus was developed in works [31] and [32]. In 2011, Ferreira [33] investigated positive solution for the following q -fractional BVP:

$$\begin{cases} \mathfrak{D}_q^\gamma\mathbf{v}(\kappa) = -\mathbf{h}(\kappa, \mathbf{v}(\kappa)), & \kappa \in (0, 1), \\ \mathbf{v}(0) = \mathfrak{D}_q\mathbf{v}(0) = 0, \\ \mathfrak{D}_q\mathbf{v}(1) = c \geq 0, \end{cases}$$

where $\gamma \in (2, 3]$, $c \in \mathbb{R}$, \mathfrak{D}_q^γ denotes the q -Riemann-Liouville derivative of fractional order γ and $\mathbf{h} : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is nonnegative continuous function. In 2015, Xinhui Li et.al. [34]

studied BVP of fractional q -difference Schringer equation:

$$\begin{cases} \mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \frac{\mathbf{m}}{\mathbf{p}}(\mathbf{E} - \mathbf{u}(\kappa))\mathbf{v}(\kappa) = 0, & \kappa \in (0, 1), \\ \mathbf{v}(0) = \mathfrak{D}_q \mathbf{v}(0) = \mathfrak{D}_q \mathbf{v}(1) = 0, \end{cases}$$

where $\gamma \in (2, 3)$, \mathbf{m}, \mathbf{E} are the mass and energy particle respectively, and \mathbf{p} is the Plank constant. Many articles have been published in the field of quantum differential problems in recent years, for information about which you can refer to [35, 36, 37, 38, 39, 40].

Here, by getting motivation from works [20, 22, 24, 33] and also filling the gap in numerical methods for studying singular equations, we are going to examine the following uncontrolled multi-singular pointwise defined equation in q -calculus:

$${}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \mathbf{h}(\kappa, \mathbf{v}(\kappa), \mathbf{v}'(\kappa), {}^c\mathfrak{D}_q^\tau \mathbf{v}(\kappa), \mathfrak{I}_q^\sigma \mathbf{v}(\kappa)) = 0, \quad \kappa \in \mathcal{K} = [0, 1] \quad (1.1)$$

under the following boundary conditions:

$$\begin{cases} \mathbf{v}'(0) = \mathbf{v}(\omega), \\ \mathbf{v}(1) = \int_0^\ell \mathbf{v}(p) dp, \\ \mathbf{v}^{(2)}(0) = \dots = \mathbf{v}^{(n-1)}(0) = 0, \quad n = [\gamma] + 1, \end{cases} \quad (1.2)$$

where $\gamma \geq 3$, $\sigma > 1$, $\tau, \omega, \ell \in (0, 1)$, $\mathbf{v} \in \mathbf{C}^1[0, 1]$, $\mathbf{h} : \mathcal{K} \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is function where $\mathbf{h}(\kappa, \dots, \dots)$ singular at some point $\kappa \in [0, 1]$ and ${}^c\mathfrak{D}_q^\gamma$ denote the q -Caputo derivative of fractional order γ . The function \mathbf{h} is multi-singular when it is singular at more than one point. Note that we will continue to do all our calculations on the time scale, namely $TS_{\kappa_0} = \{\kappa_0, \kappa_0 q, \kappa_0 q^2, \dots\} \cup \{0\}$, where $\kappa_0 \in \mathbb{R}$, and $q \in (0, 1)$.

2 Preliminaries

Notation 2.1. *It should be noted that throughout this work: $\|\cdot\|$ is the sup norm of $\mathbf{C}[0, 1]$, $\|\cdot\|_1$ is the norm of $\mathbf{L}^1[0, 1]$ and $\|\mathbf{v}\|_* = \max\{\|\mathbf{v}\|, \|\mathbf{v}'\|\}$ is the norm of $\mathbf{Y} = \mathbf{C}^1[0, 1]$.*

Notation 2.2. *Assume that the function $\psi : [0, \infty) \rightarrow [0, \infty)$ for all $\kappa > 0$ be such that $\sum_{n=1}^{\infty} \psi^n(\kappa) < \infty$. Then we denote the family of nondecreasing functions ψ by Ψ . Notice that $\forall \kappa > 0$ we have $\psi(\kappa) < \kappa$ (see [41]).*

Definition 2.3. [22] *consider the following equation*

$${}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \mathbf{h}(\kappa) = 0, \quad \kappa \in [0, 1]. \quad (2.1)$$

If there exist a set such as $\mathcal{P} \subset [0, 1]$ such that $\mu(\mathcal{P}^c) = 0$ and Eq. (2.1) hold on \mathcal{P} , then we call Eq. (2.1) pointwise defined equation where μ is the measure function.

Definition 2.4. [29] Let $\kappa \in \mathbb{R} - \{0, -1, -2, \dots\}$, then the quantum gamma function formulated as follows

$$\Gamma_q(\kappa) = \frac{(1-q)^{(\kappa-1)}}{(1-q)^{\kappa-1}},$$

also, it is worth mentioning that $\Gamma_q(\kappa + 1) = [\kappa]_q \Gamma_q(\kappa)$ holds true. we present an algorithm for calculating the quantum gamma function. Moreover, we computed for some values of q in Tables 1 and 2, also their heatmaps presented in Figures 1 and 2.

Algorithm 1 The proposed procedure To calculate $\Gamma_q(\kappa)$.

function quantum gamma = $qG(q, \kappa, r)$

$t = 1;$

for $j = 0 : r$

$t = t * (1 - q^{(j+1)}) / (1 - q^{(\kappa+j)});$

end

$qG = t / (1 - q)^{(\kappa-1)};$

end

r	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$v = 3.75$						
1	1.1498	1.3475	1.8340	2.7443	5.4843	18.2050
2	1.1492	1.3410	1.7711	2.4622	4.3021	12.1461
3	1.1492	1.3398	1.7511	2.3392	3.7104	9.1158
4	1.1492	1.3397	1.7445	2.2822	3.3808	7.3785
5	1.1492	1.3396	1.7424	2.2550	3.1853	6.2897
6	1.1492	1.3396	1.7417	2.2418	3.0649	5.5639
7	1.1492	1.3396	1.7414	2.2354	2.9888	5.0579
8	1.1492	1.3396	1.7414	2.2323	2.9401	4.6934
9	1.1492	1.3396	1.7413	2.2307	2.9085	4.4241
...
13	1.1492	1.3396	1.7413	2.2294	2.8599	3.8528
14	1.1492	1.3396	1.7413	2.2293	2.8560	3.7792
...
26	1.1492	1.3396	1.7413	2.2293	2.8487	3.5085
27	1.1492	1.3396	1.7413	2.2293	2.8486	3.5042
..
39	1.1492	1.3396	1.7413	2.2293	2.8486	3.4877
40	1.1492	1.3396	1.7413	2.2293	2.8486	3.4874
...
52	1.1492	1.3396	1.7413	2.2293	2.8486	3.4864
53	1.1492	1.3396	1.7413	2.2293	2.8486	3.4863

Table 1: Numerical results for $\Gamma_q(3.75)$ for different value of q

r	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$v = 1.6$						
1	0.9796	0.9672	0.9686	1.0111	1.1428	1.4660
2	0.9792	0.9641	0.9513	0.9662	1.0494	1.2920
3	0.9792	0.9636	0.9458	0.9461	0.9978	1.1855
4	0.9792	0.9635	0.9440	0.9366	0.9673	1.1147
5	0.9792	0.9635	0.9434	0.9321	0.9484	1.0649
6	0.9792	0.9635	0.9432	0.9298	0.9365	1.0287
7	0.9792	0.9635	0.9432	0.9288	0.9288	1.0016
8	0.9792	0.9635	0.9432	0.9282	0.9239	0.9810
9	0.9792	0.9635	0.9432	0.9280	0.9206	0.9652
10	0.9792	0.9635	0.9431	0.9279	0.9185	0.9528
...
14	0.9792	0.9635	0.9431	0.9277	0.9152	0.9245
15	0.9792	0.9635	0.9431	0.9277	0.9150	0.9206
...
25	0.9792	0.9635	0.9431	0.9277	0.9145	0.9064
26	0.9792	0.9635	0.9431	0.9277	0.9144	0.9060
...
34	0.9792	0.9635	0.9431	0.9277	0.9144	0.9048
35	0.9792	0.9635	0.9431	0.9277	0.9144	0.9047
...
40	0.9792	0.9635	0.9431	0.9277	0.9144	0.9046
41	0.9792	0.9635	0.9431	0.9277	0.9144	0.9045

Table 2: Numerical results for $\Gamma_q(1.6)$ for different value of q




Figure 1: The heatmap of Table 1.




Figure 2: The heatmap of Table 2.

Definition 2.5. [42] Suppose that $\mathbf{v}(\kappa) : [0, \infty] \rightarrow \mathbb{R}$, be a continuous function, then its fractional Riemann-Liouville quantum integral and its fractional Caputo quantum derivative are expressed respectively by

$$\mathfrak{I}_q^\gamma \mathbf{v}(\kappa) = \frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{v}(p) \, d_q p,$$

and

$${}^c \mathfrak{D}^\gamma \mathbf{v}(\kappa) = \frac{1}{\Gamma_q(n - \gamma)} \int_0^\kappa (\kappa - qp)^{n-\gamma-1} \mathfrak{D}_q^n \mathbf{v}(p) \, d_q p, \quad n = [\gamma] + 1.$$

Lemma 2.6. [43] assume that $n = [\gamma] + 1$, then the following relation holds true

$$({}^C \mathfrak{I}_q^\eta {}^C \mathfrak{D}_q^\eta \mathbf{v})(\kappa) = \mathbf{v}(\kappa) - \sum_{j=0}^{n-1} \frac{\mathbf{v}^j}{\Gamma_q(j+1)} (\mathfrak{D}_q^j \mathbf{v})(0),$$

which is deduced from it, the general solution for ${}^C \mathfrak{D}_q^\eta \mathbf{v}(\kappa) = 0$, expressed by

$$\mathbf{v}(\kappa) = r_0 + r_1 \kappa + r_2 \kappa^2 + \cdots + r_{n-1} \kappa^{n-1},$$

where $r_0, \dots, r_{n-1} \in \mathbb{R}$.

Definition 2.7. [44] Consider the two maps $\alpha : \mathbf{Y} \times \mathbf{Y} \rightarrow [0, \infty)$ and $\mathcal{T} : \mathbf{Y} \rightarrow \mathbf{Y}$. The function \mathcal{T} is called α -admissible if $\alpha(\mathbf{v}, \mathbf{u}) \geq 1$ yields that $\alpha(\mathcal{T}\mathbf{v}, \mathcal{T}\mathbf{u}) \geq 1$. Moreover, let $\psi \in \Psi$, $\mathcal{T} : \mathbf{Y} \rightarrow \mathbf{Y}$, and $(\mathbf{Y}, d_{\mathbf{Y}})$ be a complete metric space. If for every $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ we have $\alpha(\mathbf{v}, \mathbf{u}) d_{\mathbf{Y}}(\mathcal{T}\mathbf{v}, \mathcal{T}\mathbf{u}) \leq \psi(d_{\mathbf{Y}}(\mathbf{v}, \mathbf{u}))$, then the map \mathcal{T} is called α - ψ -contraction.

Lemma 2.8. [44] Suppose that $\alpha : \mathbf{Y} \times \mathbf{Y} \rightarrow [0, \infty)$, $\psi \in \Psi$, $(\mathbf{Y}, d_{\mathbf{Y}})$ be a complete metric space and $\mathcal{T} : \mathbf{Y} \rightarrow \mathbf{Y}$ is an α - ψ -contraction. If \mathcal{T} be continuous and $\exists \mathbf{v}_0 \in \mathbf{Y}$ where $\alpha(\mathbf{v}_0, \mathcal{T}\mathbf{v}_0) \geq 0$, then \mathcal{T} has a fixed point.

In the following, we present two key lemmas about the quantum Green's function and its properties.

Lemma 2.9. Consider the following fractional quantum pointwise defined problem

$${}^c \mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \mathbf{u}_0 = 0 \tag{2.2}$$

with boundary conditions

$$\begin{cases} \mathbf{v}'(0) = \mathbf{v}(\omega), \\ \mathbf{v}(1) = \int_0^\ell \mathbf{v}(p) \, d_p p, \\ \mathbf{v}^{(j)}(0) \text{ for } j = 2, \dots, [\gamma], \end{cases} \tag{2.3}$$

such that $\mathbf{u}_0 \in \mathbf{L}^1[0, 1]$, $0 < \ell, \omega < 1$ and $\gamma \in [2, 3)$. Then the function $\mathbf{v}(\kappa) = \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{u}_0(p) \, d_q p$ is a solution for problem mentioned in (2.2)-(2.3) such that

$$\mathcal{G}_q(\kappa, p) = \mathcal{L}_q(\kappa, p) + \frac{1}{1 - \ell} \int_0^\ell \mathcal{L}_q(\kappa, p) \, d_p p$$

and

$$\mathcal{L}_q(\kappa, p) = \frac{(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1}}{\Gamma_q(\gamma)},$$

where $0 \leq qp \leq \kappa \leq 1$ and $qp \leq \omega$

$$\mathcal{L}_q(\kappa, p) = \frac{(1 - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1}}{\Gamma_q(\gamma)},$$

where $0 \leq \omega \leq qp \leq \kappa \leq 1$

$$\mathcal{L}_q(\kappa, p) = \frac{(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1}}{\Gamma_q(\gamma)},$$

where $0 \leq \kappa \leq qp \leq \omega \leq 1$

$$\mathcal{L}_q(\kappa, p) = \frac{(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)},$$

where $0 \leq \kappa \leq qp \leq 1$ and $\omega \leq qp$.

Proof. Let the equation ${}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \mathbf{u}_0(\kappa) = 0$ satisfy for all $\kappa \in \mathcal{P}$ such that $\mathcal{P} \subset [0, 1]$ and measure of \mathcal{P}^c is zero. Now, we take $\mathbf{u} \in \mathbf{L}^1[0, 1] \cap \mathbf{C}^1[0, 1]$ where $\mathbf{u} = \mathbf{u}_0$ on \mathcal{P} . If \mathbf{v}_0 be a solution for Eq. (2.2)-(2.3), then for all $\kappa \in [0, 1]$ put $\mathbf{u}(\kappa) = -{}^c\mathfrak{D}_q^\gamma \mathbf{v}_0(\kappa)$. Moreover, we can write

$$\begin{aligned} \mathfrak{I}_q^\gamma(\mathbf{u}_0(\kappa)) &= \frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}_0(p) \, d_q p \\ &= \frac{1}{\Gamma_q(\gamma)} \left(\int_{[0, \kappa] \cap \mathcal{E}} (\kappa - qp)^{\gamma-1} \mathbf{u}_0(p) \, d_q p + \int_{[0, \kappa] \cap \mathcal{E}^c} (\kappa - qp)^{\gamma-1} \mathbf{u}_0(p) \, d_q p \right) \\ &= \frac{1}{\Gamma_q(\gamma)} \int_{[0, \kappa] \cap \mathcal{E}} (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &= \frac{1}{\Gamma_q(\gamma)} \left(\int_{[0, \kappa] \cap \mathcal{E}} (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + \int_{[0, \kappa] \cap \mathcal{E}^c} (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \right) \\ &= \frac{1}{\Gamma_q(\gamma)} \int_0^1 (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p = \mathfrak{I}_q^\gamma(\mathbf{u}(\kappa)). \end{aligned}$$

Now, let $\kappa \in \mathcal{P}^c \setminus \{0\}$. Take $\{\kappa_n\}$ in \mathcal{P} where $\kappa_n \rightarrow \kappa^-$, so

$$\begin{aligned} \mathfrak{I}_q^\gamma(\mathbf{u}_0(\kappa)) &= \frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}_0(p) \, d_q p \\ &= \lim_{n \rightarrow \infty} \frac{1}{\Gamma_q(\gamma)} \int_0^{\kappa_n} (\kappa_n - qp)^{\gamma-1} \mathbf{u}_0(p) \, d_q p = \lim_{n \rightarrow \infty} \mathfrak{I}_q^\gamma(\mathbf{u}_0(\kappa_n)) \\ &= \lim_{n \rightarrow \infty} \mathfrak{I}_q^\gamma(\mathbf{u}(\kappa_n)) = \lim_{n \rightarrow \infty} \frac{1}{\Gamma_q(\gamma)} \int_0^{\kappa_n} (\kappa_n - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \end{aligned}$$

$$= \frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p = \mathfrak{I}_q^\gamma(\mathbf{u}(\kappa)).$$

If $\kappa = 0 \in \mathcal{P}^c$, then $\mathfrak{I}_q^\gamma(\mathbf{u}_0(\kappa)) = \mathfrak{I}_q^\gamma(\mathbf{u}(\kappa)) = 0$, hence for all $\kappa \in [0, 1]$ we have $\mathfrak{I}_q^\gamma(\mathbf{u}_0(\kappa)) = \mathfrak{I}_q^\gamma(\mathbf{u}(\kappa)) = 0$. Therefore, for all $\kappa \in [0, 1]$ we get

$$\mathfrak{I}_q^\gamma({}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa)) = \mathfrak{I}_q^\gamma(-\mathbf{u}_0(\kappa)),$$

such that ${}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa) + \mathbf{u}_0(\kappa) = 0$ for all $\kappa \in \mathcal{P}$. Thus on $[0, 1]$ we have

$$\mathfrak{I}_q^\gamma({}^c\mathfrak{D}_q^\gamma \mathbf{v}(\kappa)) = \mathfrak{I}_q^\gamma(-\mathbf{u}(\kappa)),$$

In view of Lemma 1.1 we have

$$\mathbf{v}(\kappa) = -\frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + r_0 + r_1 \kappa.$$

which from boundary condition (2.3) deduced that

$$r_1 = -\frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p,$$

so

$$\mathbf{v}(\kappa) = -\frac{1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + r_0 - \frac{\kappa}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p.$$

But on the other hand, we have

$$\int_0^\ell \mathbf{v}(p) \, d p = \mathbf{v}(1) = \frac{-1}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + r_0 + -\frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p$$

which yields that

$$r_0 = \int_0^\ell \mathbf{v}(p) \, d p + \frac{1}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + \frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p.$$

Therefore,

$$\begin{aligned} \mathbf{v}(\kappa) &= \frac{-1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + \int_0^\ell \mathbf{v}(p) \, d p + \frac{1}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &\quad + \frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p - \frac{\kappa}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p. \end{aligned}$$

Now, for simplicity set

$$\begin{aligned} \varphi(\kappa) &= \frac{-1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p - \frac{\kappa}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &\quad + \frac{1}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + \frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p, \end{aligned}$$

then, it is clear that $\mathbf{v}(\kappa) = \varphi(\kappa) + \int_0^\ell \mathbf{v}(p) \, dp$. Now, if $\omega \leq \kappa$ then, we get

$$\begin{aligned} \varphi(\kappa) &= \frac{-1}{\Gamma_q(\gamma)} \left(\int_0^\omega + \int_\omega^\kappa \right) (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p - \frac{\kappa}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &\quad + \frac{1}{\Gamma_q(\gamma)} \left(\int_0^\omega + \int_\omega^\kappa + \int_\kappa^1 \right) (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p + \frac{1}{\Gamma_q(\gamma)} \int_0^\omega (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &= \int_0^\omega \frac{(\omega - p)^{\gamma-1} + (1 - p)^{\gamma-1} - \kappa(\omega - p)^{\gamma-1} - (\kappa - p)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p \\ &\quad + \int_\omega^\kappa \frac{(1 - p)^{\gamma-1} - (\kappa - p)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p + \int_\kappa^1 \frac{(1 - p)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p. \end{aligned}$$

But, if $\omega \geq \kappa$ then, we obtain

$$\begin{aligned} \varphi(\kappa) &= \frac{-1}{\Gamma_q(\gamma)} \int_0^\kappa (\kappa - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p - \frac{\kappa}{\Gamma_q(\gamma)} \left(\int_0^\kappa + \int_\kappa^\omega \right) (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &\quad + \frac{1}{\Gamma_q(\gamma)} \left(\int_0^\kappa + \int_\kappa^\omega + \int_\omega^1 \right) (1 - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &\quad + \frac{1}{\Gamma_q(\gamma)} \left(\int_0^\kappa + \int_\kappa^\omega \right) (\omega - qp)^{\gamma-1} \mathbf{u}(p) \, d_q p \\ &= \int_0^\kappa \frac{(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p \\ &\quad + \int_\kappa^\omega \frac{(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p + \int_\omega^1 \frac{(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \mathbf{u}(p) \, d_q p. \end{aligned}$$

Hence,

$$\varphi(\kappa) = \int_0^1 \mathcal{L}_q(\kappa, p) \mathbf{u}(p) \, d_q p,$$

so,

$$\mathbf{v}(\kappa) = \int_0^1 \mathcal{L}_q(\kappa, p) \mathbf{u}(p) \, d_q p + \int_0^\ell \mathbf{v}(p) \, dp. \quad (2.4)$$

By rewrite the Eq.(2.4), we have

$$\begin{aligned} \int_0^\ell \mathbf{v}(\kappa) \, d\kappa &= \int_0^\ell \int_0^1 \mathcal{L}_q(\kappa, p) \mathbf{u}(p) \, d_q p \, d\kappa + \int_0^\ell \int_0^\ell \mathbf{v}(p) \, dp \, d\kappa \\ &= \int_0^1 \left(\int_0^\ell \mathcal{L}_q(\kappa, p) \, d\kappa \right) \mathbf{u}(p) \, d_q p + \ell \int_0^\ell \mathbf{v}(p) \, dp, \end{aligned}$$

which implies that

$$(1 - \ell) \int_0^\ell \mathbf{v}(\kappa) \, d\kappa = \int_0^1 \left(\int_0^\ell \mathcal{L}_q(\kappa, p) \, d\kappa \right) \mathbf{u}(p) \, d_q p$$

and so

$$\int_0^\ell \mathbf{v}(\kappa) \, d\kappa = \int_0^1 \frac{1}{1-\ell} \left(\int_0^\ell \mathcal{L}_q(\kappa, p) \, d\kappa \right) \mathbf{u}(p) \, d_q p.$$

Thus,

$$\begin{aligned} \mathbf{v}(\kappa) &= \int_0^1 \mathcal{L}_q(\kappa, p) \mathbf{u}(p) \, d_q p + \int_0^1 \frac{1}{1-\ell} \left(\int_0^\ell \mathcal{L}_q(\kappa, p) \, d\kappa \right) \mathbf{u}(p) \, d_q p \\ &= \int_0^1 \left(\mathcal{L}_q(\kappa, p) + \frac{1}{1-\ell} \int_0^\ell \mathcal{L}_q(\kappa, p) \, d\kappa \right) \mathbf{u}(p) \, d_q p \\ &= \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{u}(p) \, d_q p = \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{u}_0(p) \, d_q p, \end{aligned}$$

and with this our proof is complete. \square

Lemma 2.10. Assume that $\mathcal{G}_q(\kappa, p)$ be the one defined in Lemma 2.9. Then for all $\kappa, p \in [0, 1]$, the following statments are satisfy

I. $\mathcal{G}_q(\kappa, p) \geq 0$.

II. $\mathcal{G}_q(\kappa, p) \leq \mathcal{A}_q(\gamma, \ell)(1 - qp)^{\gamma-1}$ such that $\mathcal{A}_q(\gamma, \ell) = \frac{3}{(1-\ell)\Gamma_q(\gamma)}$.

III. $\left| \frac{\partial \mathcal{G}_q}{\partial \kappa}(\kappa, p) \right| \leq \mathcal{B}_q(\gamma, \ell)(1 - qp)^{\gamma-1}$ such that $\mathcal{B}_q(\gamma, \ell) = \frac{2}{(1-\ell)\Gamma_q(\gamma-1)}$.

IV. $\mathcal{G}_q(\kappa, p) \geq \frac{(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \left(\frac{2 - \gamma^2}{2(1 - \gamma)} - \kappa \right) \geq 0$.

Proof. I. If $0 \leq qp \leq \kappa \leq 1$ and $qp \leq \omega$, then

$$(\omega - qp)^{\gamma-1} \geq \kappa(\omega - qp)^{\gamma-1} \quad \text{and} \quad (1 - qp)^{\gamma-1} \geq -(\kappa - qp)^{\gamma-1},$$

hence,

$$(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1} \geq 0,$$

and ao $\mathcal{L}_q(\kappa, p) \geq 0$. Thus $\mathcal{G}_q(\kappa, p) \geq 0$. The proof in other case is easy and we skip writing it.

II. One can see that for all $\kappa, p \in [0, 1]$ we have $\mathcal{L}_q(\kappa, p) \leq \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)}$, and so

$$\begin{aligned} \mathcal{G}_q(\kappa, p) &\leq \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} + \frac{1}{1-\ell} \int_0^\ell \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \, d\kappa \\ &= \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} + \frac{\ell}{1-\ell} \times \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} = \frac{3(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \left(1 + \frac{\ell}{1-\ell} \right) \\ &= \frac{3(1 - qp)^{\gamma-1}}{(1-\ell)\Gamma_q(\gamma)} = \mathcal{A}_q(\gamma, \ell)(1 - qp)^{\gamma-1}. \end{aligned}$$

III. Since

$$\frac{\partial \mathcal{L}_q}{\partial \kappa}(\kappa, p) = \begin{cases} \frac{-(\kappa - qp)^{\gamma-1} - (\omega - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)}, & 0 \leq p \leq \kappa \leq 1, \ p \leq \omega, \\ \frac{-(\kappa - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)}, & 0 \leq \omega \leq p \leq \kappa \leq 1, \\ \frac{-(\omega - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)}, & 0 \leq \kappa \leq p \leq \omega \leq 1, \\ 0, & 0 \leq \kappa \leq p \leq 1, \ \omega \leq p. \end{cases}$$

Therefore, we have

$$\left| \frac{\partial \mathcal{L}_q}{\partial \kappa} \right| \leq \frac{(\kappa - qp)^{\gamma-1} + (\omega - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)} \leq \frac{2(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)},$$

hence

$$\begin{aligned} \left| \frac{\partial \mathcal{G}_q}{\partial \kappa} \right| &\leq \frac{2(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)} + \frac{\ell}{1 - \ell} \times \frac{2(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)} = \frac{2(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma-1)} \left(1 + \frac{\ell}{1 - \ell} \right) \\ &= \frac{2(1 - qp)^{\gamma-1}}{(1 - \ell)\Gamma_q(\gamma-1)} = \mathcal{B}_q(\gamma, \ell)(1 - qp)^{\gamma-1}. \end{aligned}$$

IV If $0 < qp < \kappa < 1$ and $qp \leq \omega$, then

$$\kappa(1 - qp) > 0 \quad \text{and} \quad \kappa(1 - qp) - qp + \kappa > 0.$$

Thus, $\kappa - p < \kappa(1 - qp)$ and $\frac{1 - qp}{\kappa - qp} > \frac{1}{\kappa}$. Since $\frac{1}{\kappa} > 1$ and $\gamma \geq 2$, so we have

$$\left(\frac{1 - qp}{\kappa - qp} \right)^{\gamma-1} > \left(\frac{1}{\kappa} \right)^{\gamma} > \frac{1}{\kappa},$$

and

$$\begin{aligned} &(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(\omega - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1} \\ &= (1 - \kappa)(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1} \\ &> (1 - \kappa)(\omega - qp)^{\gamma-1} + (1 - qp)^{\gamma-1} - \kappa(1 - qp)^{\gamma-1} \\ &= (1 - \kappa) \left((1 - qp)^{\gamma-1} + (\omega - qp)^{\gamma-1} \right) \\ &\geq (1 - \kappa)(1 - qp)^{\gamma-1}, \end{aligned}$$

which implies that

$$\mathcal{L}_q(\kappa, p) > (1 - \kappa)(1 - qp)^{\gamma-1}.$$

If $0 < qp \leq \omega < \kappa < 1$ then

$$\begin{aligned} (1 - qp)^{\gamma-1} - (\kappa - qp)^{\gamma-1} &> (1 - qp)^{\gamma-1} - \kappa(1 - qp)^{\gamma-1} \\ &= (1 - \kappa)(1 - qp)^{\gamma-1} \end{aligned}$$

and

$$\mathcal{L}_q(\kappa, p) > \frac{(1 - \kappa)(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)}.$$

Thusly we obtain

$$\begin{aligned} \mathcal{G}_q(\kappa, p) &\geq \frac{1}{\Gamma_q(\gamma)} \left[(1 - \kappa)(1 - qp)^{\gamma-1} + \frac{1}{1 - \ell} \int_0^\ell (1 - \kappa)(1 - qp)^{\gamma-1} d\kappa \right] \\ &= \frac{1}{\Gamma_q(\gamma)} \left[(1 - \kappa)(1 - qp)^{\gamma-1} + \frac{(1 - qp)^{\gamma-1}}{1 - \ell} \times \ell \left(1 - \frac{\ell}{2}\right) \right] \\ &= \frac{(1 - qp)^{\gamma-1}}{\Gamma_q(\gamma)} \left(\frac{2 - \gamma^2}{2(1 - \gamma)} - \kappa \right) \geq 0 \end{aligned}$$

□

3 Main Results

Theorem 3.1. *Let $\mathbf{h} : \mathcal{K} \times (\mathbf{C}[0, 1])^4 \rightarrow \mathbb{R}$ be a singular function at some point $\kappa \in \mathcal{K} = [0, 1]$ such that for all $\mathbf{v}_1, \dots, \mathbf{v}_4, \mathbf{u}_1, \dots, \mathbf{u}_4 \in \mathbf{Y}$ the following inequality hold*

$$\left| \mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4) \right| \leq \sum_{i=1}^4 s_i(\kappa) \|\mathbf{v}_i - \mathbf{u}_i\|$$

where $s_1, \dots, s_4 \in \mathbf{L}^1[0, 1]$ are nonnegative realvalued maps. Moreover, let $\exists t_0 \in \mathbb{N}$ such that there exists $\mathbf{Z}_1, \dots, \mathbf{Z}_{t_0} \in \mathbf{L}^1[0, 1]$ and $\mathbf{X}_1, \dots, \mathbf{X}_{t_0} : \mathbb{R}^4 \rightarrow [0, \infty)$ where \mathbf{Z}_i are nonnegative and \mathbf{X}_i are nondecreasing and nonnegative functions for all $i = 1, \dots, t_0$. Also, let $\beta_q = \min \{1, \Gamma_q(2 - \tau), \Gamma_q(\sigma + 1)\}$, $\mathcal{C}_q(\gamma, \ell) = \max \{\mathcal{A}_q(\gamma, \ell), \mathcal{B}_q(\gamma, \ell)\}$ and assume that the following are holds true

- For all $(\mathbf{v}_1, \dots, \mathbf{v}_4) \in \mathbf{Y}^4$ and almost $\kappa \in [0, 1]$

$$\left| \mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) \right| \leq \sum_{i=1}^{t_0} \mathbf{Z}_i(\kappa) \mathbf{X}_i(\mathbf{v}_1, \dots, \mathbf{v}_4).$$

- $\lim_{y \rightarrow \infty} \frac{\mathbf{X}_i(y, y, y, y)}{y} = \rho$ where $\rho \in \mathbb{R}^+$ and for some $\nu > 0$

$$0 \leq \rho \leq \frac{\beta_q}{\mathcal{C}_q(\gamma, \ell) \sum_{i=1}^{t_0} \|\mathbf{Z}_i\| + \nu}.$$

- $\mathcal{C}_q(\gamma, \ell) \left(\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right) < 1$ where $\hat{s}_i = \int_0^1 (1-qp)^{\gamma-1} s_i \, d_q p$.

Then, the quantum multi-singular integrodifferential problem mentioned in (1.1)-(1.2) has a solution.

Proof. At first, we define the operator $\mathcal{T} : \mathbf{Y} \rightarrow \mathbf{Y}$ as follows:

$$\mathcal{T}(\kappa) = \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{h}(\kappa, \mathbf{v}(\kappa), \mathbf{v}'(\kappa), {}^c \mathfrak{D}_q^\tau \mathbf{v}(\kappa), \mathfrak{I}_q^\sigma \mathbf{v}(\kappa)).$$

Now in three steps, we show that this operator has a fixed point which is the same solution to the desired problem.

step1. we shall show that \mathcal{T} is continuous. Suppose that $\mathbf{v}_1, \mathbf{v}_2 \in \mathbf{Y}$ and $\kappa \in [0, 1]$. Afterwards, we can write

$$\begin{aligned} |\mathcal{T}_{\mathbf{v}_1}(\kappa) - \mathcal{T}_{\mathbf{v}_2}(\kappa)| &\leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}_1(p), \mathbf{v}_1'(p), {}^c \mathfrak{D}_q^\tau \mathbf{v}_1(p), \mathfrak{I}_q^\sigma \mathbf{v}_1(p)) \right. \\ &\quad \left. - \mathbf{h}(p, \mathbf{v}_2(p), \mathbf{v}_2'(p), {}^c \mathfrak{D}_q^\tau \mathbf{v}_2(p), \mathfrak{I}_q^\sigma \mathbf{v}_2(p)) \right| d_q p \\ &\leq \int_0^1 \mathcal{A}_q(\gamma, \ell) (1-qp)^{\gamma-1} \left(s_1(p) \|\mathbf{v}_1 - \mathbf{v}_2\| + s_2(p) \|\mathbf{v}_1' - \mathbf{v}_2'\| \right. \\ &\quad \left. + s_3(p) \|{}^c \mathfrak{D}_q^\tau \mathbf{v}_1 - {}^c \mathfrak{D}_q^\tau \mathbf{v}_2\| + s_4(p) \|\mathfrak{I}_q^\sigma \mathbf{v}_1 - \mathfrak{I}_q^\sigma \mathbf{v}_2\| \right) d_q p. \end{aligned}$$

But,

$$|\mathfrak{I}_q^\sigma \mathbf{v}(\kappa)| \leq \frac{1}{\Gamma_q(\sigma)} \int_0^\kappa (\kappa - qp)^\sigma |\mathbf{v}(p)| d_q p \leq \frac{\|\mathbf{v}\|}{\Gamma_q(\sigma)} \times \frac{1}{\sigma} [(\kappa - qp)^\sigma]_0^\kappa = \frac{\|\mathbf{v}\|}{\Gamma_q(\sigma+1)} \kappa^\sigma,$$

which implies that $\|\mathfrak{I}_q^\sigma \mathbf{v}\| \leq \frac{\|\mathbf{v}\|}{\Gamma_q(\sigma+1)}$. Therefore,

$$\|\mathfrak{I}_q^\sigma \mathbf{v}_1 - \mathfrak{I}_q^\sigma \mathbf{v}_2\| = \|\mathfrak{I}_q^\sigma (\mathbf{v}_1 - \mathbf{v}_2)\| \leq \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(\sigma+1)}.$$

Also, similar to the above processes for all $\mathbf{v}_1, \mathbf{v}_2 \in \mathbf{C}^1[0, 1]$, we deduce that

$$\|{}^c \mathfrak{D}_q^\tau \mathbf{v}_1 - {}^c \mathfrak{D}_q^\tau \mathbf{v}_2\| \leq \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(2-\tau)}.$$

Hence,

$$\begin{aligned} &|\mathcal{T}_{\mathbf{v}_1}(\kappa) - \mathcal{T}_{\mathbf{v}_2}(\kappa)| \\ &\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 \left(s_1(p) \|\mathbf{v}_1 - \mathbf{v}_2\| + s_2(p) \|\mathbf{v}_1' - \mathbf{v}_2'\| \right. \end{aligned}$$

$$\begin{aligned}
& + s_3(p) \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(2 - \tau)} + s_4(p) \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(\sigma + 1)} \Big) (1 - qp)^{\gamma-1} d_q p \\
& = \mathcal{A}_q(\gamma, \ell) \int_0^1 \left[\left(s_1(p) + \frac{s_4(p)}{\Gamma_q(\sigma + 1)} \right) (1 - qp)^{\gamma-1} \|\mathbf{v}_1 - \mathbf{v}_2\| \right. \\
& \quad \left. + \left(s_2(p) + \frac{s_3(p)}{\Gamma_q(2 - \tau)} \right) (1 - qp)^{\gamma-1} \|\mathbf{v}'_1 - \mathbf{v}'_2\| \right] d_q p \\
& \leq \mathcal{A}_q(\gamma, \ell) \|\mathbf{v}_1 - \mathbf{v}_2\|_* \int_0^1 \left[s_1(p) + s_2(p) + \frac{s_3(p)}{\Gamma_q(2 - \tau)} + \frac{s_4(p)}{\Gamma_q(\sigma + 1)} \right] (1 - qp)^{\gamma-1} d_q p \\
& = \mathcal{A}_q(\gamma, \ell) \left(\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2 - \tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma + 1)} \right) \|\mathbf{v}_1 - \mathbf{v}_2\|_*.
\end{aligned}$$

As well as, we have

$$\begin{aligned}
|\mathcal{T}'_{\mathbf{v}_1}(\kappa) - \mathcal{T}'_{\mathbf{v}_2}(\kappa)| & \leq \int_0^1 \left| \frac{\partial \mathcal{G}_q(\kappa, p)}{\partial \kappa} \right| \left| \mathbf{h}(p, \mathbf{v}_1(p), \mathbf{v}'_1(p), {}^c \mathfrak{D}_q^\tau \mathbf{v}_1(p), \mathfrak{I}_q^\sigma \mathbf{v}_1(p)) \right. \\
& \quad \left. - \mathbf{h}(p, \mathbf{v}_2(p), \mathbf{v}'_2(p), {}^c \mathfrak{D}_q^\tau \mathbf{v}_2(p), \mathfrak{I}_q^\sigma \mathbf{v}_2(p)) \right| d_q p \\
& \leq \int_0^1 \mathcal{B}_q(\gamma, \ell) (1 - qp)^{\gamma-1} \left(s_1(p) \|\mathbf{v}_1 - \mathbf{v}_2\| + s_2(p) \|\mathbf{v}'_1 - \mathbf{v}'_2\| \right. \\
& \quad \left. + s_3(p) \|{}^c \mathfrak{D}_q^\tau \mathbf{v}_1 - {}^c \mathfrak{D}_q^\tau \mathbf{v}_2\| + s_4(p) \|\mathfrak{I}_q^\sigma \mathbf{v}_1 - \mathfrak{I}_q^\sigma \mathbf{v}_2\| \right) d_q p \\
& \leq \mathcal{B}_q(\gamma, \ell) \int_0^1 \left(s_1(p) \|\mathbf{v}_1 - \mathbf{v}_2\| + s_2(p) \|\mathbf{v}'_1 - \mathbf{v}'_2\| \right. \\
& \quad \left. + s_3(p) \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(2 - \tau)} + s_4(p) \frac{\|\mathbf{v}_1 - \mathbf{v}_2\|}{\Gamma_q(\sigma + 1)} \right) (1 - qp)^{\gamma-1} d_q p \\
& = \mathcal{B}_q(\gamma, \ell) \left(\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2 - \tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma + 1)} \right) \|\mathbf{v}_1 - \mathbf{v}_2\|_*.
\end{aligned}$$

It follows from above inequalities

$$\begin{aligned}
\|\mathcal{T}_{\mathbf{v}_1} - \mathcal{T}_{\mathbf{v}_2}\|_* & = \max \{ \|\mathcal{T}_{\mathbf{v}_1}(\kappa) - \mathcal{T}_{\mathbf{v}_2}(\kappa)\|, \|\mathcal{T}'_{\mathbf{v}_1} - \mathcal{T}'_{\mathbf{v}_2}\| \} \\
& \leq \mathcal{C}_q(\gamma, \ell) \left(\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2 - \tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma + 1)} \right) \|\mathbf{v}_1 - \mathbf{v}_2\|_*.
\end{aligned}$$

Accordingly $\|\mathcal{T}_{\mathbf{v}_1} - \mathcal{T}_{\mathbf{v}_2}\|_* \rightarrow 0$ as $\|\mathbf{v}_1 - \mathbf{v}_2\|_* \rightarrow 0$, thus \mathcal{T} is continuous.

step2. We claim that \mathcal{T} is α -admissible. At first, since

$$\rho \times \frac{\mathcal{C}_q(\gamma, \ell) \sum_{i=1}^{t_0} \|\mathbf{Z}_i\|}{\beta_q} < 1,$$

then, we can choose λ where

$$(\rho + \lambda) \frac{\mathcal{C}_q(\gamma, \ell) \sum_{i=1}^{t_0} \|\mathbf{Z}_i\|}{\beta_q} < 1.$$

According to $\frac{\mathbf{X}(y, y, y, y)}{y} \rightarrow \rho$ as $y \rightarrow \infty$, then $\exists \zeta = \nu_\lambda > 0$ such that for all $y > \zeta = \nu_\lambda$ we have

$$\frac{\mathbf{X}(y, y, y, y)}{y} < \rho + \lambda. \quad (3.1)$$

Now, let $\mathbf{W} = \{\mathbf{v} \in \mathbf{Y} : \|\mathbf{v}\|_* < \zeta\}$ and define the map $\alpha : \mathbf{Y}^2 \rightarrow [0, \infty)$ as follows:

$$\alpha(\mathbf{v}, \mathbf{u}) = \begin{cases} 1, & \mathbf{v}, \mathbf{u} \in \mathbf{W} \\ 0, & \mathbf{v}, \mathbf{u} \notin \mathbf{W}. \end{cases}$$

If $\alpha(\mathbf{v}, \mathbf{u}) \geq 1$; then, $\|\mathbf{v}\|_* \leq \zeta$ and $\|\mathbf{u}\|_* \leq \zeta$. suppose that $\kappa \in [0, 1]$, then we get

$$\begin{aligned} |\mathcal{T}_{\mathbf{v}}(\kappa)| &\leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right| d_q p \\ &\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \sum_{i=0}^{t_0} \mathbf{Z}_i(p) \mathbf{X}_i(\mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) d_q p \\ &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \int_0^1 (1 - qp)^{\gamma-1} \sum_{i=0}^{t_0} \mathbf{Z}_i(p) \mathbf{X}_i\left(\|\mathbf{v}\|, \|\mathbf{v}'\|, \frac{\|\mathbf{v}'\|}{\Gamma_q(2-\tau)}, \frac{\|\mathbf{v}\|}{\Gamma_q(1+\sigma)}\right) d_q p \\ &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \mathbf{X}_i\left(\zeta, \zeta, \frac{\zeta}{\Gamma_q(2-\tau)}, \frac{\zeta}{\Gamma_q(1+\sigma)}\right) \left(\int_0^1 \mathbf{Z}_i(p) \sup(1 - qp)^{\gamma-1} d_q p \right) \\ &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \mathbf{X}_i\left(\frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}\right) \|\mathbf{Z}_i\|. \end{aligned}$$

It follows from (3.1) and $\frac{\zeta}{\beta} > \zeta$ that $\mathbf{X}_i(\frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}) < (\rho + \lambda) \frac{\zeta}{\beta_q}$ and so

$$\begin{aligned} |\mathcal{T}_{\mathbf{v}}(\kappa)| &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \frac{\zeta}{\beta_q} (\rho + \lambda) \|\mathbf{Z}_i\| \\ &= (\rho + \lambda) \left(\frac{\mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \|\mathbf{Z}_i\|}{\beta_q} \right) \times \zeta \\ &< \zeta, \end{aligned}$$

therefore, $\|\mathcal{T}_{\mathbf{v}}\| \leq \zeta$. In addition, we can write

$$|\mathcal{T}'_{\mathbf{v}}(\kappa)| \leq \int_0^1 \left| \frac{\partial \mathcal{G}_q(\kappa, p)}{\partial \kappa} \right| \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right| d_q p$$

$$\begin{aligned}
&\leq \mathcal{B}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \sum_{i=0}^{t_0} \mathbf{Z}_i(p) \mathbf{X}_i(\mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \, d_q p \\
&\leq \mathcal{B}_q(\gamma, \ell) \sum_{i=0}^{t_0} \mathbf{X}_i\left(\zeta, \zeta, \frac{\zeta}{\Gamma_q(2-\tau)}, \frac{\zeta}{\Gamma_q(1+\sigma)}\right) \left(\int_0^1 \sup \mathbf{Z}_i(p) (1 - qp)^{\gamma-1} \, d_q p\right) \\
&\leq \mathcal{B}_q(\gamma, \ell) \sum_{i=0}^{t_0} \mathbf{X}_i\left(\frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}, \frac{\zeta}{\beta_q}\right) \|\mathbf{Z}_i\| \\
&= (\rho + \lambda) \left(\frac{\mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \|\mathbf{Z}_i\|}{\beta_q}\right) \zeta < \zeta
\end{aligned}$$

and so

$$\|\mathcal{T}_{\mathbf{v}}\|_* = \max\{\|\mathcal{T}_{\mathbf{v}}\|, \|\mathcal{T}'_{\mathbf{v}}\|\} \leq \zeta.$$

Therefore, we draw conclusions that $\mathcal{T}_{\mathbf{v}} \in \mathbf{W}$ and $\mathcal{T}_{\mathbf{u}} \in \mathbf{W}$, which implies that $\alpha(\mathcal{T}_{\mathbf{v}}, \mathcal{T}_{\mathbf{u}}) \geq 1$. hence, \mathcal{T} is α -admissible.

Step3. In view of $\mathbf{W} \neq \emptyset$, afterwards $\exists \mathbf{v}_0 \in \mathbf{W}$ such that $\mathcal{T}_{\mathbf{v}_0} \in \mathbf{W}$. Thus, $\alpha(\mathbf{v}_0, \mathcal{T}_{\mathbf{v}_0}) \geq 1$. For convenience, set

$$\mathcal{J}_q(\gamma, \ell) := \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \mathcal{C}_q(\gamma, \ell) < 1,$$

and for all $\kappa \geq 0$ put $\psi(\kappa) = \mathcal{J}_q(\gamma, \ell)\kappa$. Since

$$\sum_{n=1}^{\infty} \psi^n(\kappa) = \sum_{n=1}^{\infty} \mathcal{J}_q^n(\gamma, \ell)\kappa = \frac{\mathcal{J}_q(\gamma, \ell)}{1 - \mathcal{J}_q(\gamma, \ell)} \kappa < \infty,$$

and $\psi : [0, \infty) \rightarrow [0, \infty)$ is nondecreasing, we have $\psi \in \Psi$. Now, we obtain

$$\begin{aligned}
|\mathcal{T}_{\mathbf{v}}(\kappa) - \mathcal{T}_{\mathbf{u}}(\kappa)| &\leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right. \\
&\quad \left. - \mathbf{h}(p, \mathbf{u}(p), \mathbf{u}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{u}(p), \mathfrak{I}_q^\sigma \mathbf{u}(p)) \right| \, d_q p \\
&\leq \mathcal{A}_q(\gamma, \ell) \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \|\mathbf{v} - \mathbf{u}\|_*,
\end{aligned}$$

which yields

$$\|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\| \leq \mathcal{A}_q(\gamma, \ell) \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \|\mathbf{v} - \mathbf{u}\|_*.$$

Also, in a similar way

$$|\mathcal{T}'_{\mathbf{v}}(\kappa) - \mathcal{T}'_{\mathbf{u}}(\kappa)| \leq \int_0^1 \left| \frac{\partial \mathcal{G}_q(\kappa, p)}{\partial \kappa} \right| \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right| \, d_q p$$

$$\begin{aligned}
& - \mathbf{h}(p, \mathbf{u}(p), \mathbf{u}'(p), {}^c \mathfrak{D}_q^\tau \mathbf{u}(p), \mathfrak{I}_q^\sigma \mathbf{u}(p)) \Big| d_q p \\
& \leq \mathcal{B}_q(\gamma, \ell) \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \|\mathbf{v} - \mathbf{u}\|_*,
\end{aligned}$$

which yields

$$\|\mathcal{T}'_{\mathbf{v}} - \mathcal{T}'_{\mathbf{u}}\| \leq \mathcal{B}_q(\gamma, \ell) \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \|\mathbf{v} - \mathbf{u}\|_*.$$

Thus for all $\mathbf{v}, \mathbf{u} \in \mathbf{W}$ we obtain

$$\begin{aligned}
\|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\|_* & \leq \mathcal{C}_q(\gamma, \ell) \left[\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2-\tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma+1)} \right] \|\mathbf{v} - \mathbf{u}\|_* \\
& = \mathcal{J}_q(\gamma, \ell) \|\mathbf{v} - \mathbf{u}\|_* = \psi(\|\mathbf{v} - \mathbf{u}\|_*).
\end{aligned}$$

Hence for all $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ we get

$$\alpha(\mathbf{v}, \mathbf{u}) \|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\|_* \leq \psi(d(\mathbf{v}, \mathbf{u})).$$

In view of Lemma 2.8, the operator \mathcal{T} has a fixed point which is the solution to the problem mentioned in (1.1)-(1.2). \square

Theorem 3.2. *Assume that the function $\mathbf{h} : \mathcal{K} \times (\mathbf{C}[0, 1])^4 \rightarrow \mathbb{R}$ defined for all $\mathbf{v}_1, \dots, \mathbf{v}_4 \in \mathbf{C}[0, 1]$ and almost all $\kappa \in [0, 1]$. In addition, let $\exists t_1 \in \mathbb{N}$ and there exists nondecreasing maps $\mathbf{X}_1, \dots, \mathbf{X}_{t_1} : \mathbb{R}^4 \rightarrow \mathbb{R}$ such that $\mathbf{X}_i(y, y, y, y) \geq 0$ and $\lim_{y \rightarrow 0^+} \frac{\mathbf{X}_i(y, y, y, y)}{y} = \rho_i$ for all $y \geq 0$, $i = 1, \dots, t_1$ and some $0 \leq \rho_i < 1$. Moreover, for all $\mathbf{v}_1, \dots, \mathbf{v}_4, \mathbf{u}_1, \dots, \mathbf{u}_4 \in \mathbf{Y}$ there exists $s_1, \dots, s_{t_1} : [0, 1] \rightarrow [0, \infty)$ s.t.*

$$\begin{aligned}
& \left| \mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4) \right| \\
& \leq \sum_{i=1}^{t_1} s_i(\kappa) \mathbf{X}_i(\mathbf{v}_1 - \mathbf{u}_1, \mathbf{v}_2 - \mathbf{u}_2, \mathbf{v}_3 - \mathbf{u}_3, \mathbf{v}_4 - \mathbf{u}_4)
\end{aligned}$$

where $\mathbf{v}_j \geq \mathbf{u}_j \geq 0$. If

$$\mathcal{C}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - qp)^{\gamma-1} s_i\|_1 \leq 1.$$

Then, the quantum multi-singular integrodifferential problem mentioned in (1.1)-(1.2) has a solution.

Proof. we present our proof in three steps.

Step1. Assume that $\lambda > 0$ be given and put

$$\mathbf{q}_1 := \mathcal{C}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 + 1.$$

Moreover, according to $\mathbf{X}_i(\frac{y}{\beta}, \dots, \frac{y}{\beta}) \rightarrow 0$ as $y \rightarrow 0^+$ then for each $i = 1, \dots, t_1$ there exists $\zeta_i > 0$ such that $\mathbf{X}_i(\frac{y}{\beta}, \dots, \frac{y}{\beta}) < \frac{\lambda}{q_1}$ where $0 < y \leq \zeta_i$. If $\zeta = \min\{\zeta_i : 1 \leq i \leq t_1\}$, then for all $0 < y \leq \zeta$ we get $\mathbf{X}_i(\frac{y}{\beta}, \dots, \frac{y}{\beta}) < \frac{\lambda}{q_1}$. If $\mathbf{v}_n \rightarrow \mathbf{v}$ then $\exists n_0 \in \mathbb{N}$ such that $\forall n \geq n_0$ we have $\|\mathbf{v}_n - \mathbf{v}\|_* < \zeta$. Afterwards for all $\kappa \in [0, 1]$ and $n \geq n_0$ we have

$$\begin{aligned}
& |\mathcal{T}_{\mathbf{v}_n}(\kappa) - \mathcal{T}_{\mathbf{v}}(\kappa)| \\
& \leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \right. \\
& \quad \left. - \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right| d_q p \\
& \leq \int_0^1 \mathcal{G}_q(\kappa, p) \sum_{i=0}^{t_1} s_i(p) \mathbf{X}_i(\|\mathbf{v}_n - \mathbf{v}\|, \|(\mathbf{v}_n - \mathbf{v})'\|, \|{}^c\mathfrak{D}_q^\tau(\mathbf{v}_n - \mathbf{v})\|, \|\mathfrak{I}_q^\sigma(\mathbf{v}_n - \mathbf{v})\|) d_q p \\
& \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \mathbf{X}_i\left(\zeta, \zeta, \frac{\zeta}{\Gamma_q(2-\tau)}, \frac{\zeta}{\Gamma_q(1+\sigma)}\right) \left(\int_0^1 s_i(p) (1-qp)^{\gamma-1} d_q p \right) \\
& \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \mathbf{X}_i\left(\frac{\zeta}{\beta}, \frac{\zeta}{\beta}, \frac{\zeta}{\beta}, \frac{\zeta}{\beta}\right) \left(\int_0^1 s_i(p) (1-qp)^{\gamma-1} d_q p \right) \\
& \leq \mathcal{A}_q(\gamma, \ell) \frac{\lambda}{t_*} \sum_{i=0}^{t_1} \|(1-\kappa)^{\gamma-1} s_i\|_1 \zeta < \lambda,
\end{aligned}$$

so for $n \geq n_0$ we obtain $\|\mathcal{T}_{\mathbf{v}_n} - \mathcal{T}_{\mathbf{v}}\| < \lambda$. Also, similarly $\forall \kappa \in [0, 1]$ and $n \geq n_0$ we get

$$\begin{aligned}
& |\mathcal{T}'_{\mathbf{v}_n}(\kappa) - \mathcal{T}'_{\mathbf{v}}(\kappa)| \leq \int_0^1 \left| \frac{\partial \mathcal{G}_q(\kappa, p)}{\partial \kappa} \right| \left| \mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \right. \\
& \quad \left. - \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right| d_q p \\
& \leq \mathcal{B}_q(\gamma, \ell) \frac{\lambda}{t_*} \sum_{i=0}^{t_1} \|(1-\kappa)^{\gamma-1} s_i\|_1 < \lambda,
\end{aligned}$$

and so $\|\mathcal{T}'_{\mathbf{v}_n} - \mathcal{T}'_{\mathbf{v}}\| \leq \lambda$. Thus

$$\|\mathcal{T}'_{\mathbf{v}_n} - \mathcal{T}'_{\mathbf{v}}\|_* = \max\{\|\mathcal{T}_{\mathbf{v}_n} - \mathcal{T}_{\mathbf{v}}\|, \|\mathcal{T}'_{\mathbf{v}_n} - \mathcal{T}'_{\mathbf{v}}\|\} \leq \lambda,$$

which implies that $\mathcal{T}_{\mathbf{v}_n} \rightarrow \mathcal{T}_{\mathbf{v}}$ as $\mathbf{v}_n \rightarrow \mathbf{v}$. Therefore, \mathcal{T} is continuous.

Step2. Notice that, for all $i = 1, \dots, t_1$ we have $\lim_{y \rightarrow 0^+} \frac{\mathbf{X}_i(y, y, y, y)}{y} = \rho_i < 1$; So $\forall \lambda_i > 0$, $\exists \zeta_i := \zeta(\lambda_i) > 0$ where $0 < \frac{y}{\beta} < \zeta_i$ and it follows that

$$\mathbf{X}_i\left(\frac{y}{\beta}, \dots, \frac{y}{\beta}\right) \leq (\rho_i + \lambda_i) \frac{y}{\beta}$$

where $\beta = \min\{\Gamma_q(2 - \tau), \Gamma_q(1 + \sigma)\}$. Now assume that λ_i^0 be such that $\rho_i + \lambda_i^0 < 1$ and $\zeta_i^0 := \zeta(\lambda_i^0)$. Put:

$$\zeta = \min\{\zeta_1^0, \dots, \zeta_{t_1}^0\}, \quad \rho = \max\{\rho_1, \dots, \rho_{t_1}\} \quad \text{and} \quad \lambda_0 = \min\{\lambda_1^0, \dots, \lambda_{t_1}^0\},$$

which for all $0 < \frac{y}{\beta} < \zeta_i$ implies that

$$\rho + \lambda_0 < 1 \quad \text{and} \quad \mathbf{X}_i\left(\frac{y}{\beta}, \dots, \frac{y}{\beta}\right) \leq (\rho + \lambda_0) \frac{y}{\beta}.$$

Moreover,

$$\mathbf{X}_i\left(\frac{y}{\beta}, \dots, \frac{y}{\beta}\right) \leq (\rho + \lambda_0) \frac{\zeta}{\beta} \beta \leq (\rho + \lambda_0) \zeta \leq (\rho + \lambda_0) \lambda_0.$$

Define the map $\boldsymbol{\alpha} : \mathbf{Y}^2 \rightarrow [0, \infty)$ as follows:

$$\boldsymbol{\alpha}(\mathbf{v}, \mathbf{u}) = \begin{cases} 1, & \|\mathbf{v} - \mathbf{u}\|_* \leq \zeta \\ 0, & \text{otherwise.} \end{cases}$$

Now, let $\boldsymbol{\alpha}(\mathbf{v}, \mathbf{u}) \geq 1$ then

$$\begin{aligned} & |\mathcal{T}_{\mathbf{v}}(\kappa) - \mathcal{T}_{\mathbf{u}}(\kappa)| \\ & \leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right. \\ & \quad \left. - \mathbf{h}(p, \mathbf{u}(p), \mathbf{u}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{u}(p), \mathfrak{I}_q^\sigma \mathbf{u}(p)) \right| \mathrm{d}_q p \\ & \leq \int_0^1 \mathcal{G}_q(\kappa, p) \sum_{i=0}^{t_1} s_i(p) \left| \mathbf{X}_i((\mathbf{v} - \mathbf{u}), (\mathbf{v} - \mathbf{u})', {}^c\mathfrak{D}_q^\tau (\mathbf{v} - \mathbf{u}), \mathfrak{I}_q^\sigma (\mathbf{v} - \mathbf{u})) \right| \mathrm{d}_q p \\ & \leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \sum_{i=0}^{t_1} s_i(p) \left| \mathbf{X}_i(\|\mathbf{v} - \mathbf{u}\|, \|(\mathbf{v} - \mathbf{u})'\|, \|{}^c\mathfrak{D}_q^\tau (\mathbf{v} - \mathbf{u})\|, \|\mathfrak{I}_q^\sigma (\mathbf{v} - \mathbf{u})\|) \right| \mathrm{d}_q p \\ & \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \mathbf{X}_i\left(\zeta, \zeta, \frac{\zeta}{\Gamma_q(2 - \tau)}, \frac{\zeta}{\Gamma_q(1 + \sigma)}\right) \left(\int_0^1 s_i(p) (1 - qp)^{\gamma-1} \mathrm{d}_q p \right) \\ & \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \mathbf{X}_i\left(\frac{\zeta}{\beta}, \frac{\zeta}{\beta}, \frac{\zeta}{\beta}, \frac{\zeta}{\beta}\right) \left(\int_0^1 s_i(p) (1 - qp)^{\gamma-1} \mathrm{d}_q p \right) \\ & \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - qp)^{\gamma-1} s_i\|_1 (\rho + \lambda) \zeta \\ & \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - qp)^{\gamma-1} s_i\|_1 \zeta \leq \zeta \end{aligned}$$

so, $\|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\| \leq \zeta$ which yields that $\alpha(\mathcal{T}_{\mathbf{v}}, \mathcal{T}_{\mathbf{u}}) = 1$. Now, put

$$\mathbf{q}_2 = \mathcal{C}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 (\rho + \lambda_0)$$

then follows from assumption that $\mathbf{q}_2 < 1$. If $\psi(\kappa) = \mathbf{q}_2 \kappa$ then $\psi \in \Psi$. If $\|\mathbf{v} - \mathbf{u}\| \leq \zeta$, then we have

$$\|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\| \leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 (\rho + \lambda_0) \|\mathbf{v} - \mathbf{u}\|_* \leq \mathbf{q}_2 \|\mathbf{v} - \mathbf{u}\|_*$$

and

$$\|\mathcal{T}'_{\mathbf{v}} - \mathcal{T}'_{\mathbf{u}}\| \leq \mathcal{B}_q(\gamma, \ell) \sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 (\rho + \lambda_0) \|\mathbf{v} - \mathbf{u}\|_* \leq \mathbf{q}_2 \|\mathbf{v} - \mathbf{u}\|_*.$$

Hence, $\|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\|_* \leq \mathbf{q}_2 \|\mathbf{v} - \mathbf{u}\|_* = \psi(\|\mathbf{v} - \mathbf{u}\|_*)$ and so for all $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ we get

$$\alpha(\mathbf{v}, \mathbf{u}) \|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\|_* \leq \psi(\|\mathbf{v} - \mathbf{u}\|_*).$$

step3. Now, we claim that $\exists \mathbf{v}_0 \in \mathbf{Y}$ such that $\alpha(\mathbf{v}_0, \mathcal{T}_{\mathbf{v}_0}) = 1$. To achieve this goal we shall show that $\|\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0\| \leq \zeta$ for some $\mathbf{v}_0 \in \mathbf{Y}$. According to $\mathbf{X}_i(y, y, y, y) \rightarrow 0$ as $y \rightarrow 0^+$, then $\forall \mathcal{L} > 0$ there exists $n = n(\lambda)$ where $0 < y \leq \frac{\mathcal{L}}{n}$ which implies that $\mathbf{X}_i(y, y, y, y) \leq \lambda$ for all $1 \leq i \leq t_1$. Therefore, $\mathbf{X}_i(\frac{y}{n}, \frac{y}{n}, \frac{y}{n}, \frac{y}{n}) \leq \lambda$. Put

$$\mathcal{L} := \max \left\{ 1, \frac{1}{\Gamma_q(1 + \sigma)}, \frac{1}{\Gamma_q(2 - \tau)} \right\} = \max \left\{ \frac{1}{\Gamma_q(1 + \sigma)}, \frac{1}{\Gamma_q(2 - \tau)} \right\}$$

and take λ_m such that

$$\sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 \lambda_m \mathcal{C}_q(\gamma, \ell) < \zeta.$$

Put $\mathbf{b}_1 = n(\lambda_m)$ and take $\mathbf{b}_2 \in \mathbb{N}$ where

$$\sum_{i=0}^{t_1} \|(1 - \kappa)^{\gamma-1} s_i\|_1 \lambda_m \mathcal{C}_q(\gamma, \ell) \leq \zeta - \frac{1}{\mathbf{b}_2}.$$

Now, if $\mathbf{b}_* = \max\{\mathbf{b}_1, \mathbf{b}_2\}$ then for all $1 \leq i \leq t_1$ we have $\mathbf{X}_i(\frac{y}{\mathbf{b}_*}, \frac{y}{\mathbf{b}_*}, \frac{y}{\mathbf{b}_*}, \frac{y}{\mathbf{b}_*}) \leq \lambda_m$. Define the map \mathbf{v}_0 as follows:

$$\mathbf{v}_0(\kappa) = \begin{cases} 0, & \kappa \leq \frac{1}{\mathbf{b}_*+1}, \\ \frac{6\mathbf{b}_*^2}{6\mathbf{b}_*^2 + 5\mathbf{b}_* + 2} \left(\frac{\kappa^3}{3} - \frac{2\mathbf{b}_* + 1}{2\mathbf{b}_*(\mathbf{b}_* + 1)} \kappa^2 + \frac{\kappa}{\mathbf{b}_*(\mathbf{b}_* + 1)} \right) + \frac{1}{\mathbf{b}_* + 2}, & \frac{1}{\mathbf{b}_*+1} < \kappa < \frac{1}{\mathbf{b}_*}, \\ \frac{1}{\mathbf{b}_*}, & \kappa \geq \frac{1}{\mathbf{b}_*}, \end{cases}$$

It is clear that $\mathbf{v}_0 \in \mathbf{C}[0, 1]$ and for all $\kappa \in [0, 1]$ we have $0 \leq \mathbf{v}_0 \leq \frac{1}{\mathbf{b}_*}$ and $\mathbf{v}_0(\frac{1}{\mathbf{b}_*+1}) = 0$. In addition, we get

$$\mathbf{v}'_0(\kappa) = \begin{cases} 0, & \kappa \leq \frac{1}{\mathbf{b}_*+1}, \\ \frac{6\mathbf{b}_*^2}{6\mathbf{b}_*^2 + 5\mathbf{b}_* + 2} \left(\kappa^2 - \frac{2\mathbf{b}_* + 1}{\mathbf{b}_*(\mathbf{b}_* + 1)} \kappa + \frac{1}{\mathbf{b}_*(\mathbf{b}_* + 1)} \right), & \frac{1}{\mathbf{b}_*+1} < \kappa < \frac{1}{\mathbf{b}_*}, \\ \frac{1}{\mathbf{b}_*}, & \kappa \geq \frac{1}{\mathbf{b}_*}. \end{cases}$$

Hence, $\mathbf{v}'_0 \in \mathbf{C}[0, 1]$ and $\mathbf{v}'_0(\frac{1}{\mathbf{b}_*+1}) = \mathbf{v}'_0(\frac{1}{\mathbf{b}_*}) = 0$, which implies that $\mathbf{v}_0 \in \mathbf{C}^1[0, 1]$. Moreover, we obtain

$$\begin{aligned} \mathbf{v}'_0(\kappa) &\leq \frac{6\mathbf{b}_*^2}{6\mathbf{b}_*^2 + 5\mathbf{b}_* + 2} \left(\frac{1}{\mathbf{b}_*^2} - \frac{2\mathbf{b}_* + 1}{2\mathbf{b}_*(\mathbf{b}_* + 1)^2} + \frac{1}{\mathbf{b}_*^4(\mathbf{b}_* + 1)} \right) \\ &\leq 1 \times \frac{1}{\mathbf{b}_*} \times \left(\frac{1}{\mathbf{b}_*} - \frac{2\mathbf{b}_*}{(\mathbf{b}_* + 1)^2} + \frac{1}{\mathbf{b}_*^3(\mathbf{b}_* + 1)} \right) \leq \frac{1}{\mathbf{b}_*}, \end{aligned}$$

which yields $\|\mathbf{v}_{\mathbf{b}_*}\| \leq \frac{1}{\mathbf{b}_*}$. So, we can write

$$\begin{aligned} &|\mathcal{T}_{\mathbf{v}_0}(\kappa) - \mathbf{v}_0(\kappa)| \\ &= \left| \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \, d_q p - \mathbf{v}_0(p) \right| \\ &\leq \int_0^1 \mathcal{G}_q(\kappa, p) |\mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \, d_q p| + \frac{1}{\mathbf{b}_*} \\ &\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \sum_{i=0}^{t_0} s_i(p) \mathbf{X}_i \left(\|\mathbf{v}_0\|, \|\mathbf{v}_0'\|, \frac{\|\mathbf{v}_0'\|}{\Gamma_q(2-\tau)}, \frac{\|\mathbf{v}_0\|}{\Gamma_q(1+\sigma)} \right) \, d_q p + \frac{1}{\mathbf{b}_*} \\ &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \left[\left(\int_0^1 (1 - qp)^{\gamma-1} s_i(p) \, d_q p \right) \mathbf{X}_i \left(\frac{\mathcal{L}}{\mathbf{b}_*}, \frac{\mathcal{L}}{\mathbf{b}_*}, \frac{\mathcal{L}}{\mathbf{b}_*}, \frac{\mathcal{L}}{\mathbf{b}_*} \right) \right] + \frac{1}{\mathbf{b}_*} \\ &\leq \mathcal{A}_q(\gamma, \ell) \sum_{i=0}^{t_0} \|(1 - \kappa)^{\gamma-1} s_i\| \lambda_1 + \frac{1}{\mathbf{b}_*} \leq \zeta \end{aligned}$$

and so $\|\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0\| \leq \zeta$. In a same way

$$\begin{aligned} |\mathcal{T}'_{\mathbf{v}_0}(\kappa) - \mathbf{v}'_0(\kappa)| &\leq \int_0^1 \left| \frac{\partial \mathcal{G}_q(\kappa, p)}{\partial \kappa} \right| |\mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \, d_q p| + \frac{1}{\mathbf{b}_*} \\ &\leq \mathcal{B}_q(\gamma, \ell) \sum_{i=0}^{t_0} \|(1 - \kappa)^{\gamma-1} s_i\| \lambda_1 + \frac{1}{\mathbf{b}_*} \leq \zeta. \end{aligned}$$

It follows $\|(\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0)'\| \leq \zeta$. Thus,

$$\|\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0\|_* = \max \{ \|\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0\|, \|(\mathcal{T}_{\mathbf{v}_0} - \mathbf{v}_0)'\| \} \leq \lambda,$$

which implies that $\alpha(\mathbf{v}_0, \mathcal{T}_{\mathbf{v}_0}) = 1$. Thanks to Lemma 2.8, the multi-singular problem mentioned in (1.1)-(1.2) has a solution. \square

Theorem 3.3. *Let \mathbf{E}^c be a null subset of $[0, 1]$, namely its measure is zero, and for all $\mathbf{v}_1, \dots, \mathbf{v}_4 \in \mathbf{Y}$ and $\kappa \in \mathbf{E}$ the function $\mathbf{h} : [0, 1] \times \mathbf{Y}^4 \rightarrow [0, \infty)$ be such that $\mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) < \infty$ and continuous with respect to $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$. Moreover, let $\exists t_1 \in \mathbb{N}$ such that there exists maps $s_i : [0, 1] \rightarrow [0, \infty)$, $s_1, \dots, s_{t_1} \in \mathbf{L}^1[0, 1]$ and $\mathbf{X}_1, \dots, \mathbf{X}_{t_1} : \mathbb{R}^4 \rightarrow [0, \infty)$, $\mathbf{S} : \mathbb{R}^4 \rightarrow [0, \infty)$ such that*

$$\|\mathbf{S}\|_1^* := \sup_{\mathbf{v} \in \mathbf{C}[0,1]} \int_0^1 \mathbf{S}(\mathbf{v}(\kappa), \mathbf{v}(\kappa), \mathbf{v}(\kappa), \mathbf{v}(\kappa)) \, d_q \kappa < \infty$$

and

$$\|\mathbf{X}_i\|_\infty := \sup_{y \in \mathbb{R}} \{ \mathbf{X}_i(y, y, y, y) \} < \infty,$$

also

$$|\mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4)| \leq \sum_{i=1}^{t_1} s_i(\kappa) \mathbf{X}_i(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) + \mathbf{S}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4).$$

In addition, let there exists maps $\psi : \mathbb{R}^4 \rightarrow [0, \infty)$, s.t $\|\psi\|^* := \min\{\psi(\mathbf{v}_1, \dots, \mathbf{v}_4)\}$ and $z : [0, 1] \rightarrow [0, \infty)$ where

$$\|z\|_1^{\mathbf{L}} := \int_0^1 (1 - \kappa)^{\gamma-1} z(\kappa) \, d_q \kappa$$

and for all $\mathbf{v}_1, \dots, \mathbf{v}_4 \in \mathbf{Y}$ and almost $\kappa \in [0, 1]$

$$\mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) \geq z(\kappa) \psi(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4).$$

Also, let there exists $g_1, g_2, g_3, g_4 \in \mathbf{L}^1[0, 1]$ where

$$\sum_{i=1}^4 \|g_i\|_1 < \frac{1}{\mathcal{C}_q(\gamma, \ell)},$$

and $\Xi : [0, \infty) \rightarrow [0, \infty)$, $\Xi_\beta \in \Psi$ such that for all $\lambda > 0$ we set $\Xi_\lambda(\mathbf{v}) := \Xi(\frac{\mathbf{v}}{\lambda})$ and

$$|\mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4)| \leq \sum_{i=0}^4 g_i(\kappa) \Xi(\|\mathbf{v}_i - \mathbf{u}_i\|)$$

with $\|\mathbf{v}_i\|, \|\mathbf{u}_i\| \in [\zeta_1, \zeta_2]$ where

$$\zeta_1 \leq \frac{\|\psi\|^* \|z\|_1^{\mathbf{L}} (4 - \gamma^2 - 2\gamma)}{2\Gamma_q(\gamma)(1 - \gamma)} \quad \text{and} \quad \zeta_2 \geq \mathcal{C}_q(\gamma, \ell) \left[\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right].$$

Then, the quantum pointwise multi-singular integrodifferential problem mentioned in (1.1)-(1.2) has a solution.

Proof. Assume that $\{\mathbf{v}_n\}$ be a sequence in \mathbf{Y} such that $\|\mathbf{v}_n - \mathbf{v}\|_* \rightarrow 0$, which implies that $\mathbf{v}_n \rightarrow \mathbf{v}$ and $\mathbf{v}'_n \rightarrow \mathbf{v}'$. in view of the following inequalities:

$$\|{}^c\mathfrak{D}_q^\tau(\mathbf{v}_n - \mathbf{v})\| \leq \frac{\|(\mathbf{v}_n - \mathbf{v})'\|}{\Gamma_q(2 - \tau)} \quad \text{and} \quad \|\mathfrak{I}_q^\sigma(\mathbf{v}_n - \mathbf{v})\| \leq \frac{\|\mathbf{v}_n - \mathbf{v}\|}{\Gamma_q(1 + \sigma)}$$

we obtain ${}^c\mathfrak{D}_q^\tau \mathbf{v}_n \rightarrow {}^c\mathfrak{D}_q^\tau \mathbf{v}$ and $\mathfrak{I}_q^\sigma \mathbf{v}_n \rightarrow \mathfrak{I}_q^\sigma \mathbf{v}$. It follows from the continuity of $\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4)$ that $\mathbf{h}(\kappa, \mathbf{v}_n, \mathbf{v}'_n, {}^c\mathfrak{D}_q^\tau \mathbf{v}_n, \mathfrak{I}_q^\sigma \mathbf{v}_n) \rightarrow \mathbf{h}(\kappa, \mathbf{v}, \mathbf{v}', {}^c\mathfrak{D}_q^\tau \mathbf{v}, \mathfrak{I}_q^\sigma \mathbf{v})$. suppose that $\mathbf{v} \rightarrow \mathbf{Y}$ and $\kappa \in \mathcal{K} = [0, 1]$ then, we can write

$$|\mathcal{T}_\mathbf{v}(\kappa)| \leq \mathcal{A}_q(\gamma, \ell) \left[\int_0^1 (1 - qp)^{\gamma-1} \left(\sum_{i=1}^{t_1} s_i(p) \mathbf{X}_i(\mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right. \right. \\ \left. \left. + \mathbf{S}(\mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right) \mathrm{d}_q p \right].$$

Put, $\mathbf{v}_*(p) := \max \{ \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p) \}$, then $\mathbf{v}_* \in \mathbf{C}[0, 1]$ and afterwards for all $\kappa \in [0, 1]$ we have

$$|\mathcal{T}_\mathbf{v}(\kappa)| \leq \mathcal{A}_q(\gamma, \ell) \left[\sum_{i=1}^{t_1} \mathbf{X}_i \left(\|\mathbf{v}\|, \|\mathbf{v}'\|, \frac{\|\mathbf{v}'\|}{\Gamma_q(2 - \tau)}, \frac{\|\mathbf{v}\|}{\Gamma_q(1 + \sigma)} \right) \right] \times \int_0^1 s_i(p) \mathrm{d}_q p \\ + \int_0^1 \mathbf{S}(\mathbf{v}_*(p), \mathbf{v}_*(p), \mathbf{v}_*(p), \mathbf{v}_*(p)) \mathrm{d}_q p \\ \leq \mathcal{A}_q(\gamma, \ell) \left(\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right).$$

In a same way, we get

$$|\mathcal{T}_{\mathbf{v}'}(\kappa)| \leq \mathcal{B}_q(\gamma, \ell) \left(\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right).$$

Hence, for all $\mathbf{v} \in \mathbf{Y}$ we obtain

$$\|\mathcal{T}_\mathbf{v}\|_* \leq \mathcal{C}_q(\gamma, \ell) \left(\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right) < \infty. \quad (3.2)$$

In view of the Lebesgue dominated convergence theorem, for all $\kappa \in [0, 1]$ we deduce that

$$\mathcal{T}_{\mathbf{v}_n}(\kappa) = \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{h}(p, \mathbf{v}_n(p), \mathbf{v}'_n(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}_n(p), \mathfrak{I}_q^\sigma \mathbf{v}_n(p)) \mathrm{d}_q p \\ \rightarrow \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \mathrm{d}_q p = \mathcal{T}_\mathbf{v}(\kappa).$$

Thus \mathcal{T} is continuous.

Define the map $\alpha : \mathbf{Y}^2 \rightarrow [0, \infty)$ as follows:

$$\alpha(\mathbf{v}, \mathbf{u}) = \begin{cases} 1, & \|\mathbf{v}\|_*, \|\mathbf{u}\|_* \in [\zeta_1, \zeta_2] \\ 0, & \text{otherwise.} \end{cases}$$

Assume that $\alpha(\mathbf{v}, \mathbf{u}) \geq 1$, then $\|\mathbf{v}\|_*, \|\mathbf{u}\|_* \in [\zeta_1, \zeta_2]$ and afterwards for all $\kappa \in [0, 1]$ we can write

$$\begin{aligned} |\mathcal{T}_{\mathbf{v}}(\kappa)| &= \left| \int_0^1 \mathcal{G}_q(\kappa, p) \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \, d_q p \right| \\ &\geq \frac{1}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} \left(\frac{2 - \ell^2}{2(1 - \ell)} - \kappa \right) z(p) \\ &\quad \times \psi(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \, d_q p \\ &\geq \|\psi\|^* \left[\frac{-\kappa}{\Gamma_q(\gamma)} \int_0^1 (1 - qp)^{\gamma-1} z(p) \, d_q p \right. \\ &\quad \left. + \frac{2 - \ell^2}{2(1 - \ell)} \int_0^1 (1 - qp)^{\gamma-1} z(p) \, d_q p \right] \\ &\geq \|\psi\|^* \|z\|_1^{\mathbf{L}} \left(\frac{1}{\Gamma_q(\gamma)} + \frac{2 - \ell^2}{2(1 - \ell)} \right) \\ &= \|\psi\|^* \|z\|_1^{\mathbf{L}} \left(\frac{4 - \ell^2 - 2\ell}{2\Gamma_q(\gamma)(1 - \ell)} \right), \end{aligned}$$

which implies that

$$\|\mathcal{T}_{\mathbf{v}}\| \geq \frac{\|\psi\|^* \|z\|_1^{\mathbf{L}} (4 - \ell^2 - 2\ell)}{2\Gamma_q(\gamma)(1 - \ell)},$$

and

$$\|\mathcal{T}_{\mathbf{v}}\|_* := \{\|\mathcal{T}_{\mathbf{v}}\|, \|\mathcal{T}_{\mathbf{v}}'\|\} \geq \frac{\|\psi\|^* \|z\|_1^{\mathbf{L}} (4 - \ell^2 - 2\ell)}{2\Gamma_q(\gamma)(1 - \ell)} \geq \zeta_1.$$

It follows from Eq. (3.2)

$$\|\mathcal{T}_{\mathbf{v}}\|_* \leq \mathcal{C}_q(\gamma, \ell) \left(\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right) \leq \zeta_2,$$

therefore, $\alpha(\mathcal{T}_{\mathbf{v}}, \mathcal{T}_{\mathbf{u}}) \geq 1$. On the other hands, if $\mathbf{v}_0 \in [\zeta_1, \zeta_2]$ it is obviously $\alpha(\mathbf{v}_0, \mathcal{T}_{\mathbf{v}_0}) \geq 1$. Now, let $\mathbf{v}, \mathbf{u} \in [\zeta_1, \zeta_2]$, then we have

$$\alpha(\mathbf{v}, \mathbf{u}) |\mathcal{T}_{\mathbf{v}}(\kappa) - \mathcal{T}_{\mathbf{u}}(\kappa)|$$

$$\begin{aligned}
&\leq \int_0^1 \mathcal{G}_q(\kappa, p) \left| \mathbf{h}(p, \mathbf{v}(p), \mathbf{v}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{v}(p), \mathfrak{I}_q^\sigma \mathbf{v}(p)) \right. \\
&\quad \left. - \mathbf{h}(p, \mathbf{u}(p), \mathbf{u}'(p), {}^c\mathfrak{D}_q^\tau \mathbf{u}(p), \mathfrak{I}_q^\sigma \mathbf{u}(p)) \right| d_q p \\
&\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \left[g_1(p) \Xi(|\mathbf{v} - \mathbf{u}|) + g_2(p) \Xi(|\mathbf{v}' - \mathbf{u}'|) \right. \\
&\quad \left. + g_3(p) \Xi(|{}^c\mathfrak{D}_q^\tau \mathbf{v} - {}^c\mathfrak{D}_q^\tau \mathbf{u}|) + g_4(p) \Xi(|\mathfrak{I}_q^\sigma \mathbf{v} - \mathfrak{I}_q^\sigma \mathbf{u}|) \right] d_q p \\
&\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \left[g_1(p) \Xi(\|\mathbf{v} - \mathbf{u}\|) + g_2(p) \Xi(\|\mathbf{v}' - \mathbf{u}'\|) \right. \\
&\quad \left. + g_3(p) \Xi\left(\frac{\|\mathbf{v}' - \mathbf{u}'\|}{\Gamma_q(2 - \tau)}\right) + g_4(p) \Xi\left(\frac{\|\mathbf{v} - \mathbf{u}\|}{\Gamma_q(\sigma + 1)}\right) \right] d_q p \\
&\leq \mathcal{A}_q(\gamma, \ell) \int_0^1 (1 - qp)^{\gamma-1} \Xi\left(\frac{\|\mathbf{v} - \mathbf{u}\|_*}{\beta}\right) \sum_{i=1}^4 g_i(p) d_q p \\
&\leq \mathcal{A}_q(\gamma, \ell) \Xi\left(\frac{\|\mathbf{v} - \mathbf{u}\|_*}{\beta}\right) \sum_{i=1}^4 \int_0^1 (1 - qp)^{\gamma-1} g_i(p) d_q p \\
&\leq \sum_{i=1}^4 \|g_i\|_1 \Xi\left(\frac{\|\mathbf{v} - \mathbf{u}\|_*}{\beta}\right) \\
&\leq \Xi_\beta(\|\mathbf{v} - \mathbf{u}\|_*)
\end{aligned}$$

which for all $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ implies that

$$\alpha(\mathbf{v}, \mathbf{u}) \|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\| \leq \Xi_\beta(\|\mathbf{v} - \mathbf{u}\|_*).$$

Moreover, for all $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ we get

$$\alpha(\mathbf{v}, \mathbf{u}) \|\mathcal{T}'_{\mathbf{v}} - \mathcal{T}'_{\mathbf{u}}\| \leq \Xi_\beta(\|\mathbf{v} - \mathbf{u}\|_*),$$

Hence for all $\mathbf{v}, \mathbf{u} \in \mathbf{Y}$ we have

$$\alpha(\mathbf{v}, \mathbf{u}) \|\mathcal{T}_{\mathbf{v}} - \mathcal{T}_{\mathbf{u}}\|_* \leq \Xi_\beta(\|\mathbf{v} - \mathbf{u}\|_*).$$

Thanks to Lemma 2.8, the operator \mathcal{T} has a fixed point which is a solution for our quantum multi-singular problem mentioned in (1.1)-(1.2). \square

4 Examples

Example 4.1. *Regard the following quantum multi-singular fractional problem*

$${}^c\mathfrak{D}_q^{\frac{15}{4}} \mathbf{v}(\kappa) + \frac{1}{75\sqrt{\pi}(\mathbf{f}(\kappa))^{\frac{1}{5}}} + \left(\|\mathbf{v}\| + \|\mathbf{v}'\| + \|{}^c\mathfrak{D}_q^{\frac{4}{5}} \mathbf{v}\| + \|\mathfrak{I}_q^{\frac{3}{8}} \mathbf{v}\| \right) = 0, \quad (4.1)$$

under boundary conditions

$$\begin{cases} \mathbf{v}'(0) = \mathbf{v}(\frac{2}{3}), \\ \mathbf{v}(1) = \int_0^{\frac{1}{2}} \mathbf{v}(p) \, dp, \\ \mathbf{v}''(0) = 0, \end{cases} \quad (4.2)$$

such that

$$\mathbf{f}(\kappa) = \begin{cases} 0, & \kappa \in [0, 1] \cap Q, \\ \kappa, & \kappa \in (0, 1) \cap Q^c. \end{cases}$$

In this case put: $\gamma = \frac{15}{4}$, $\tau = \frac{4}{5}$, $\sigma = \frac{3}{8}$, $\omega = \frac{2}{3}$, $\ell = \frac{1}{2}$. Also, $s_1(\kappa) = s_2(\kappa) = s_3(\kappa) = s_4(\kappa) = \mathbf{Z}_1(\kappa) = \mathbf{Z}_2(\kappa) = \mathbf{Z}_3(\kappa) = \mathbf{Z}_4(\kappa) = \frac{1}{\kappa^{\frac{1}{5}}}$ which implies that $\|s\|_1 = \|\mathbf{Z}\|_1 = \frac{1}{60\sqrt{\pi}}$, and for $i = 1, \dots, 4$ we set $\mathbf{X}_i(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) = \|\mathbf{v}_i\|$. Then we have

$$\mathcal{A}_q(\gamma, \ell) = \mathcal{A}_q(\frac{15}{4}, \frac{1}{2}) = \frac{3}{(1 - \frac{1}{2})\Gamma_q(\frac{15}{4})} = \frac{3}{\frac{1}{2}\Gamma_q(\frac{15}{4})},$$

and

$$\mathcal{B}_q(\gamma, \ell) = \mathcal{B}_q(\frac{15}{4}, \frac{1}{2}) = \frac{2}{(1 - \frac{1}{2})\Gamma_q(\frac{15}{4} - 1)} = \frac{2}{\frac{1}{2}\Gamma_q(\frac{11}{4})},$$

hence

$$\mathcal{C}_q(\gamma, \ell) = \mathcal{C}_q(\frac{15}{4}, \frac{1}{2}) = \max \left\{ \mathcal{A}_q(\frac{15}{4}, \frac{1}{2}), \mathcal{B}_q(\frac{15}{4}, \frac{1}{2}) \right\} = \max \left\{ \frac{3}{\frac{1}{2}\Gamma_q(\frac{15}{4})}, \frac{2}{\frac{1}{2}\Gamma_q(\frac{11}{4})} \right\},$$

which their numerical values are presented in Table 3. Also, the heatmap of Table 3 is presented in Figure 3.

	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$\mathcal{A}_q(\frac{15}{4}, \frac{1}{2})$	5.2212	4.4789	3.4456	2.6915	2.1063	1.7210
$\mathcal{B}_q(\frac{15}{4}, \frac{1}{2})$	3.7799	3.5700	3.2659	3.0235	2.8126	2.6559
$\mathcal{C}_q(\frac{15}{4}, \frac{1}{2})$	5.2212	4.4789	3.4456	3.0235	2.8126	2.6559

Table 3: Numerical resual for $\mathcal{A}_q(\frac{15}{4}, \frac{1}{2})$, $\mathcal{B}_q(\frac{15}{4}, \frac{1}{2})$ and $\mathcal{C}_q(\frac{15}{4}, \frac{1}{2})$.




Figure 3: The heatmap of Table 3.

Moreover,

$$\beta_q = \min \left\{ \Gamma_q(2 - \tau), \Gamma_q(1 + \sigma) \right\} = \min \left\{ \Gamma_q\left(\frac{6}{5}\right), \Gamma_q\left(\frac{11}{8}\right) \right\},$$

and its numerical values are given in Table 4.

	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$\Gamma_q(\frac{6}{5})$	0.9802	0.9679	0.9529	0.9419	0.9326	0.9257
$\Gamma_q(\frac{11}{8})$	0.9754	0.9587	0.9379	0.9225	0.9094	0.8996
β_q	0.9802	0.9679	0.9529	0.9419	0.9326	0.9257

Table 4: Numerical resual for $\Gamma_q(\frac{6}{5})$, $\Gamma_q(\frac{11}{8})$ and β_q .

Consider the map \mathbf{h} as follows:

$$\mathbf{h}(\kappa, \mathbf{v}(\kappa), \mathbf{v}'(\kappa), {}^c\mathfrak{D}_q^\tau \mathbf{v}(\kappa), \mathfrak{I}_q^\sigma \mathbf{v}(\kappa)) := \frac{1}{75\sqrt{\pi}(\mathbf{f}(\kappa))^{\frac{1}{5}}} + \left(\|\mathbf{v}\| + \|\mathbf{v}'\| + \|{}^c\mathfrak{D}_q^{\frac{4}{5}} \mathbf{v}\| + \|\mathfrak{I}_q^{\frac{3}{8}} \mathbf{v}\| \right),$$

then we can write

$$\left| \mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4) \right| = \frac{\|\mathbf{v}_1\| - \|\mathbf{u}_1\| + \cdots + \|\mathbf{v}_4\| - \|\mathbf{u}_4\|}{75\sqrt{\pi}(\mathbf{f}(\kappa))^{\frac{1}{5}}}$$

$$\begin{aligned}
&\leq \frac{\|\mathbf{v}_1 - \mathbf{u}_1\| + \cdots + \|\mathbf{v}_4 - \mathbf{u}_4\|}{75\sqrt{\pi}(\mathbf{f}(\kappa))^{\frac{1}{5}}} \\
&= \frac{1}{75\sqrt{\pi}\kappa^{\frac{1}{5}}} \sum_{i=1}^4 \|\mathbf{v}_i - \mathbf{u}_i\| \\
&= \sum_{i=1}^4 s_i(\kappa) \|\mathbf{v}_i - \mathbf{u}_i\|,
\end{aligned}$$

and

$$\begin{aligned}
|\mathbf{h}(\kappa, \mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4)| &= \frac{\|\mathbf{v}_1\| + \cdots + \|\mathbf{v}_4\|}{75\sqrt{\pi}(\mathbf{f}(\kappa))^{\frac{1}{5}}} \\
&= \frac{\|\mathbf{v}_1\| + \cdots + \|\mathbf{v}_4\|}{75\sqrt{\pi}\kappa^{\frac{1}{5}}} \\
&= \sum_{i=1}^4 \mathbf{Z}(\kappa) \mathbf{X}_i(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4).
\end{aligned}$$

In addition, we have $\lim_{y \rightarrow \infty} \frac{\mathbf{X}(y, y, y, y)}{y} = 1 = \rho$ and

$$\mathfrak{A}_q := \frac{\beta_q}{\mathcal{C}_q(\gamma, \ell) \sum_{i=1}^4 \|\mathbf{Z}\|_1} = \frac{\beta_q}{\mathcal{C}_q(\frac{15}{4}, \frac{1}{2}) \times \frac{4}{60\sqrt{\pi}}} > 1 \quad (4.3)$$

where the establishment of inequality (4.3) is shown in Table 5.

q	$\mathfrak{A}_q > 1$
0.08	1.5888
0.17	1.8288
0.33	2.3382
0.49	2.6364
0.66	2.8061
0.81	2.9497

Table 5: Numerical resual for \mathfrak{A}_q .

Now, take $\nu > 0$ such that

$$\frac{\beta_q}{\mathcal{C}_q(\frac{15}{4}, \frac{1}{2}) \times \frac{1}{15\sqrt{\pi}} + \nu} \geq 1$$

On other hands,

$$\hat{s} = \int_0^1 (1 - qp)^{\frac{15}{4}-1} s(p) \, d_q p \leq \int_0^1 s(p) \, d_q p = \|s\|_1 = \frac{1}{60\sqrt{\pi}},$$

which implies that

$$\begin{aligned} \mathfrak{A}_q^* &= \mathcal{C}_q(\gamma, \ell) \left(\hat{s}_1 + \hat{s}_2 + \frac{\hat{s}_3}{\Gamma_q(2 - \tau)} + \frac{\hat{s}_4}{\Gamma_q(\sigma + 1)} \right) \\ &= \mathcal{C}_q\left(\frac{15}{4}, \frac{1}{2}\right) \left(\hat{s} + \hat{s} + \frac{\hat{s}}{\Gamma_q(\frac{6}{5})} + \frac{\hat{s}}{\Gamma_q(\frac{11}{8})} \right) \\ &\leq \mathcal{C}_q\left(\frac{15}{4}, \frac{1}{2}\right) \left(1 + 1 + \frac{1}{\Gamma_q(\frac{6}{5})} + \frac{1}{\Gamma_q(\frac{11}{8})} \right) \frac{1}{60\sqrt{\pi}} < 1 \end{aligned}$$

and the data related to the correctness of this inequality are given in Table 6. By using Theorem 3.1 our problem which formulated in (4.1)-(4.2) has a solution.

q	$\mathfrak{A}_q^* < 1$
0.08	0.6240
0.17	0.5393
0.33	0.4189
0.49	0.3703
0.66	0.3466
0.81	0.3289

Table 6: Numerical resual for \mathfrak{A}_q^* .

Example 4.2. Regard the following quantum multi-singular fractional problem

$${}^c \mathfrak{D}_q^{\frac{13}{5}} \mathbf{v}(\kappa) + \frac{0.06}{\kappa^{\frac{1}{3}} \left(\kappa - \frac{1}{4} \right)^{\frac{1}{6}}} \left(1 - \left(\frac{4}{5} \right)^{\frac{2}{5}} \left(\mathbf{v} + \mathbf{v}' + {}^c \mathfrak{D}_q^{\frac{3}{20}} \mathbf{v} + \mathfrak{I}_q^{\frac{4}{7}} \mathbf{v} \right) \right) = 0, \quad (4.4)$$

under boundary conditions

$$\begin{cases} \mathbf{v}'(0) = \mathbf{v}(\frac{3}{4}), \\ \mathbf{v}(1) = \int_0^{\frac{1}{8}} \mathbf{v}(p) \, dp. \end{cases} \quad (4.5)$$

In this case put: $\gamma = \frac{13}{5}$, $\tau = \frac{3}{20}$, $\sigma = \frac{4}{7}$, $\omega = \frac{3}{4}$, $\ell = \frac{1}{8}$ and $t_1 = 1$. Then we have

$$\mathcal{A}_q(\gamma, \ell) = \mathcal{A}_q\left(\frac{13}{5}, \frac{1}{8}\right) = \frac{3}{(1 - \frac{1}{8})\Gamma_q(\frac{13}{5})} = \frac{3}{\frac{7}{8}\Gamma_q(\frac{13}{5})},$$

and

$$\mathcal{B}_q(\gamma, \ell) = \mathcal{B}_q\left(\frac{13}{5}, \frac{1}{8}\right) = \frac{2}{(1 - \frac{1}{8})\Gamma_q(\frac{13}{5} - 1)} = \frac{2}{\frac{7}{8}\Gamma_q(\frac{8}{5})},$$

hence

$$\mathcal{C}_q(\gamma, \ell) = \mathcal{C}_q\left(\frac{13}{5}, \frac{1}{8}\right) = \max\left\{\mathcal{A}_q\left(\frac{13}{5}, \frac{1}{8}\right), \mathcal{B}_q\left(\frac{13}{5}, \frac{1}{8}\right)\right\} = \max\left\{\frac{3}{\frac{7}{8}\Gamma_q(\frac{13}{5})}, \frac{2}{\frac{7}{8}\Gamma_q(\frac{8}{5})}\right\},$$

which their numerical values are presented in Table 7. Also, the heatmap of Table 7 is presented in Figure 4.

	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$\mathcal{A}_q(\frac{13}{5}, \frac{1}{8})$	3.2791	3.1378	2.9333	2.7692	2.6250	2.5165
$\mathcal{B}_q(\frac{13}{5}, \frac{1}{8})$	2.3344	2.3724	2.4235	2.4638	2.4996	2.5271
$\mathcal{C}_q(\frac{13}{5}, \frac{1}{8})$	3.2791	3.1378	2.9333	2.7692	2.6250	2.5271

Table 7: Numerical resual for $\mathcal{A}_q(\frac{13}{5}, \frac{1}{8})$, $\mathcal{B}_q(\frac{13}{5}, \frac{1}{8})$ and $\mathcal{C}_q(\frac{13}{5}, \frac{1}{8})$ in Example 4.2.




Figure 4: The heatmap of Table 7 in Example 4.2.

Consider the map \mathbf{h} as follows:

$$\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) := \frac{0.06}{\kappa^{\frac{1}{3}}(\kappa - \frac{1}{4})^{\frac{1}{6}}} \left(1 - \left(\frac{4}{5}\right)^{\frac{2}{5}(\mathbf{v}_1 + \dots + \mathbf{v}_4)} \right),$$

further,

$$s(\kappa) = \frac{0.06}{\kappa^{\frac{1}{3}}(\kappa - \frac{1}{4})^{\frac{1}{6}}} \quad \text{and} \quad \mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4) = 1 - \left(\frac{4}{5}\right)^{\frac{2}{5}(\mathbf{v}_1 + \dots + \mathbf{v}_4)}.$$

It is obvious that for all $y \geq 0$ we have $\mathbf{X}(y, y, y, y) \geq 0$ and the function \mathbf{X} is nondecreasing. Now, assume that for $i = 1, \dots, 4$ we have $0 \leq \mathbf{u}_i \leq \mathbf{v}_i$ such that $(\mathbf{v}_1, \dots, \mathbf{v}_4), (\mathbf{u}_1, \dots, \mathbf{u}_4) \in \mathbf{Y}^4$. Since

$$\left(\frac{4}{5}\right)^{\mathbf{u}_i} \leq \left(\frac{4}{5}\right)^{\mathbf{v}_i},$$

then

$$\left(\frac{4}{5}\right)^{\mathbf{u}_i} \left(\left(\frac{4}{5}\right)^{\mathbf{u}_i} - \left(\frac{4}{5}\right)^{\mathbf{v}_i} \right) \leq \left(\frac{4}{5}\right)^{\mathbf{u}_i} - \left(\frac{4}{5}\right)^{\mathbf{v}_i},$$

which implies that

$$\left(\frac{4}{5}\right)^{\mathbf{u}_i} \left(\left(\frac{4}{5}\right)^{\mathbf{u}_i} - \left(\frac{4}{5}\right)^{\mathbf{v}_i} \right) \leq \left(\frac{4}{5}\right)^{\mathbf{u}_i} \left(1 - \left(\frac{4}{5}\right)^{\mathbf{v}_i - \mathbf{u}_i} \right).$$

Hence,

$$\left(1 - \left(\frac{4}{5}\right)^{\mathbf{v}_i} \right) - \left(1 - \left(\frac{4}{5}\right)^{\mathbf{u}_i} \right) \leq 1 - \left(\frac{4}{5}\right)^{\mathbf{v}_i - \mathbf{u}_i}.$$

From the placement of \mathbf{v}_i and \mathbf{u}_i with $\frac{2}{5} \sum_{i=1}^4 \mathbf{v}_i$ and $\frac{2}{5} \sum_{i=1}^4 \mathbf{u}_i$, follows that:

$$\left(1 - \left(\frac{4}{5}\right)^{\frac{2}{5} \sum_{i=1}^4 \mathbf{v}_i} \right) - \left(1 - \left(\frac{4}{5}\right)^{\frac{2}{5} \sum_{i=1}^4 \mathbf{u}_i} \right) \leq 1 - \left(\frac{4}{5}\right)^{\frac{2}{5} \sum_{i=1}^4 \mathbf{v}_i - \mathbf{u}_i}.$$

Thus,

$$\mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4) - \mathbf{X}(\mathbf{u}_1, \dots, \mathbf{u}_4) \leq \mathbf{X}(\mathbf{v}_1 - \mathbf{u}_1, \dots, \mathbf{v}_4 - \mathbf{u}_4)$$

as well as

$$\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \dots, \mathbf{u}_4) \leq s(\kappa) \mathbf{X}(\mathbf{v}_1 - \mathbf{u}_1, \dots, \mathbf{v}_4 - \mathbf{u}_4).$$

and

$$\lim_{y \rightarrow 0^+} \frac{\mathbf{X}(y, y, y, y)}{y} = \lim_{y \rightarrow 0^+} \frac{1 - \left(\frac{4}{5}\right)^{4 \times \frac{2}{5} y}}{y} = -4 \frac{2}{5} \ln \frac{4}{5} = 0.3570 < 1$$

Now all the assumptions of Theorem 3.2 are satisfied, so our problem, namely (4.4)-(4.5), has a solution.

Example 4.3. Regard the following quantum multi-singular fractional problem

$${}^c \mathfrak{D}_q^{\frac{17}{6}} \mathbf{v}(\kappa) + \frac{0.02}{\mathbf{f}(\kappa)} \mathbf{X}(\mathbf{v}(\kappa) + \mathbf{v}'(\kappa) + {}^c \mathfrak{D}_q^{\frac{7}{12}} \mathbf{v}(\kappa) + \mathfrak{I}_q^{\frac{11}{13}} \mathbf{v}(\kappa)) + 2 = 0, \quad (4.6)$$

under boundary conditions

$$\begin{cases} \mathbf{v}'(0) = \mathbf{v}(\frac{2}{5}), \\ \mathbf{v}(1) = \int_0^{\frac{2}{9}} \mathbf{v}(p) \, dp. \end{cases} \quad (4.7)$$

such that

$$\mathbf{f}(\kappa) = \begin{cases} 0, & \kappa \in [0, 1] \cap Q, \\ \sqrt{\kappa}, & \kappa \in (0, 1) \cap Q^c. \end{cases}$$

In this case put: $\gamma = \frac{17}{6}$, $\tau = \frac{7}{12}$, $\sigma = \frac{11}{13}$, $\omega = \frac{2}{5}$, $\ell = \frac{2}{9}$ and $t_1 = 1$. Then we obtain

$$\mathcal{A}_q(\gamma, \ell) = \mathcal{A}_q(\frac{17}{6}, \frac{2}{9}) = \frac{3}{(1 - \frac{2}{9})\Gamma_q(\frac{17}{6})} = \frac{3}{\frac{7}{9}\Gamma_q(\frac{17}{6})},$$

and

$$\mathcal{B}_q(\gamma, \ell) = \mathcal{B}_q(\frac{17}{6}, \frac{2}{9}) = \frac{2}{(1 - \frac{2}{9})\Gamma_q(\frac{17}{6} - 1)} = \frac{2}{\frac{7}{9}\Gamma_q(\frac{11}{6})},$$

hence

$$\mathcal{C}_q(\gamma, \ell) = \mathcal{C}_q(\frac{17}{6}, \frac{2}{9}) = \max \left\{ \mathcal{A}_q(\frac{17}{6}, \frac{2}{9}), \mathcal{B}_q(\frac{17}{6}, \frac{2}{9}) \right\} = \max \left\{ \frac{3}{\frac{7}{9}\Gamma_q(\frac{17}{6})}, \frac{2}{\frac{7}{9}\Gamma_q(\frac{11}{6})} \right\},$$

which their numerical values are presented in Table 8. Also, the heatmap of Table 8 is presented in Figure 5.

	$q = 0.08$	$q = 0.17$	$q = 0.33$	$q = 0.49$	$q = 0.66$	$q = 0.81$
$\mathcal{A}_q(\frac{17}{6}, \frac{2}{9})$	3.6204	3.3937	3.0657	2.8053	2.5803	2.4142
$\mathcal{B}_q(\frac{17}{6}, \frac{2}{9})$	2.5979	2.6201	2.6508	2.6754	2.6975	2.7145
$\mathcal{C}_q(\frac{17}{6}, \frac{2}{9})$	3.6204	3.3937	3.0657	2.8053	2.6975	2.7145

Table 8: Numerical resual for $\mathcal{A}_q(\frac{17}{6}, \frac{2}{9})$, $\mathcal{B}_q(\frac{17}{6}, \frac{2}{9})$ and $\mathcal{C}_q(\frac{17}{6}, \frac{2}{9})$ in Example 4.3.

Consider the map $\mathbf{X} : \mathbb{R}^4 \rightarrow [0, \infty)$ as follows:

$$\mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4) = \begin{cases} \frac{1}{2} \sum_{i=1}^4 \frac{\|\mathbf{v}_i\|}{1 + \|\mathbf{v}_1\|}, & \mathbf{v}_i, \dots, \mathbf{v}_4 \in [0, 23], \\ |\sin(\mathbf{v}_1 + \dots + \mathbf{v}_4)|, & \mathbf{v}_1, \dots, \mathbf{v}_4 \in (-\infty, 0], \\ -\frac{23}{48} \left(\frac{1}{4} \sum_{i=1}^4 \mathbf{v}_i - 24 \right), & \mathbf{v}_i, \dots, \mathbf{v}_4 \in [23, 24]. \end{cases}$$




Figure 5: The heatmap of Table 8 in Example 4.3.

which implies that $\|\mathbf{X}\|_\infty = 1$. Moreover, define $\psi(\mathbf{v}_1, \dots, \mathbf{v}_4) := \mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4)$, also put $g(\kappa) = z(\kappa) = s(\kappa) = \frac{0.02}{\sqrt{\kappa}}$ and $\Xi(\kappa) = \frac{\kappa}{2}$. If we set $\mathbf{S}(\mathbf{v}_1, \dots, \mathbf{v}_4) = 2$ and

$$\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) := \frac{0.02}{\mathbf{f}(\kappa)} \mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4) + 2,$$

further, we take $\zeta_1 = 0$ and $\zeta_2 = 23$. Then for all $\mathbf{v}_1, \dots, \mathbf{v}_4 \in \mathbf{Y}$ and $\kappa \in \mathbb{Q}^c \cap [0, 1]$ we get

$$\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) < \infty$$

and $\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4)$ is continuous respect to its components. Therefore, we have

$$\begin{aligned} \mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) &= \frac{1}{\mathbf{f}(\kappa)} \mathbf{X}(\mathbf{v}_1, \dots, \mathbf{v}_4) + 2 \\ &\geq z(\kappa) \psi(\mathbf{v}_1, \dots, \mathbf{v}_4) \end{aligned}$$

and

$$\begin{aligned} |\mathbf{h}(\kappa, \mathbf{v}_1, \dots, \mathbf{v}_4) - \mathbf{h}(\kappa, \mathbf{u}_1, \dots, \mathbf{u}_4)| &\leq \sum_{i=1}^4 \frac{1}{2} \times \frac{0.02}{\sqrt{\kappa}} \|\mathbf{v}_i - \mathbf{u}_i\| \\ &= \sum_{i=1}^4 g(\kappa) \Xi(\|\mathbf{v}_i - \mathbf{u}_i\|). \end{aligned}$$

It is worth noting that $\Xi_\beta \in \Psi$ and

$$\sum_{i=1}^4 \|g\|_1 = 4 \times 0.02 = 0.08 < \frac{1}{\mathcal{C}_q(\frac{17}{6}, \frac{2}{9})}.$$

In addition

$$\begin{aligned} \frac{\|\psi\|^* \|z\|_1^L (4 - \gamma^2 - 2\gamma)}{2\Gamma_q(\gamma)(1 - \gamma)} &= \frac{1 \times 0.02 \times (4 - \frac{289}{36} - \frac{17}{3})}{2\Gamma_q(\frac{17}{6})(1 - \frac{17}{6})} \\ &= \frac{0.0969}{1.8333\Gamma_q(\frac{17}{6})} \geq 0 = \zeta_1, \end{aligned}$$

and

$$\begin{aligned} \mathcal{C}_q(\gamma, \ell) \left[\sum_{i=1}^{t_1} \|\mathbf{X}_i\|_\infty \|s_i\|_1 + \|\mathbf{S}\|_1^* \right] &= \mathcal{C}_q\left(\frac{17}{6}, \frac{2}{9}\right) [1 \times 0.02 + 2] \\ &= \mathcal{C}_q\left(\frac{17}{6}, \frac{2}{9}\right) \times 2.02 \leq 23 = \zeta_2. \end{aligned}$$

Now, all the assumptions of Theorem 3.3 are satisfied, so our problem, namely (4.6)-(4.7), has a solution.

5 Conclusion

Today, computer capabilities and software packages are essential for solving complex problems. Therefore, physical phenomenon models must be comprehensible to computers, particularly if they have certain complications like singularity. Our study presents numerical algorithms that investigate the quantitative and qualitative aspects of a pointwise defined multisingular differential equation using quantum fractional operators. We ensured the existence of the solution in three different conditions by using the α - ψ -contraction theorem. Other researchers can also use our method to examine well-known equations with singularities and strong singularities.

Declarations

Availability of Data and Materials

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Funding

Not Applicable.

Authors' contributions

Conceptualization, M.G., S.R.; Formal analysis, R.G., F.S.A., M.G., S.R.; Funding acquisition, R.G.; Methodology, R.G., F.S.A., M.G., S.R.; Software, M.G., S.R.; All authors read and approved the final manuscript. The authors declare that the study was realized in collaboration with equal responsibility.

Competing interests

The authors declare that they have no competing interests.

References

- [1] Baleanu, D.; Jajarmi, A.; Sajjadi, S. S.; Mozyrska, D. A new fractional model and optimal control of a tumor-immune surveillance with non-singular derivative operator, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, **2019**, 29(8), 083127. <https://doi.org/10.1063/1.5096159>

- [2] Hajipour, M.; Jajarmi, A.; Malek, A.; Baleanu, D. Positivity-preserving sixth-order implicit finite difference weighted essentially non-oscillatory scheme for the nonlinear heat equation, *Applied Mathematics and Computation*, **2018**, 325, 146–158. <https://doi.org/10.1016/j.amc.2017.12.026>
- [3] George, R.; Houas, M.; Ghaderi, M.; Rezapour, Sh.; Elagan, S. K. On a coupled system of pantograph problem with three sequential fractional derivatives by using positive contraction-type inequalities, *Results in Physics*, **2022**, 39, 105687. <https://doi.org/10.1016/j.rinp.2022.105687>
- [4] Goswami, A.; Singh, J.; Kumar, D.; Sushila, An efficient analytical approach for fractional equal width equations describing hydro-magnetic waves in cold plasma, *Physica A: Statistical Mechanics and its Applications*, **2019**, 524, 563–575. <https://doi.org/10.1016/j.physa.2019.04.058>
- [5] Singh, J.; Kumar, D.; Kumar, S. New treatment of fractional FornbergWhitham equation via Laplace transform, *Ain Shams Engineering Journal*, **2013**, 4(3), 557–562. <https://doi.org/10.1016/j.asej.2012.11.009>
- [6] B. J. West, P. Grigolini, Applications of Fractional Calculus in Physics, *World Scientific, Singapore*, **1998**.
- [7] A. Carpinteri, F. Mainardi, Fractals and fractional calculus in continuum mechanics, *Springer-Verlag Wien, New York*, **1997**.
- [8] H. Sun, Y. Zhang, D. Baleanu, W. Chen, Y. Chen, A new collection of real world applications of fractional calculus in science and engineering, *Communications in Nonlinear Science and Numerical Simulation*, **2018**, 64, 213–231. <https://doi.org/10.1016/j.cnsns.2018.04.019>
- [9] R. Hilfer, Experimental evidence for fractional time evolution in glass forming materials, *Chemical physics*, **2002**, 284(1-2), 399–408. [https://doi.org/10.1016/S0301-0104\(02\)00670-5](https://doi.org/10.1016/S0301-0104(02)00670-5)
- [10] Sousa, J. V. da. C.; De Oliveira, E. C. On the ψ -Hilfer fractional derivative, *Communications in Nonlinear Science and Numerical Simulation*, **2018**, 60, 72–91. <https://doi.org/10.1016/j.cnsns.2018.01.005>
- [11] Abbas, S.; Benchohra, M.; Lazreg, J. E.; Zhou, Y. A survey on Hadamard and Hilfer fractional differential equations: analysis and stability, *Chaos, Solitons and Fractals*, **2017**, 102, 47–71. <https://doi.org/10.1016/j.chaos.2017.03.010>
- [12] Ali Dokuyucu, M.; Celik, E.; Bulut, H.; Mehmet Baskonus, H. Cancer treatment model with the Caputo-Fabrizio fractional derivative, *The European Physical Journal Plus*, **2018**, 133, 1–6. <https://doi.org/10.1140/epjp/i2018-11950-y>
- [13] Owolabi, K. M.; Atangana, A. Analysis and application of new fractional Adams–Bashforth scheme with Caputo–Fabrizio derivative, *Chaos, Solitons and Fractals*, **2017**, 105, 111–119. <https://doi.org/10.1016/j.chaos.2017.10.020>
- [14] Syam, M. I.; Al-Refai, M. Fractional differential equations with Atangana–Baleanu fractional derivative: analysis and applications, *Chaos, Solitons and Fractals*, **2019**, 2, 100013. <https://doi.org/10.1016/j.csf.2019.100013>
- [15] Gómez-Aguilar, J. F.; Escobar-Jiménez, R. F.; López-López, M. G.; Alvarado-Martínez, V. M. Atangana–Baleanu fractional derivative applied to electromagnetic waves in dielectric media, *Journal of Electromagnetic Waves and Applications*, **2016**, 30(15), 1937–1952. <https://doi.org/10.1080/09205071.2016.1225521>
- [16] Z. Heydarpour, M. N. Parizi, R. Ghorbanian, M. Ghaderi, Sh. Rezapour, A. mosavi, A study on a special case of the Sturm-Liouville equation using the Mittag-Leffler function and a new type of contraction, *AIMS Mathematics*, **2022**, 7(10), 10665–10684. <https://doi.org/10.3934/math.20221004>
- [17] Awadalla, M.; Noupoue, Y. Y. Y.; Asbeh, K. A.; Ghiloufi, N.; Rezapour, A. mosavi, Modeling drug concentration level in blood using fractional differential equation based on Psi-Caputo derivative, *Journal of Mathematics*, **2022**, 9006361. <https://doi.org/10.1155/2022/9006361>
- [18] Samet, B.; Aydi, H. Lyapunov-type inequalities for an anti-periodic fractional boundary value problem involving ψ -Caputo fractional derivative, *Journal of inequalities and applications*, **2018**, 1–11. <https://doi.org/10.1186/s13660-018-1850-4>

- [19] George, R.; Al-shammari, F.; Ghaderi, M.; Rezapour, Sh. On the boundedness of the solution set for the ψ -Caputo fractional pantograph equation with a measure of non-compactness via simulation analysis, *AIMS Mathematics*, **2023**, 8(9), 20125–20142. <https://doi.org/10.3934/math.20231025>
- [20] Agarwal, R. P.; O'Regan, D.; Staněk, S. Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations, *Journal of Mathematical Analysis and Applications*, **2010**, 371(1), 57–68. <https://doi.org/>
- [21] Wenquan, F.; Shurong, S.; Zhenlai, H.; Yige, Z. Existence of solutions for a singular system of nonlinear fractional differential equations, *Computers and Mathematics with Applications*, **2011**, 62(3), 1370–1378. <https://doi.org/10.1016/j.camwa.2011.03.076>
- [22] SeakWeng, V. Positive solutions of singular fractional differential equations with integral boundary conditions, *Mathematical and Computer Modelling*, **2013**, 57(5-6), 1053–1059. <https://doi.org/10.1016/j.mcm.2012.06.024>
- [23] Nyamoradi, N.; Bashiri, T.; Vaezpour, S. M.; Baleanu, D. Uniqueness and existence of positive solutions for singular fractional differential equations, *Electronic Journal of Differential Equations*, **2014**, 130, 1–13.
- [24] Malekpour, A.; Shabibi, M.; Nuraei, R. Existence of a solution for a multi singular point-wise defined fractional differential system, *Journal Of Mathematical Extension*, **2022**, 16(4), 1–36. <https://doi.org/0.30495/JME.2022.1118>
- [25] Guo, L.; Liu, L.; Wu, Y. Existence of positive solutions for singular fractional differential equations with infinite-point boundary conditions, *Nonlinear Analysis: Modelling and Control*, **2016**, 21(5), 635–650. <https://doi.org/10.15388/NA.2016.5.5>
- [26] Xingqiu, Z. Positive solutions for a class of singular fractional differential equation with infinite-point boundary value conditions, *Applied Mathematics Letters*, **2015**, 39, 22–27. <https://doi.org/10.1016/j.aml.2014.08.008>
- [27] Khan, A.; Khan, H.; Gómez-Aguilar, J. F.; Abdeljawad, T. Existence and Hyers-Ulam stability for a non-linear singular fractional differential equations with Mittag-Leffler kernel, *Chaos, Solitons and Fractals*, **2019**, 127, 422–427. <https://doi.org/10.1016/j.chaos.2019.07.026>
- [28] Jackson, F. H. q -Difference equation, *American Journal of Mathematics*, **1910** 32(4), 305–314.
- [29] Jackson, F. H. On Q -Definite Integrals, *The Quarterly Journal of Pure and Applied Mathematics*, **1910** 41, 193–203.
- [30] V. Kac, P. Cheung, Quantum Calculus, *Springer-Verlag, New york*, **2002**.
- [31] Atici, F. M.; Eloe, P. w. Fractional q -calculus on a time scale, *Journal of Nonlinear Mathematical Physics*, **2007**, 14(3), 341–352. <https://doi.org/10.2991/jnmp.2007.14.3.4>
- [32] Rajković, P. M.; Marinković, S. D.; Stanković, M. S. Fractional integrals and derivatives in q -calculus, *Applicable analysis and discrete mathematics*, **2007**, 14(3), 311–323.
- [33] Ferreira, R. A. c. Positive solutions for a class of boundary value problems with fractional q -differences, *Computers and Mathematics with Applications*, **2011**, 61(2), 367–373. <https://doi.org/10.1016/j.camwa.2010.11.012>
- [34] Li, X.; Han, Z.; Li, X. Boundary value problems of fractional q -difference Schringer equations, *Applied Mathematics Letters*, **2015**, 46, 100–105. <https://doi.org/10.1016/j.aml.2015.02.013>
- [35] Guo, F.; Kang, Sh.; Chen, F. Existence and uniqueness results to positive solutions of integral boundary value problem for fractional q -derivatives, *Advances in Difference Equations*, **2018**, 379, 1–15. <https://doi.org/10.1186/s13662-018-1796-3>
- [36] Yo, Ch.; Wang, J. Existence of solutions for nonlinear second-order q -difference equations with first-order q -derivatives, *Advances in Difference Equations*, **2013**, 124, 1–11. <https://doi.org/10.1186/1687-1847-2013-124>

- [37] El-Emam, F. Z. Convolution conditions for two subclasses of analytic functions defined by Jackson q -difference operator, *Journal of the Egyptian Mathematical Society*, **2022**, 7, 1–10. <https://doi.org/10.1186/s42787-022-00141-2>
- [38] Lio, Z. G. On the q -derivative and q -series expansions, *International Journal of Number Theory*, **2013**, 9(8), 2069–2089. <https://doi.org/10.1142/S1793042113500759>
- [39] Mahmood, Sh.; Jabeen, M.; Malik, S. N.; Srivastava, H. M.; Manzoor, R.; Riaz, S. M. J. Some Coefficient Inequalities of q -Starlike Functions Associated with Conic Domain Defined by q -Derivative, *Journal of Function Spaces*, **2018**, 8492072. <https://doi.org/10.1155/2018/8492072>
- [40] Zhai, C.; Ren, J. The unique solution for a fractional q -difference equation with three-point boundary conditions, *Indagationes Mathematicae*, **2018**, 29(3), 948–961. <https://doi.org/10.1016/j.indag.2018.02.002>
- [41] Ali, M. U.; Kamran, T.; Karapinar, E. A new approach to α - ψ -contractive non-self multivalued mappings, *Journal of Inequalities and Applications*, **2014**, 71, 1–9. <https://doi.org/10.1186/1029-242X-2014-71>
- [42] J. R. Graef, L. Kong, Positive solutions for a class of higher order boundary value problems with fractional q -derivatives, *Applied Mathematics and Computation*, **2012**, 218(19), 9682–9689. <https://doi.org/10.1016/j.amc.2012.03.006>
- [43] M. El-Shahed, F. M. Al-Askar, Positive Solutions for Boundary Value Problem of Nonlinear Fractional q -Difference Equation, *International Scholarly Research Notices*, **2011**, 218(19), 385459. <https://doi.org/10.5402/2011/385459>
- [44] Samet, B.; Vetro, C.; Vetro, P. Fixed point theorem for α - ψ contractive type mappings, *Nonlinear analysis: theory, methods and applications*, **2012**, 75(4), 2154–2165. <https://doi.org/10.1016/j.na.2011.10.014>