

Oxidant and Antioxidant Balance in Patients with COVID-19

Kubra Aykac¹, Yasemin Ozsurekci², Burcu Ceylan Cura Yayla¹, Sibel Lacinel Gurlevik², Pembe Derin Oygur², Nuriye Bodur Uluc³, Medine Aysin Tasar³, Fatma Sebnem Erdinc⁴, Gunay Ertem⁴, Salim Neselioglu⁵, Ozcan Erel⁵, Ali Bülent Cengiz², Mehmet Ceyhan²

¹Department of Pediatric Infectious Diseases, University of Health Science, Ankara Training and Research Hospital, Ankara, Turkey

²Department of Pediatric Infectious Diseases, Hacettepe University Faculty of Medicine, Ankara, Turkey

³Department of Pediatric Diseases, University of Health Science, Ankara Training and Research Hospital, Ankara, Turkey

⁴Department of Infectious Diseases, University of Health Science, Ankara Training and Research Hospital, Ankara, Turkey

⁵Department of Clinical Biochemistry, Yildirim Beyazıt University Faculty of Medicine, Ankara, Turkey

Kubra Aykac, Department of Pediatric Infectious Diseases, University of Health Science, Ankara Training and Research Hospital, Ankara, Turkey

kubraklnc.kk@gmail.com

+90 533 922 00 87

ORCID: [0000-0002-0974-4765](https://orcid.org/0000-0002-0974-4765)

Running head: Oxidative Stress in Patients with COVID-19

Abstract

Background: A crucial balance exists between oxidant and antioxidant mechanisms in the functional immune system. We aimed to evaluate the contributions of balance between these systems to coronavirus disease 2019 (COVID-19), a devastating pandemic caused by viral infection.

Method: We analyzed serum oxidant and antioxidant stress parameters according to the clinical and demographic characteristics of children and adults with COVID-19 and compared them against the values of healthy controls. Serum native thiol (NT), total thiol (TT), disulfide, total antioxidant status, total oxidant status, and ischemia-modified albumin levels were evaluated and compared between groups.

Results: A total of 79 children and 74 adults were evaluated in the present study, including 46 children and 40 adults with COVID-19, 33 healthy children, and 34 healthy adults. TT, NT, and disulfide levels were significantly lower in the adult COVID-19 group than in all other groups ($p = 0.001$, $p = 0.001$, and $p = 0.005$, respectively). Additionally, TT and NT levels were significantly lower in both pediatric and adult COVID-19 cases with severe disease course than mild/moderate course. TT and NT levels were identified as predictors for the diagnosis of the adult COVID-19 cases and as independent predictors for disease severity in both children and adults with COVID-19.

Conclusion: Parameters that reveal the oxidant and antioxidant capacity, including TT and NT, appear to be good candidates for the accurate prediction of the clinical course among patients with COVID-19.

Key words: oxidative stress; thiols; COVID-19; pediatrics; severity

1. Introduction

The current coronavirus disease 2019 (COVID-19) pandemic is the most devastating global disease experienced in a century, and, unfortunately, its clinical manifestations, pathogenesis, and treatment remain unclear^{1,2}. The clinical course of this disease has ranged from asymptomatic to severe pneumonia and death. Additionally, the course of the disease differs in children and adults, and we are currently unable to predict the future prognosis of patients at the time of diagnosis³. Therefore, an in-depth understanding of the disease pathogenesis of the disease is necessary to predict the occurrence of severe pneumonia, identify potentially critical patients, develop appropriate treatments, and reduce hospitalization rates and mortality.

Oxidative stress may be a key player in COVID-19 pathogenesis because it plays a significant role in the response to infections.⁴. Studies have demonstrated that oxidative stress regulates the host immune system in viral diseases, such as hepatitis B, hepatitis C, herpes simplex virus, and influenza^{5, 6, 7, 8}. Additionally, oxidative stress can contribute to the pathogenesis of several lung diseases in children, such as pneumonia, asthma, acute bronchiolitis, cystic fibrosis, acute respiratory distress syndrome, and chronic neonatal lung disease^{9, 10, 11}. Reviews have reported a potential link between oxidative stress and the pathogenesis, severity, and mortality risk of patients infected with severe acute respiratory syndrome coronavirus (SARS-CoV)^{4 12, 13}. However, to the best of our knowledge, no clinical study has been performed in humans with COVID-19 to investigate the oxidant and antioxidant stress parameters.

Because we suspected a potential relationship between oxidative stress and COVID-19, in this study, we evaluated the oxidant and oxidative stress statuses in both children and adults with COVID-19 compared with health controls and examined the use of oxidative stress parameters as a predictor of disease severity.

2. Materials and Methods

We conducted a prospective cohort study including patients diagnosed with COVID-19 between April 10, 2020, and August 10, 2020, who were admitted to University of Health Sciences, Ankara Educating and Training Hospital, and Hacettepe University Ihsan Dogramaci Children's Hospital. The study was approved by the Public Health Agency, Turkish Ministry of Health and Ethics Board of University of Health Sciences, Ankara Educating and Training Hospital, Turkey (E-322). All parents or guardians provided written informed consent.

We diagnosed cases via reverse transcriptase-polymerase chain reaction (RT-PCR), according to the national COVID-19 guidelines established by the Coronavirus Scientific Advisory Board in Turkey in response to new data regarding the disease¹⁴. The demographic and clinical characteristics of patients were obtained from medical records maintained by the two hospitals. We categorized the severity of pediatric COVID-19 cases based on the clinical characteristics and the results of laboratory examinations and radiologic imaging, as defined by Dong et al.³.

The severity of adult cases was defined according to the criteria found in the World Health Organization (WHO) interim guidelines¹⁵. At the time of admission, serum samples were obtained from pediatric and adult cases diagnosed with COVID-19, and all samples were stored at -80°C until oxidative stress parameters were measured. Age and gender-matched pediatric and adult healthy volunteers were enrolled as controls. Serum native thiol (NT), total thiol (TT), disulfide (DD), total antioxidant status (TAS), total oxidant status (TOS), and ischemia-modified albumin (IMA) levels were evaluated in both patients and controls and compared between groups.

The analysis of thiol/DD levels was performed as described by Erel and Neselioglu, using a method based on the reduction of DD bonds into reactive thiol groups in the presence of

NaBH₄¹⁶. Serum TAS and TOS levels were measured using automated colorimetric measurement methods, as described by Erel [17, 18]. Serum IMA levels were measured using a colorimetric assay previously described by Bar-Or et al.¹⁹.

Statistical analyses

All statistical analyses were performed using SPSS, version 19.0 (SPSS, Inc., Chicago, IL, USA), and a trial version of Medcalc statistical software. Descriptive statistics were used to summarize the participants' baseline characteristics, as the median and interquartile range (IQR) for continuous variables and as frequency distributions for categorical variables. The student's t-test was used to compare continuous variables with a normal distribution, whereas the Mann–Whitney U test was applied to comparisons of non-normally distributed variables. Cut-off values used to distinguish between the case and control groups were determined using receiver operator characteristic (ROC) analysis, after which the sensitivity and specificity were calculated. Two-tailed tests were used for all analyses, and $p < 0.05$ was considered significant.

3. Results

Serum samples were collected from a total of 46 children (median age: 10 years; IQR: 3.7–14 years; 41.3% male) and 40 adults (median age: 50 years; IQR: 43–64 years; 52.5% male) who were diagnosed with COVID-19. The age- and gender-matched control group consisted of 33 healthy children and 34 healthy adults. Based on disease status, the patients were divided into 4 groups: Group 1, children with COVID-19; Group 2, adults with COVID-19; Group 3, healthy children; and Group 4, healthy adults. In Group 1, 10 (21.7%) children had severe/critical disease and 36 (78.3%) children had mild/moderate disease. In Group 2, 18 (45%) adults had severe/critical disease, and 22 (55%) adults had mild/moderate disease. Two

pediatric cases and no adult cases resulted in fatalities. The pediatric patients who died were aged 2 and 7 years, and neither had any preexisting comorbidities.

When the parameters of pediatric and adult patients with COVID-19 were compared, significantly reduced serum levels of NT, TT, and DD were detected in the adult group ($p = 0.001$, $p = 0.001$, and $p = 0.005$, respectively). No significant differences were in NT, TT, and DD levels were observed for any other 2-group comparisons (Group 1 vs. Group 3 or Group 2 vs. Group 4). No significant differences in TAS, TOS, or IMA values were observed between adult and pediatric COVID-19 patients (Group 1 vs. Group 2). The demographic, clinical, and laboratory parameters of patients with COVID-19 are shown in Table 1.

When the parameters of pediatric patients with COVID-19 and healthy children were evaluated according to disease course, significantly lower serum TT and NT levels were observed in children with a severe/critical disease presentation compared with those with a mild or moderate disease presentation ($p = 0.006$, $p = 0.003$, and $p = 0.006$, $p = 0.002$, respectively) (Table 2). Consistently, significantly lower serum TT and NT levels were also observed in adults with a severe/critical disease course compared with patients who had a mild or moderate disease course ($p = 0.004$, $p = 0.01$, and $p = 0.003$, $p = 0.01$, respectively, Table 3).

The calculated cut-off values for TT and NT used to distinguish pediatric patients with COVID-19 were 440 and 480 $\mu\text{mol/L}$, respectively, which resulted in sensitivities of 50% and 52% and specificities of 72% and 72%. The area under the ROC curve (AUC) values (with 95% confidence intervals [95% CI]) were 0.614 (0.486–0.742) for NT and 0.618 (0.490–0.745) for TT ($p = 0.08$ and $p = 0.07$, respectively, Fig. 1A). The calculated cut-off values for TT and NT used to distinguish adult COVID-19 patients were 379 and 420 $\mu\text{mol/L}$, respectively, which resulted in sensitivities of 58% and 58% and specificities of 82% and

82%. The AUC (95% CI) values were 0.701 (0.58–0.822) for NT and 0.699 (0.578–0.82) for TT ($p = 0.003$ and $p = 0.003$, respectively, Fig. 1B).

We further analyzed whether these oxidant/antioxidant stress parameters could be used to predict COVID-19 disease severity. Adult and pediatric patients were separately divided into a severe group, which contained all severe and critically ill cases, and a non-severe group, which included both mild and moderate cases. The ROC curves for TT and NT were calculated separately for children and adults. The results for children showed that the AUC values were 0.889 for NT and 0.885 for TT ($p = 0.001$ and $p = 0.001$, respectively, Fig. 1C). The results for adults showed that the AUC values were 0.707 for NT and 0.705 for TT ($p = 0.002$ and $p = 0.002$, respectively, Fig. 1D).

4. Discussion

We found that TT and NT levels were significantly lower in both pediatric and adult COVID-19 cases with severe disease course in the present study. Our findings supported the potential utility of using thiol levels as a predictor of COVID-19 severity, and the higher thiol levels observed in mild or moderate cases may serve as an indirect indicator of the improved antioxidant status in lung tissues due to better control of the proinflammatory processes in the lungs²⁰. Additionally, our data showed that the TT and NT levels were lower in adults than in children with COVID-19, and these levels appeared to be good predictors of COVID-19 diagnosis, particularly among adults. The better thiol/DD balance observed in children might indicate they might be better protected from the harmful effects of oxidative stress than adults. Although severe patients presented with low thiol levels in both groups, the milder clinical course that is generally observed in children might be explained by these differences in the oxidative statuses of pediatric and adult cases.

The imbalance between oxidant production and antioxidant mechanisms is referred to as oxidative stress and can result in oxidative damage caused by the peroxidation of lipids and the oxidation of DNA (11). Reactive oxygen species (ROS) are produced by activated inflammatory cells in response to infectious diseases²¹. Viral infections have been found to be associated with a decreased antioxidant defense, in addition to increased neutrophil infiltration and ROS release²². The high neutrophil to lymphocyte ratio observed in severe COVID-19 cases²³, the neutrophilic infiltration of pulmonary capillaries, and the increase in neutrophil activation, including the circulation of neutrophil extracellular traps (NETs), have all been suggested to serve as indicators of COVID-19^{24,25}. Thiols are major total body antioxidants that play crucial roles against ROS, protecting against damage induced by free radicals, and thiol levels decrease during the neutralization of ROS in cases of high levels of oxidative stress^{26,27}. Consistent with our findings, relative to those in healthy individuals, the TT and NT levels have been reported to be reduced during many infectious/inflammatory processes, such as acute tonsillopharyngitis, meningism, meningitis, acute appendicitis, Crimean–Congo hemorrhagic fever (CCHF), and brucellosis^{18,29,30,31,32}. Moreover, in an unpublished study performed at our center, thiol levels were found to be useful as a severity predictor for community acquired pneumonia (CAP), and the high thiol levels observed in outpatients with CAP were suggested as a potential indirect indicator of improved antioxidant status in the lung tissue, indicating the status of the proinflammatory processes in the lungs. We postulate that excessive ROS production might occur in both severe pediatric and adult COVID-19 cases; however, this requires further detailed confirmation. An in-depth understanding of the critical disease course would allow for the identification of promising avenues for the investigation of the potential pathophysiological mechanisms underlying COVID-19 and for the development of reasonable COVID-19 management strategies.

Thiol groups are among the most abundant and important antioxidant molecules found in both cells and plasma and appear to serve as the primary regulators of oxidative stress. Thiols are organic compounds contained within protein structures. Thiol groups can form reversible DD bridges through the effect of oxidants to maintain dynamic thiol-DD homeostasis¹⁶. The thiol-DD balance can be affected by oxidative stress and is crucial for viral entry, viral reactivity, and fusion into the host cell^{33,34,35,36}. Hati et al. investigated the molecular basis of SARS-CoV-2 cellular recognition and reported that the binding affinity was significantly impaired when the DD bonds in both angiotensin-converting enzyme II (ACE2) and the SARS-CoV/CoV-2 spike proteins were reduced to thiol groups³⁶, which indicated that the reduction or absence of oxidative stress could have significant beneficial effects during the early stage of viral infection by preventing viral protein binding with host cells.

To gain entry into human cells, a spike glycoprotein on the viral envelope of SARS-CoV-2 interacts with the ACE2 receptor, which is expressed in various human organs. Therefore, the spike glycoprotein appears to represent a promising potential target for the development of specific drug treatments and vaccines³⁷. Due to the urgent necessity of identifying effective anti-SARS-CoV-2 therapeutic agents, the typical drug development process may require too much time; therefore, a focus on the potential use of currently FDA-approved drugs might represent a reasonable therapeutic approach. Disulfiram (DSF), which is a thiol-reacting FDA-approved drug, appears to be an acceptable candidate drug that is often used over the long term by patients with chronic alcoholism^{38,39}. DSF was found to present antimycobacterial properties against *Mycobacterium tuberculosis*⁴⁰. DSF, along with other thiol-reactive compounds (l-captopril and bithionol) and quercetin, are defined as urease inhibitors and have been recognized as potential antimicrobial strategies for the treatment of *Citrullus vulgaris*⁴¹. The human opportunistic pathogen *Pseudomonas aeruginosa* is also targeted by DSF metabolites⁴². Carbamate kinase is an essential enzyme required for the

survival of *Giardia lamblia*, and DSF inhibits this kinase, killing the trophozoites *in vitro*⁴³. Recently, DSF was reported to possess antiviral activity against the hepatitis C virus⁴⁴ and was determined to serve as an inhibitor of the papain-like proteases associated with SARS-CoV-1 and Middle East respiratory syndrome coronavirus (MERS-CoV)⁴⁵. Finally, both DSF and its metabolites have clearly been demonstrated to penetrate into human cells to block the proteolytic functions of 3CL^{pro}, which is a protein that is crucial for SARS-CoV-2 replication. The results of the present study indicated that thiol-mediated inflammation should be considered in severe/critically ill COVID-19 patients and that thiol-reacting agents, such as DSF, might represent a reasonable therapeutic option. However, this assumption requires additional clinical confirmation.

Several limitations of this study should be noted. First, the sample sizes were relatively small; therefore, our result cannot be generalized and require confirmation in a larger cohort to reveal the actual disease nature. Second, the detailed analyses of many of the oxidant and antioxidant parameters mentioned in the literature could not be performed in this study or linked with the disease course. For instance, the decreased expression of the antioxidant enzyme superoxide dismutase 3 (SOD3) in the lungs of a COVID-19 patient was associated with disease severity⁴⁶. However, we believe that the findings of our study will be able to guide physicians and improve our understanding of the potential physio-pathological pathways involved in COVID-19.

In conclusion, serum oxidant and antioxidant stress parameters could change in response to disease severity in COVID-19 patients. We found that serum TT and NT levels might serve as significant predictors of disease severity in both pediatric and adult COVID-19 patients. Evaluating the thiol-DD balance, which serves as a marker of oxidative stress, appears to be essential for understanding the pathogenesis of various viral infections, including COVID-19. In addition, Parlak et al. also concluded that the thiol/DD status might represent an important

variable for monitoring the treatment response in pneumonia⁴⁷, which is a serious complication of COVID-19. Our findings shed light on the disease pathogenesis, emphasizing the significance of oxidative stress and the potential use of both new and currently available therapeutic options, such as thiol-reacting agents and metabolites for the treatment of COVID-19.

5. Credit author statement

KA: Data curation, Methodology, Writing- Original draft preparation. YO: Data curation, Writing- Writing- Reviewing and Editing. BCCY: Conceptualization, Writing- Original draft preparation. SLG: Data curation. PDO: Data curation. NBU: Data curation. MAT: Supervision. FSE: Supervision. SE: Supervision. SN: Data curation. OE: Supervision. ABC: Supervision. MC: Supervision

Funding

None to report.

Declaration of Competing Interest

The authors report no declarations of interest.

References

- 1-World Health Organization (WHO): Coronavirus disease (COVID-19) outbreak situation. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> Accessed 29 Sep 2020
- 2- Shen KL, Yang YH, Jiang RM, et al. Updated diagnosis, treatment and prevention of COVID-19 in children: experts' consensus statement (condensed version of the second edition). *World J Pediatr.* 2020;16(3):232-239. doi:10.1007/s12519-020-00362-4
- 3-Dong Y, Mo X, Hu Y et al. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. *Pediatrics.* 2020 Mar 16
- 4- Delgado-Roche L, Mesta F. Oxidative Stress as Key Player in Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV) Infection. *Arch Med Res.* 2020;51(5):384-387. doi:10.1016/j.arcmed.2020.04.019
- 5-Hunt N.H., van Reyk D.M., Fragonas J.C., Jeitner T.M., Goldstone S.D. (1994) Redox Mechanisms in T Cell Activation. In: Pasquier C., Olivier R.Y., Auclair C., Packer L.

(eds) Oxidative Stress, Cell Activation and Viral Infection. Molecular and Cell Biology Updates. Birkhäuser Basel. https://doi.org/10.1007/978-3-0348-7424-3_21

6- Ha HL, Shin HJ, Feitelson MA, Yu DY. Oxidative stress and antioxidants in hepatic pathogenesis. *World J Gastroenterol*. 2010;16(48):6035-6043. doi:10.3748/wjg.v16.i48.6035

7-Gonzalez-Dosal R, Horan KA, Rahbek SH, Ichijo H, Chen ZJ, Mieyal JJ ,et al. HSV infection induces production of ROS, which potentiate signaling from pattern recognition receptors: role for S-glutathionylation of TRAF3 and 6. *Plos Pathog* 2011;7:e1002250.

8- Gonzalez-Dosal R, Horan KA, Rahbek SH, et al. HSV infection induces production of ROS, which potentiate signaling from pattern recognition receptors: role for S-glutathionylation of TRAF3 and 6. *PLoS Pathog*. 2011;7(9):e1002250. doi:10.1371/journal.ppat.1002250

9-Aykac K, Ozsurekci Y, Tanir Basaranoglu S, et al. Oxidant and antioxidant balance in children with bacteremia [published online ahead of print, 2020 Sep 3]. *Minerva Pediatr*. 2020;10.23736/S0026-4946.20.05748-5. doi:10.23736/S0026-4946.20.05748-5

10-J. Ciencewicki, S. Trivedi, S.R. Kleeberger. Oxidants and the pathogenesis of lung diseases. *J Allergy Clin Immunol*. 2008;122:456-470.

11-H.S. Park, S.R. Kim, Y.C. Lee. Impact of oxidative stress on lung diseases. *Respirology*. 2009;14:27-38.

12-Derouiche S. Oxidative Stress Associated with SARS-Cov-2 (COVID-19) Increases the Severity of the Lung Disease - A Systematic Review. *J Infect Dis Epidemiol* 2020;6:121.

13-Zeliger HI, Kahaner H. Can the Oxidative Stress Index Predict the Severity of COVID-19? *EJMED, European Journal of Medical and Health Sciences* Vol. 2, No. 2, April 2020

14-The Coronavirus Scientific Advisory Board (Turkey).
https://covid19bilgi.saglik.gov.tr/depo/rehberler/COVID-19_Rehberi.pdf Last update: 14 April 2020.

15-World Health Organization. (2020). Clinical management of COVID-19: interim guidance, 27 May 2020. World Health Organization. <https://apps.who.int/iris/handle/10665/332196>. License: CC BY-NC-SA 3.0 IGO

16- Erel O, Neselioglu S. A novel and automated assay for thiol/disulphide homeostasis. *Clin Biochem*. 2014;47(18):326-332. doi:10.1016/j.clinbiochem.2014.09.026

17-Erel, O. A new automated colorimetric method for measuring total oxidant status. *Clin. Biochem*. 2005;38:1103–1111.

18-Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin Biochem* 2004;37:277-85.

19-Bar-Or D, Lau E, Winkler JV. A novel assay for cobalt-albumin binding and its potential as a marker for myocardial ischemia-a preliminary report. *J Emerg Med*. 2000;19:311-5.

20-Leelarungrayub J, Borisuthibandit T, Yankai A, Boontha K. Changes in oxidative stress from tracheal aspirates sampled during chest physical therapy in hospitalized intubated infant patients with pneumonia and secretion retention. *Ther Clin Risk Manag*. 2016;12:1377-1386. Published 2016 Sep 6. doi:10.2147/TCRM.S112972

21-Gantt KR, Goldman TL, McCormick ML, et al. Oxidative responses of human and murine macrophages during phagocytosis of *Leishmania chagasi*. *J Immunol*. 2001;167(2):893-901. doi:10.4049/jimmunol.167.2.893

22-Laforge M, Elbim C, Frère C, et al. Tissue damage from neutrophil-induced oxidative stress in COVID-19 [published correction appears in *Nat Rev Immunol*. 2020 Aug 10;]. *Nat Rev Immunol*. 2020;20(9):515-516. doi:10.1038/s41577-020-0407-1

- 23-Fu J, et al. The clinical implication of dynamic neutrophil to lymphocyte ratio and D-dimer in COVID-19: a retrospective study in Suzhou China. *Thromb. Res.* 2020;192:3–8. doi: 10.1016/j.thromres.2020.05.006.
- 24-Barnes BJ, et al. Targeting potential drivers of COVID-19: neutrophil extracellular traps. *J. Exp. Med.* 2020;217:e20200652. doi: 10.1084/jem.20200652.
- 25-Golonka RM, Saha P, Yeoh BS, et al. Harnessing innate immunity to eliminate SARS-CoV-2 and ameliorate COVID-19 disease. *Physiol Genomics.* 2020;52(5):217-221. doi:10.1152/physiolgenomics.00033.2020
- 26-Biswas S, Chida AS, Rahman I. Redox modifications of protein-thiols: emerging roles in cell signaling. *Biochem Pharmacol.* 2006;71:551-64.
- 27-Gulpamuk B, Tekin K, Sonmez K et al. The significance of thiol/disulfide homeostasis and ischemia-modified albumin levels to assess the oxidative stress in patients with different stages of diabetes mellitus. *Scand J Clin Lab Invest.* 2018;78:136-142.
- 28-Esen R, Aslan M, Kucukoglu ME et al. Serum paraoxonase activity, total thiols levels, and oxidative status in patients with acute brucellosis. *Wien Klin Wochenschr.* 2015;127:427-33.
- 29-Kara SS, Erel O, Demirdag TB et al. Alteration of thiol-disulphide homeostasis in acute tonsillopharyngitis. *Redox Rep.* 2017;22:205-209.
- 30-Aycicek A, Iscan A, Erel O, Akcali M, Selek S. Total Antioxidant/Oxidant Status in Meningism and Meningitis. *Pediatr Neurol.* 2006;35:382-6.
- 31-Ozyazici S, Karateke F, Turan U et al. A Novel Oxidative Stress Mediator in Acute Appendicitis: Thiol/Disulphide Homeostasis Mediators Inflamm. 2016;2016:6761050.
- 32-Karadag-Oncel E, Erel O, Ozsurekci Y et al. Plasma oxidative stress and total thiol levels in Crimean-Congo hemorrhagic fever. *Jpn J Infect Dis.* 2014;67:22-26.
- 33-Suhail S, Zajac J, Fossum C et al. Role of Oxidative Stress on SARS-CoV (SARS) and SARS-CoV-2 (COVID-19) Infection: A Review. *The Protein Journal.* 2020;

- 34-Markovic I, Stantchev TS, Fields KH, Tiffany LJ, Tomić M, Weiss CD, Broder CC, Strebel K, Clouse KA (2004) Thiol/disulfide exchange is a prerequisite for CXCR4-tropic HIV-1 envelope-mediated T-cell fusion during viral entry. *Blood* 2004;103:1586–1594
- 35-Lavillette D, Barbouche R, Yao Y, Boson B, Cosset F, Jones IM, Fenouillet E (2006) Significant redox insensitivity of the functions of the SARS-CoV spike glycoprotein: comparison with HIV envelope. *J Biol Chem* 2006;281:9200–9204
- 36-Hati S, Bhattacharyya S. Impact of Thiol-Disulfide Balance on the Binding of Covid-19 Spike Protein with Angiotensin-Converting Enzyme 2 Receptor. *ACS Omega*. 2020;5(26):16292-16298. Published 2020 Jun 23. doi:10.1021/acsomega.0c02125
- 37-Ni W, Yang X, Yang D, et al. Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19. *Crit Care*. 2020;24(1):422. Published 2020 Jul 13. doi:10.1186/s13054-020-03120-0
- 38-Lobo-Galo, N. , Terrazas-López, M. , & Martínez-Martínez, A. (2020). FDA-approved thiol-reacting drugs that potentially bind into the SARS-CoV-2 main protease, essential for viral replication. *Journal of Biomolecular Structure and Dynamics* , 1–12. 2020;
- 39-Yoshimura, A., Kimura, M., Nakayama, H., Matsui, T., Okudaira, F., Akazawa, S., Ohkawara, M., Cho, T., Kono, Y., Hashimoto, K., Kumagai, M., Sahashi, Y., Roh, S., & Higuchi, S. (2014). Efficacy of disulfiram for the treatment of alcohol dependence assessed with a multicenter randomized controlled trial. *Alcoholism, Clinical and Experimental Research*, 38(2), 572–578. doi: 10.1111/acer.12278
- 40-Dalecki, A. G., Haeili, M., Shah, S., Speer, A., Niederweis, M., Kutsch, O., & Wolschendorf, F. (2015). Disulfiram and copper ions kill *Mycobacterium tuberculosis* in a synergistic manner. *Antimicrobial Agents and Chemotherapy*, 59(8), 4835–4844. doi: 10.1128/AAC.00692-15

- 41- Díaz-Sánchez ÁG, Alvarez-Parrilla E, Martínez-Martínez A, et al. Inhibition of Urease by Disulfiram, an FDA-Approved Thiol Reagent Used in Humans. *Molecules*. 2016;21(12):1628. Published 2016 Nov 26. doi:10.3390/molecules21121628
- 42- Zaldívar-Machorro VJ, López-Ortiz M, Demare P, Regla I, Muñoz-Clares RA. The disulfiram metabolites S-methyl-N,N-diethyldithiocarbamoyl sulfoxide and S-methyl-N,N-diethylthiocarbamoyl sulfone irreversibly inactivate betaine aldehyde dehydrogenase from *Pseudomonas aeruginosa*, both in vitro and in situ, and arrest bacterial growth. *Biochimie*. 2011;93(2):286-295. doi:10.1016/j.biochi.2010.09.022
- 43- Galkin A, Kulakova L, Lim K, et al. Structural basis for inactivation of Giardia lamblia carbamate kinase by disulfiram. *J Biol Chem*. 2014;289(15):10502-10509. doi:10.1074/jbc.M114.553123
- 44- Lee YM, Duh Y, Wang ST, Lai MM, Yuan HS, Lim C. Using an Old Drug to Target a New Drug Site: Application of Disulfiram to Target the Zn-Site in HCV NS5A Protein. *J Am Chem Soc*. 2016;138(11):3856-3862. doi:10.1021/jacs.6b00299
- 45-Lin MH, Moses DC, Hsieh CH, et al. Disulfiram can inhibit MERS and SARS coronavirus papain-like proteases via different modes. *Antiviral Res*. 2018;150:155-163. doi:10.1016/j.antiviral.2017.12.015
- 46- Abouhashem AS, Singh K, Azzazy HME, Sen CK. Is Low Alveolar Type II Cell *SOD3* in the Lungs of Elderly Linked to the Observed Severity of COVID-19?. *Antioxid Redox Signal*. 2020;33(2):59-65. doi:10.1089/ars.2020.8111
- 47-Parlak ES, Alisik M, Hezer H, Karalezli A, Hasanoglu HC, Erel O. Evaluation of dynamic thiol/disulfide redox state in community-acquired pneumonia. *Saudi Med J*. 2018;39(5):495-499. doi:10.15537/smj.2018.5.22111

Figure Legends

Figure 1. The ROC curve of serum native/total thiol levels on admission for both pediatric and adult patients with COVID-19 and pediatric and adult cases with severe disease course or not. **A.** The AUC of the ROC curve for serum native/total thiol levels was calculated in children **B.** The AUC of the ROC curve for serum native/total thiol levels was calculated in adults. **C.** The AUC of the ROC curve for serum native/total thiol levels was calculated in children with severe disease **D.** The AUC of the ROC curve for serum native/total thiol levels was calculated in adults with severe disease