

Table 2. Characteristics of 2 groups after ablation

	Group 1 n=9	Group 2 n=22	p value
echocardiographic parameters			
follow-up period of echocardiography - d	206(181-234)	292(129-449)	0.845
LAD - mm	41.4(38.7-44.1)	37.4(33.7-42.9)	0.0817
LAV - ml	62.5(57.5-67.3)	45.2(38.0-51.6)	0.0016
LAVI - ml/m ²	34.0(32.6-42.8)	23.7(20.9-27.3)	0.000554
LVEF - %	41.1(35.7-54.1)	64.1(61.8-66.5)	0.000018
LV end-diastolic diameter - mm	49.8(48.5-58.7)	46.9(43.1-49.9)	0.00845
LV end-systolic diameter - mm	37.7(35.5-50.5)	31.2(28.1-33.7)	0.00118
Moderate or severe MR	2(22.2)	1(4.5)	0.195
NT-proBNP - pg/ml	337(72-1242)	104(57-134)	0.344
Recurrence of AF	2(22.2)	7(31.8)	0.689
Time from ablation to recurrence - d	635	176(105-397)	0.667

Abbreviations: LAD, left atrial diameter; LAV, left atrial volume; LAVI, left atrial volume index; LVEF, left ventricular ejection fraction; LV, left ventricular; MR, mitral regurgitation; AF, atrial fibrillation