

**Table 3.** The mean IIEF, ICIQ-MLUTS, SF-36 and their subgroup scores of the patients in group 1 and 2 before and after the cystoscopies.

	Rigid Cystoscopy						Flexible Cystoscopy					
	Group 1 (n=20)			Group 2 (n=21)			Group 1 (n=20)			Group 2 (n=21)		
	Before cystoscopy Mean ± SD	After cystoscopy Mean ± SD	p value	Before cystoscopy Mean ± SD	After cystoscopy Mean ± SD	p value	Before cystoscopy Mean ± SD	After cystoscopy Mean ± SD	p value	Before cystoscopy Mean ± SD	After cystoscopy Mean ± SD	p value
IIEF- EF	14.20 ± 9.05	13.60 ± 10.05	0.17	12.21 ± 9.52	13.38 ± 8.51	0.27	13.95 ± 10.06	14.20 ± 9.05	0.64	13.38 ± 8.51	12.42 ± 8.84	0.30
IIEF- Total	36.45 ± 20.08	34.50 ± 21.34	0.16	31.25 ± 19.17	29.04 ± 19.87	0.45	34.45 ± 22.96	36.45 ± 20.08	0.82	29.04 ± 19.87	26.80 ± 20.13	0.06
ICIQ-MLUTS-Voiding	3.55 ± 4.90	3.40 ± 4.46	0.76	5.04 ± 6.27	5.00 ± 4.56	0.73	3.20 ± 3.27	4.20 ± 4.88	0.11	4.57 ± 4.23	4.19 ± 4.52	0.38
ICIQ-MLUTS-Incontinence	3.45 ± 5.23	3.30 ± 5.02	0.95	3.09 ± 3.92	5.80 ± 7.18	0.15	3.00 ± 3.11	4.75 ± 6.48	0.06	3.61 ± 4.75	3.80 ± 4.56	0.83
SF-Physical	83.73 ± 13.90	83.32 ± 14.17	0.88	70.75 ± 21.57	72.61 ± 20.22	0.75	78.61 ± 17.22	83.73 ± 13.90	0.13	72.61 ± 20.22	74.21 ± 20.28	0.27
SF-Mental	74.52 ± 15.00	75.98 ± 13.76	0.58	65.17 ± 18.30	71.19 ± 15.21	0.08	72.10 ± 16.50	74.52 ± 15.00	0.24	71.19 ± 15.21	70.45 ± 17.88	0.10
SF- Physical functioning	79.50 ± 21.02	83.25 ± 20.08	0.40	70.71 ± 28.11	68.57 ± 27.84	0.59	74.00 ± 25.98	79.50 ± 21.02	0.12	68.57 ± 27.84	72.38 ± 25.13	0.28
SF-Role functioning/physical	90.00 ± 40.06	92.50 ± 36.36	0.70	65.47 ± 42.92	76.19 ± 36.63	0.14	83.75 ± 37.41	90.00 ± 40.06	0.41	76.19 ± 36.63	85.71 ± 38.38	0.08
SF-Pain	91.60 ± 13.32	88.00 ± 17.08	0.37	86.38 ± 18.82	85.90 ± 18.98	0.83	90.50 ± 12.24	91.60 ± 13.32	0.73	85.90 ± 18.98	82.09 ± 22.61	0.52
SF-General health	71.10 ± 14.53	67.63 ± 16.89	0.22	60.42 ± 16.16	59.80 ± 15.42	0.72	65.47 ± 15.94	71.10 ± 14.53	0.09	59.80 ± 15.42	56.66 ± 15.90	0.12
SF-Energy/Fatigue	63.75 ± 16.37	62.25 ± 17.35	0.38	58.09 ± 10.77	57.38 ± 10.77	0.65	62.25 ± 15.76	63.75 ± 16.37	0.26	57.38 ± 10.77	57.14 ± 11.78	0.77
SF- Social functioning	88.75 ± 19.40	87.50 ± 20.27	0.70	77.97 ± 25.58	79.76 ± 24.83	0.72	91.25 ± 14.67	88.75 ± 19.40	0.49	79.76 ± 24.83	75.59 ± 26.36	0.33

SF-Role functioning/ emotional	$80.00 \pm 29.41$	$90.00 \pm 36.03$	0.10	$73.49 \pm 40.69$	$80.95 \pm 37.37$	0.12	$73.33 \pm 38.38$	$80.00 \pm 29.41$	0.33	$80.95 \pm 37.37$	$88.88 \pm 53.05$	0.59
SF-Mental health	$65.60 \pm 18.73$	$64.20 \pm 17.19$	0.76	$61.14 \pm 13.39$	$66.66 \pm 14.47$	0.06	$61.60 \pm 18.82$	$65.60 \pm 18.73$	0.08	$66.66 \pm 14.47$	$60.19 \pm 15.15$	0.10

IIEF: International Index of Erectile Function, EF: Erectile Function, ICIQ-MLUTS: International Consultation on Incontinence Questionnaire-Male Lower Urinary Tract Symptoms, SF: Short Form

Wilcoxon Signed Ranks Test was performed.