

Table 1. Demographic information and medical data of patients by respondents

Respondent	Adolescent (N=50)	Mother^a (N=94)	Father^a (N=48)
Age in years, mean \pm SD	15.8 \pm 2.3	9.4 \pm 5.1	8.4 \pm 5.3
Female, n (%)	24 (48.0)	44 (46.8)	27 (56.3)
Gene mutation, n (%)			
Δ F508del homozygous	11 (22.0)	16 (17.0)	10 (20.8)
Δ F508del heterozygous	14 (28.0)	23 (24.5)	14 (29.2)
Other mutations	22 (44.0)	49 (52.1)	21 (43.8)
BMI z-score, mean \pm SD	-0.8 \pm 1.7	-0.9 \pm 1.4	-1.1 \pm 1.4
FEV1% pred ^b , mean \pm SD	79.2 \pm 26.1	84.0 \pm 22.7	81.3 \pm 27.8
Comorbid conditions, n (%)			
CF related diabetes mellitus	3 (6.0)	1 (1.1)	2 (4.2)
Hepatobiliary disorder	7 (14.0)	16 (17.0)	11 (22.9)
Pancreatic insufficiency	39 (78.0)	75 (79.8)	40 (83.3)
BIPAP use	1 (2.0)	3 (3.2)	2 (4.2)
Oxygen use	1 (2.0)	1 (1.1)	2 (4.2)
Chronic MRSA infection	9 (18.0)	15 (16.0)	7 (14.6)
Chronic PA infection	25 (5.0)	30 (31.9)	15 (31.3)
^a Characteristics of CF patients whose parents completed the screening.			
^b pred = predicted			