

Table 1. Baseline participant characteristics

	<i>P. aeruginosa</i> group			MRSA group		
	IV (N=79)	Non-IV (N=23)	p-value	IV (N = 36)	Non-IV (N = 12)	p-value
Diagnosis age (year), mean ± SD	1.5 ± 3.3	2.2 ± 4.4	0.483	1.1 ± 2.4	1.9 ± 2.8	0.255
Age at isolation (year), mean ± SD	5.1 ± 4.9	8.1 ± 5.8	0.02*	6.7 ± 5.1	10.4 ± 6.6	0.078
Female, n (%)	37 (46.8)	14 (60.9)	0.236	19 (52.8)	7 (58.3)	0.738
Pancreatic insufficiency, n (%)	67 (84.8)	20 (87)	1	31 (86.1)	9 (75)	0.314
CF related diabetes, n (%)	1 (1.3)	0 (0)	1	0 (0)	0 (0)	-
Hepatobiliary disease, n (%)	15 (19)	4 (17.4)	1	9 (25)	6 (50)	0.106
ABPA, n (%)	2 (2.5)	1 (4.3)	0.539	1 (2.8)	0 (0)	1
Bipap use, n (%)	2 (2.5)	0 (0)	1	1 (2.8)	0 (0)	1
Oxygen use, n (%)	0 (0)	0 (0)	-	1 (2.8)	0 (0)	1
Inhaled corticosteroid, n (%)	16 (20.3)	4 (17.4)	1	10 (27.8)	1 (8.3)	0.248
Dornase alfa, n (%)	78 (98.7)	22 (95.7)	0.402	35 (97.2)	12 (100)	1
* p < 0.05						