

TABLE 1 : Baseline characteristics by study groups

	CM (n=59)	MM (n=165)	P value
Male	40 (67.8)	105 (63.7)	0.566
Age, yrs	70.9 ± 8.1	69.6 ± 7.5	0.280
BMI	24.6 ± 2.6	25.0 ± 3.4	0.478
Combined procedure	27 (45.8)	112 (67.9)	0.003
PAF in isolated LAAC	0 (0)	4 (7.5)	0.775
PAF in combined procedure	15 (55.6)	56 (50)	0.863
Mean CHA ₂ DS ₂ -VASc score	4.1 ± 1.5	4.3 ± 1.5	0.401
Mean HAS-BLED score	2.8 ± 1.2	2.6 ± 1.0	0.227
Existing comorbidities, n (%)			
Previous stroke/TIA	30 (50.8)	108 (65.5)	0.048
Hypertension	48 (81.4)	124 (75.6)	0.333
Diabetes mellitus	14 (23.7)	46 (27.9)	0.537
CAD	11 (18.6)	35 (21.2)	0.675
Echocardiographic characteristics			
LAD, mm	48.8 ± 6.2	48.0 ± 6.2	0.358
LVEF, %	50.2 ± 5.7	49.5 ± 5.5	0.425

Values are mean ± SD or n (%).

BMI: body mass index; PAF: paroxysmal atrial fibrillation; CAD: coronary artery disease; LAD: left atrial diameter; LVEDD: left ventricular end-diastolic dimension; LVEF: left ventricular ejection fraction; LAAC: left atrial appendage closure.