

TABLE 1 Clinical Characteristics of Patients with Retinoblastoma Having Hepatic Metastases

		Case 1	Case 2	Case 3	Case4	Case5
Gender		Male	Male	Female	Male	Male
Age at first symptom (months)		12	66	16	20	16
Symptom	leukocoria	√	√	√	√	√
	strabismus	√	√			
	visual impairment	√	√			√
	exophthalmos			√		√
	Peri-orbit neoplasm		√		√	√
Laterality		Left	Right	Bilateral	Bilateral	Left
Family history		No	No	No	No	No
Interval between first symptom and initiation of treatment (months)		7	0	1	0	2
Interval between first symptom and hepatic metastasis (months)		14	22	1	41	8
IIRC stage	Right	/	E	E	C	/
	Left	E	/	E	E	E
CT features	Single focal or not	Multiple	Single	Single	Multiple	Multiple
	Calcification	√	×	×	×	×
	Density	Low	Low	Low	Low	Low
	Enhancement	No	yes	No	No	Yes
	Post-treatment change	Decreased	Decrease d	Decrease d	No reaction	Decreased
Involvement of other systematic organs	CNS	√	√	√	√	√
	Bone	√	√	√	√	√
	Bone marrow	√	√	×	√	√
	Lymph node	√	×	√	√	√

IIRC, International Integrated Reporting Council; CT, computed tomography; CNS, Central Nervous System