

**Table 1.**

	<b>Male (N=209)</b>	<b>Female (N=101)</b>	<b>Total (N=310)</b>	<b><i>p</i> value</b>
<b>Obstructive Sleep Apnea Syndrome (OSAS)</b>	109 (52.2%); 5 also had PLMD	50 (49.5%); 2 also had PLMD	159 (51.3%); 7 also had PLMD	0.661972
<b>Primary Snoring</b>	57 (27.3%)	24 (23.8 %)	81 (26.1%)	0.509682
<b>Periodic Limb Movement Disorder (PLMD)</b>	19 (9.1%) (5)	13 (12.9%) (2)	31 (10%) (7)	0.30522
<b>Idiopathic hypersomnia</b>	8 (3.8%)	4 (4.0%)	12 (3.9 %)	1
<b>Central apnea</b>	6 (2.9%)	2 (2.0%)	8 (2.9%)	1
<b>Enuresis</b>	6 (2.9%)	1 (1.0%)	7 (2.3%)	0.4339
<b>Bruxism</b>	1 (0.5%)	4 (4.0%)	5 (1.6%)	0.0404.
<b>Sleep tremors</b>	3 (1.4%)	2 (2.0%)	5 (1.6%)	0.6622
<b>Narcolepsy</b>	3 (1.4%)	0 (0%)	3 (1.0%)	0.5535
<b>Sleep seizures</b>	1 (0.6%)	2 (2.0%)	3 (1.0%)	0.2485
<b>Sleep walking</b>	0	1 (1.0%)	1 (0.3%)	0.3258
<b>Insomnias</b>	1 (0.6%)	0	1 (0.3%)	1