

# 2016 Monsoon Convection and its place in the Large-Scale Circulation using Doppler Radars

A. J. Doyle<sup>1</sup>, T. H. M. Stein<sup>1</sup>, A. G. Turner<sup>1,2</sup>

<sup>1</sup>Department of Meteorology, University of Reading, Reading, UK

<sup>2</sup>National Centre for Atmospheric Science, University of Reading, Reading, UK

## Key Points:

- Indian Doppler Weather radars are used to analyse cell-top height over the 2016 monsoon season
- Cell-top height exhibits clear diurnal and intraseasonal variation
- Cell-top height varies by region and is affected by local features and the wider monsoon circulation

---

Corresponding author: Alex Doyle, [a.j.doyle@pgr.reading.ac.uk](mailto:a.j.doyle@pgr.reading.ac.uk)

## Abstract

Convective cloud development during the Indian monsoon helps moisten the atmospheric environment and drive the monsoon trough northwards each year, bringing a large amount of India's annual rainfall. Therefore, an increased understanding of how monsoon convection develops in observations will help inform model development. In this study, 139 days of India Meteorological Department Doppler weather radar data is analysed for 7 sites across India during the 2016 monsoon season. Convective cell-top heights (CTH) are objectively identified through the season, and compared with near-surface (at 2 km height) reflectivity. These variables are analysed over three time scales of variability during the monsoon: monsoon progression, active-break periods and the diurnal cycle. We find a modal maximum in CTH around 6–8 km for all sites. Reflectivity increases with CTH, at first sharply, then less sharply above the freezing level. Bhopal and Mumbai exhibit lower CTH for monsoon break periods compared to active periods. A clear diurnal cycle in CTH is seen at all sites except Mumbai. The phase of the diurnal cycle depends on the surface type being land or ocean for south-eastern India, with the frequency of oceanic cells typically exhibiting an early morning peak compared to those over land, consistent with the observed diurnal cycle of precipitation. The cell characteristics discovered are discussed in light of the differences in large-scale synoptic and mesoscale mechanisms responsible for different cell regimes. Our findings confirm that Indian monsoon convective regimes are partly regulated by the large-scale synoptic environment within which they are embedded.

## 1 Introduction

The Indian summer monsoon (ISM) is responsible for one of the largest modes of seasonal variability in wind direction and precipitation seen worldwide, and brings more than 80% of India's annual rainfall during the Boreal summer months of June to September. This fact, along with the vast, growing population across South Asia whose livelihoods depend upon monsoon rains, makes it one of the most active areas of research in the atmospheric sciences. The ISM is a complex phenomenon, exhibiting substantial variability on intraseasonal and interannual time scales, and monsoon forecasts remain a challenge as a result of this complexity. S. A. Rao et al. (2019) raised three areas of improvement required in numerical weather prediction (NWP) models for improved ISM forecasts: resolution, physics, and ocean coupling. Whilst substantial improvements have been made in recent years, meaningful forecast skill exists only out to four days (Turner et al., 2011; Durai & Roy Bhowmik, 2014; S. A. Rao et al., 2019). The representation of monsoon clouds in NWP models is one of the most challenging physical aspects of the ISM due to complex and numerous sub-gridscale processes. General circulation models (GCMs) used for forecasting have resolutions too large to explicitly capture convective processes, a crucial component of the monsoon. Errors due to parametrization of deep convection in GCMs are quick to grow over time (e.g. Martin et al., 2010). A more detailed view of the monsoon circulation and underlying processes from observations is necessary to better comprehend the intimate coupling between the land surface, boundary layer, and cloud development (Turner et al., 2019). In this vein, we present here an observation-driven analysis of the ISM's distinctive cloud regimes in relation to the large-scale environment.

ISM cloud development has further implications for monsoon progression. On average, the monsoon arrives at the tip of south-eastern India on 1 June; however, the monsoon onset does not necessarily fall on the same date each year. The leading edge of the monsoon then propagates gradually northwards and westwards, typically reaching the north-west of the country and neighbouring Pakistan by mid-July. By September, the monsoon weakens and withdraws gradually from north-western regions, moving back south-east. Developing convection over India during early Boreal summer is tied to the progression of the ISM (Parker et al., 2016; Menon et al., 2018). During the early stages of

the monsoon at a given location, convection is typically shallow in nature, with moist tropical air increasing the relative humidity near the surface. However, this moisture is initially confined to the lower levels, with any convection vertically capped as a result. This is attributed to a mid-level wedge of dry air at approximately 600 hPa, near the freezing level, with winds originating from the desert north-west (Parker et al., 2016). However, moisture starts to penetrate at this level from below with the arrival of early monsoon convection (Menon et al., 2018). Johnson et al. (1996) suggested the increase in moisture near the freezing level in the tropics is likely associated with the direct results of melting of ice precipitation. Menon et al. (2018) postulated that due to a mix of surface moistening and atmospheric moistening through additional convection, this moisture then propagates towards the surface, driving the monsoon forwards, perpendicular to the prevailing low-level winds. Furthermore, the advance of the tropical low-level circulation against the retreating dry mid-level north-westerlies as described above is not a steady process (Volonté et al., 2019) and exhibits significant variation on several spatiotemporal scales. Therefore, an enhanced understanding of monsoon cloud development in relation to the large-scale environment that it is embedded within is needed.

Part of the inherent challenge in ISM prediction is that monsoon rainfall stems from a large range of cloud sizes, from individual single-cell storms to large mesoscale convective systems (MCS) and monsoon depressions. Large-scale variability in fields such as temperature, winds, and moisture content all directly affect cloud development. This study will explore cloud regimes and the large-scale environment in relation to three time scales of variability that play a primary role in the ISM. Firstly, monsoon propagation from the south-east to the north of India each year (and its retreat again later in the Boreal summer) is non-steady with substantial interannual variability in arrival times. Accurate modelling of monsoon arrival in a region relies upon a precise understanding of the large-scale wind pattern changes that control monsoon propagation. Secondly, systematic variation in convection exists as a result of active and break periods (Rajeevan et al., 2010), a key component of intraseasonal variability during the ISM. These occur as a manifestation of fluctuations in the Boreal summer intraseasonal oscillation, the northward movement of clouds and convection from the equator to the Indian monsoon region (e.g. Jiang et al., 2004), varying on 30–60 day time scales. Models have been found to lack sufficient variability on intraseasonal time scales over both land and ocean (Martin et al., 2017). Thirdly, monsoon clouds typically exhibit a strong signal on diurnal time scales associated with the solar cycle and heating of the surface. However, the diurnal cycle is poorly modelled in the tropics, leading to underestimates in daily mean rainfall (Martin et al., 2017). Currently, there is limited understanding on the main drivers of monsoon convection in different locations, and the relative importance of different factors such as local orography, proximity to the coast, prevailing wind direction, amongst others.

To characterise cloud types, we will use data from a network of operational Indian Doppler Weather radars, managed by the India Meteorological Department (IMD). Similar studies have been performed in the past on radars in the tropics. V. V. Kumar et al. (2013) analysed convective cloud-top height, objectively defined using the C-band polarimetric (CPOL) research radar in Darwin, northern Australia. They related four discrete convective cumulus modes to rainfall events, in an effort to understand what pattern of cloud structures and their temporal progression correlates to the most rainfall. This showed that cloud-top height can be used as a robust proxy for convective cell development, and using cloud-top height statistically in this manner forms the basis of the present study. The IMD Doppler radars have also been used before for ISM cloud research. Sindhu and Bhat (2018) analysed the dynamics and microphysics of monsoonal MCSs using four IMD operational S-band radars, chosen to represent four different climate regions. They also used storm area and height to understand the temporal progression of storms. Following this, Sindhu and Bhat (2019) then looked at monsoon storm statistics and the robustness of radar-derived precipitation estimates using an IMD C-band radar at New Delhi. They subset storms into simple (less intense) and more intense

categories, and found a positive relationship between maximum echo-top height (ETH) and precipitation amount. However, this relationship was found to be weaker for the less intense storms. Here, we build on these studies by comprehensively analysing statistics of cloud height across multiple sites for an entire season, allowing a direct comparison of monsoon cloud regimes in different climate regions.

The overarching aim for this study is to find the spatiotemporal characteristics of convective clouds across India during the monsoon season in 2016, which coincides with the period of radar data available to us (15 May to 30 September 2016). The 2016 season was the year of the INCOMPASS field campaign (Interaction of Convective Organisation with Monsoon Precipitation, Atmosphere, Surface and Sea; Turner et al., 2019), which enabled the provision of these data, and is considered a representative year for analysis (June–September all-India rainfall was 97% of the long-term average). To summarise, local radar-derived fields of cell-top height (CTH) and near-surface reflectivity are calculated for the 7 radars across the entire 2016 monsoon season and statistically analysed. Patterns in these cells are interpreted over major time scales of variability during the ISM, encompassing the seasonal cycle, active and break events and the diurnal cycle in 2016. Finally, these patterns are related to the large-scale monsoon circulation, in order to assess large-scale patterns of cloud development.

The paper is structured as follows: section 2 explains in detail how the radars are used to calculate CTH and describes any auxiliary datasets used; section 3 presents the season-average cell statistics in the context of the large-scale environment; section 4 assesses cell patterns during 2016 active and break periods; section 5 examines the diurnal cycle in cell activity at each location, and investigates how it compares between land and ocean, and how it is modulated by monsoon progression and active-break spells. Finally, the main conclusions of this study and key areas requiring future investigation are presented in section 6.

## 2 Data and Methods

In the following section we introduce the radar data, first detailing the procedure toward analysing radar reflectivity on discrete vertical levels. Then, objective definitions for ETH and CTH are given. Additional relevant methods and datasets are specified in section 2.3.


### 2.1 Radar data

Access to 139 days of reflectivity data (15 May to 30 September 2016) from 7 Indian Doppler Weather radars (see Figure 1a) will allow for investigation of individual clouds during the 2016 monsoon season across India. These 7 radars are part of a wider network managed by the IMD, and are chosen for analysis as they are all S-band (approx. 10 cm wavelength), are robustly calibrated (see below) and represent 4 distinct climate regions with different monsoon rainfall climatology and arrival dates. The south-east sites (Chennai, Machilipatnam) represent coastal India, and are outside the core monsoon circulation. Mumbai in western India represents the west coast and Western Ghats region. Northern India (Bhopal, Lucknow) represents inland regions within the core monsoon circulation. North-eastern India (Agartala, Kolkata) is a near-coastal region where the monsoon winds typically begin to curve back to the north (Figure 1b). Two radars per region were identified for analysis if possible, to allow comparison within each region. However, Mumbai was the only radar with data available on the west coast.

Each radar volume scan consists of 10 elevation angles from  $0.2^\circ$  to  $21^\circ$ , and takes approximately 7 minutes, repeating every 10 minutes. The beamwidth is approximately  $1^\circ$  and the radar gatesizes are 500 m for all radars except Mumbai, which has a 300 m gatesize. Geographical coordinates and further technical specifications of each radar are



shown in Table 1. A rigorous calibration procedure is applied for each radar following the methods of Warren et al. (2018) and Louf et al. (2019), with the basic procedure as follows. Ground clutter (available from the total power data) is used to establish changes in calibration for each day over the season. Variability in the ground clutter distribution is corrected to an arbitrary reflectivity baseline, i.e. the relative calibration adjustment Louf et al. (2019). The corrected radar data is then compared to the Global Precipitation Measurement (GPM) Satellite Ku-band radar product to assess the absolute calibration (Warren et al., 2018) over the season. The median difference between the radar and satellite reflectivity is averaged over several volume matches for each site. This offset is applied to the reflectivity data, which is iteratively corrected against GPM until the difference is calculated to be less than 1 dB, at which point the radar is assumed to be calibrated to within that error. Further quality control procedures are applied to the radar data automatically by the IMD. This includes the removal of ground clutter and anomalous propagation echos in the reflectivity data by specialist software, and is described in more detail in Roy Bhowmik et al. (2011).



**Figure 1.** (a) Topographical map of India and surrounding regions, with some major features labelled. Also shown is the network of 17 Indian Meteorological Department Doppler weather radars made available to INCOMPASS, with diamond markers, and rings denoting 100 km range from the radar. The blue rings (with bold labels and red markers) are the 7 radars used in this study. The network has since been expanded further. (b) Recent climatological (2008–2017) precipitation (India only) and 850 hPa winds from IMD gridded data and ERA5 respectively. Regions above 1.5 km in altitude are shaded grey.

The primary method surrounds the construction of a vertical profile of radar data in Cartesian coordinates - also known as a Constant Altitude Plan Position Indicator (CAPPI) - transformed from its native polar coordinates. This grid is constructed using the Wradlib module for Python (Heistermann et al., 2013). The Cartesian grid consists of 28 vertical levels at 0.5 km resolution from a base of 2 km, chosen to be above any clutter near the surface but below the freezing level so as to avoid brightband effects (usually at around 5 km over India during the Boreal summer; Harris Jr et al., 2000).

A range of 40–100 km is considered to avoid the ‘cone of silence’ close to the radar up to the maximum height of the grid at 16.0 km, and because of increasing noise level for the radars as range increases as well as beam broadening effects. Large cumulonimbus towers will often exceed 16 km during the monsoon season. However, the largest elevation angle in the IMD radar scanning strategy is  $21^\circ$ . The centre of this beam will intersect the 16 km level when it is a horizontal distance of approximately 40 km from the radar. Any further restrictions in the size of our horizontal domain in order to capture higher clouds would reduce the number of sampled cells. As we have only one season of data, we opted to prioritise the size of the horizontal domain. Furthermore, with only ten elevation angles, the gaps between sweeps becomes increasingly large with height, giving lower confidence in cell reflectivity at higher altitudes. Cells penetrating above this threshold are designated a CTH of 16.5 km, though we do not know the true CTH.

The horizontal resolution of the grid is  $1 \text{ km} \times 1 \text{ km}$ . At each elevation, for each Cartesian pixel, we find all grid points within 3 km and calculate the average reflectivity, weighted by the inverse square of its distance from the radar site. This distance was chosen as a trade-off between a smooth Cartesian grid of reflectivity and the retention of original features. Warren and Protat (2019) recommends interpolation in dBZ for low reflectivity signals such as ETH. As ETH is a major part of this study, we interpolate in dBZ units here. Whilst Wradlib was used to construct the Cartesian 3-D grid, its CAPPI algorithm introduced significant artificial annular ring features at regular intervals around the radar above and including the brightband region. These occur as a result of increasingly large spatial gaps between sweeps and the effects of melting precipitation. We therefore apply our own algorithm, whereby the vertical profile (0.5 km grid) is calculated by linearly interpolating vertically across all elevation scans for each Cartesian pixel. This removes most of the annular ring artefacts, although some anomalies remain present after CAPPI construction, especially in the brightband (5–7 km) region and at higher altitudes.

## 2.2 Derivation of echo-top height and cell-top height

To calculate radar-derived ETH, that is to say, the highest observed altitude of reflectivity for each pixel, a threshold for what constitutes a meteorological value must be applied. This is dependent on the noise level of the radar, which itself is dependent on range. The choice of threshold must be greater than the noise level at 100 km (the outer-edge of our domain). This threshold is set to 5 dBZ, and is higher than the minimum detectable signal for all radars at 100 km, as shown in Table 1. Specifically then, searching upwards from 2 km, the pixel nearest to 5 dBZ in the vertical column is taken as the ETH, with the condition that there must be retrievals greater than 5 dBZ at each 0.5 km level in the vertical column below the ETH. This approach is similar to that of V. V. Kumar et al. (2013).

Next, we define cell-top height (CTH), which is the maximum ETH in a convective region. A convective region is any connected region of convective pixels, which is determined from the base CAPPI at 2 km following the method of Steiner et al. (1995):

1. Any pixel  $\geq 40$  dBZ is automatically considered convective.
2. Any pixel greater than a background state by the reflectivity difference depicted in Figure 7 of Steiner et al. (1995) is convective.
3. Each convective pixel has a convective radius of a certain size as a function of intensity. A cell then is any connected set of convective pixels, and hence is termed a cell here, as a typical cumulonimbus cloud would have convective and stratiform segments.

The assumption is that if a pixel is convective at 2 km, it is convective throughout its vertical extent. Due to small 2–12 km vertical shear at all sites during the mon-

**Table 1.** Relevant specifications for the seven Doppler radars used in this study, grouped by region. The noise at 100 km is the minimum detectable signal calculated using a random day of clear-sky data for each radar. A clear-sky day is simply determined as any day with no measured rainfall that is also persistently clear in infra-red satellite imagery.

Site <sup>a</sup>	Lat (°N)	Lon (°E)	Alt (m)	Gatesize (m)	Beamwidth (°)	Noise at 100 km (dB)
Chennai (VOMM)	13.07	80.29	35	500	1.00	2
Machilipatnam (VOMP)	16.18	81.15	35	500	1.00	2
Bhopal (VABP)	23.24	77.42	570	500	0.93	0
Lucknow (VILK)	26.77	80.88	143	500	0.97	3
Agartala (VEAT)	23.89	91.25	35	500	0.93	-1
Kolkata (VECC)	22.57	88.35	35	500	1.00	2
Mumbai (VABB)	18.90	72.81	100	300	1.00	2

<sup>a</sup>The elevation angles for all radars are: 0.2°, 1.0°, 2.0°, 3.0°, 4.5°, 6.0°, 9.0°, 12.0°, 16.0°, and 21.0°.

soon season ( $< 10^{-3}\text{s}^{-1}$ ), this is considered a fair assumption. Therefore, any inaccuracies stemming from tilting of the convective core due to vertical shear or cloud movement within the 7-minute radar volume are further reduced by taking all connected pixels when calculating CTH. One limitation of applying the method of Steiner et al. (1995) in this manner is that elevated convection with no convective signal at 2 km will not appear in this study. However, due to very moist conditions in the lower troposphere during the monsoon, such occurrences are assumed to be small in number.

Attenuation is considered to be negligible for S-band radars, except for the most intense convective storms where hail is present (Testud et al., 2000). Such storms do occur during the ISM but are assumed to constitute a small proportion of our large, final sample. Therefore, attenuation correction is not performed. Only cells with a surface area at 2 km of at least 4 km<sup>2</sup> (4 pixels) are included in this analysis, as cells smaller than this dominate the statistics but do not represent true convection. Finally, any cell touching the inner or outer edges of the 40-100 km domain is not considered so that we always sample entire cells.

### 2.3 Auxiliary Datasets and Method

The convective cell statistics presented are considered in the context of the local and large-scale meteorological state during the 2016 monsoon season. The Integrated Multi-satellitE Retrievals for GPM product (hereafter referred to as IMERG), provides half-hourly robust precipitation estimates (Huffman et al., 2019), at 0.1° horizontal resolution. This allows visualisation of the large-scale precipitation field.

IMD gridded daily rainfall accumulations (Rajeevan et al., 2006), with 1° spatial resolution, are used to designate active and break days in 2016 (and are also used in Figure 1b). We follow the method of Ravi Kiran et al. (2009) with a few adjustments: June–September 2016 IMD gridded daily rainfall is compared to the recent climatological (2008–2017) rainfall, averaged across a central India zone (18–27 N, 73–82 E). A standardised rainfall anomaly is then calculated for each day, defined as the climatological average subtracted from the 2016 value, divided by the standard deviation of the climatological average. A difference between each day’s standardised anomaly and the 2016 mean standardised anomaly that is greater than (less than) 0.7 (-0.7) defines an active (break) day. Finally, an active/break day must be part of a set of 3 consecutive active/break days in order to be included in our statistics. The active and break days derived in this manner may not necessarily be part of true active and break periods, but instead more closely represent active and break spells with respect to the 2016 mean state, and their corresponding large-scale pattern changes. As a result, we identify a larger and more equal number of active and break

days, lending itself to a more robust and fair comparison. The final active and break days used for this study are presented in Table 2. Handily, there are 32 active and 32 break days across the season (26% of the full sample in each case).

ERA5 (ECMWF Re-Analysis) estimates of several atmospheric variables at multiple levels are also used to complement this analysis over the 2016 season (Hersbach et al., 2018). Owing to the hourly resolution of ERA5, the diurnal cycle can be explicitly determined. Finally, twice daily (00 UTC and 12 UTC) atmospheric soundings are available in the near vicinity ( $< 25$  km) of all sites analysed here except Machilipatnam, and can be freely downloaded from the University of Wyoming Atmospheric Soundings web-page (<http://weather.uwyo.edu/upperair/sounding.html>). As an alternative for Machilipatnam, sounding data 280 km up the coast in Vishakapatnam (see Figure 1a) is used. Naturally, this should then be viewed with more caution when making direct comparisons. In the case of both IMERG and ERA5, when comparing fields at local radar sites, the average of all grid-boxes with centroids in the 40–100 km radar domain is used.

**Table 2.** 2016 Indian monsoon season active and break spells. The method used to obtain these dates is described in the text.

Date	Active/Break
1–3 June	Break
9–17 June	B
28–30 June	Active
2–7 July	A
9–13 July	A
20–26 July	B
2–4 August	A
12–14 August	B
19–21 August	A
3–12 September	B
14–17 September	A
21–28 September	A

### 3 Individual Cell Statistics

We start by examining the number of cells of different heights for each site, and relate these patterns to near-surface cell reflectivity. Table 3 shows the number of cells at each site and as an average per radar volume. Agartala has by far the largest number of cells, situated in the very convectively active north-eastern India region, with prevailing low-level winds originating from over the very moist Bay of Bengal (Figure 1b). Mumbai, unfortunately, has noticeably fewer volumes over the season, owing to substantial periods of missing data from July onwards, and so should be analysed with more caution. Considering the ratio of number of cells to number of radar volumes as a measure of convective activity (fourth column of Table 3), Agartala is by far the most convectively active, then all other sites average approximately 3 or 4 cells per radar volume over the entire season.

The probability of cell occurrence in each 0.5 km CTH bin is shown in Figure 2a. There is consistency between all 7 sites in the shape of this function, with a peak in CTH at approximately 6–8 km, around or just a bit above the average freezing level during the monsoon (Harris Jr et al., 2000). The two northern India sites - Bhopal and Lucknow - have the maximum CTH occurrence at a greater height, as well as the highest proportion of very deep (CTH  $> 10$  km) cells. Several sites (Bhopal, Lucknow, Chennai, Machilipatnam) display the hint of a secondary peak in CTH at 13–14 km, associated with deep convection. However, this peak is not obvious, as Figure 2 is an average over the entire season, throughout which different cloud regimes, some dominated by shallower convection, will occur. Indeed, taking especially convectively active weeks displays the secondary peak much more clearly (not shown). Figure 2a suggests the presence of at least 2 cumulus modes, one around the freezing level, and one in the upper troposphere. This will be explored in more detail later in this section.

**Table 3.** Total number of complete radar volumes, the number of cells (including CTH > 16 km), and average number of cells per radar volume for all 7 sites, grouped as in Table 1, from 15 May to 30 September 2016. The percentage of cells with CTH > 16 km is also shown. Note that with a radar volume of 10 minutes, if there were 0 missing scans through the season, the number of volumes would equate to 20016 scans.


Site	Volumes	Cells	Cells per Volume	CTH > 16 km (%)
Chennai (VOMM)	17839	59009	3.3	2.9
Machilipatnam (VOMP)	18589	53822	2.9	2.3
Bhopal (VABP)	13546	53641	4.0	5.5
Lucknow (VILK)	6914	22218	3.2	6.3
Agartala (VEAT)	14711	108928	7.4	2.3
Kolkata (VECC)	13018	50654	3.9	1.9
Mumbai (VABB)	3553	11071	3.1	0.2

Mumbai exhibits a larger proportion of shallow convection (CTH < 5 km), and a smaller proportion of deep convection relative to other sites. This is consistent with previous observational analysis for the region. Das et al. (2017) found rain in the Western Ghats region to be dominated by shallow-convective systems (echo tops below the melting layer). Mid-tropospheric air from Mumbai radiosonde soundings in 2016 is found to be drier on average compared to the other regions (not shown), consistent with the findings of S. Kumar (2017). Das et al. (2017) also commented on the role of orography, showing that convective clouds over the Western Ghats precipitate out quickly owing to rapid collision-coalescence of cloud droplets due to orographic uplift.

The vertical wind profiles in Figure 2b show the presence of the tropical easterly jet (TEJ) upwards of 8 km, a major component of the large-scale monsoon circulation in Boreal summer (Krishnamurti & Bhalme, 1976). It forms as a result of the meridional temperature gradient between the subtropics and equatorial Indian Ocean in the upper troposphere. Mumbai and south-eastern India exhibits westerly or south-westerly winds in the low and mid-troposphere, with strong easterly winds associated with the TEJ in the upper troposphere. Interestingly, the transition to easterly at around 6–8 km above the surface is where the winds are weakest, and this is approximately the height of the modal CTH for these sites. The presence of stronger winds aloft associated with the TEJ may be an inhibiting factor for vertical cell growth, as the strong wind shear can spread cloud tops above 8 km (Sathiyamoorthy et al., 2004). This can act as one constraint on CTH, especially during peak monsoon months when the TEJ is strongest. For northern and north-eastern India, there is a deep layer of southerly winds in the lower and middle-troposphere, with a small amount of gradual backing of these winds with height. Wind speeds only begin to increase above 8 km, again above the modal peak in CTH for these sites. The increase in wind speed is more subtle for Lucknow (VILK), where weaker upper-level wind shear may contribute to the marginally higher proportion of very deep cells here (> 12 km), due to less spreading of cloud tops.

As aforementioned, Figure 2a suggests the presence of multiple cumulus modes (preferential detrainment of cloud tops at certain levels), and we investigate this and its associated mechanisms in more detail here, by analysing the relationship between CTH, the lapse rate in temperature (Figure 2c) and near-surface reflectivity (Figure 3). This analysis is important because the different cumulus modes in Figure 2a are likely to have different characteristics and may interact with the large-scale environment in different ways. Firstly, Figure 3 shows that 2 km cell reflectivity is clearly a definitive indicator of CTH for all sites. More intense cells typically have higher cell tops (and are greater in horizontal extent, not shown here). Reflectivity values of 20 dBZ indicate light rain (< 1 mm hr<sup>-1</sup>), 30 dBZ moderate rain (> 1 mm hr<sup>-1</sup>), and 40 dBZ heavy rain (> 10





**Figure 2.** (a) Percentage of different CTHs for the 7 radars used in this study as shown in Figure 1, from 2.5 km above the surface, every 0.5 km in height, (b) wind barbs from local soundings every 1 km in height with barb increments denoting 5, 10 and 50 knots respectively and, (c) mean temperature lapse rate every 1 km in height also from local soundings. Radars in different regions are grouped by colour. The PDF does not include CTH > 16 km (see Table 3).

mm hr<sup>-1</sup>) and so higher cells are also associated with heavier precipitation. Mumbai exhibits a higher average 2 km reflectivity for shallow CTH (27 dBZ at 4 km compared to 24 dBZ at 4 km for the composite of all sites), suggesting low cell tops are associated with heavier rainfall here. The ability for shallow convection (with warmer cloud tops) to produce high rainfall rates in this region has implications for conventional satellite rainfall algorithms which relate precipitation to cloud top brightness temperature (Shige & Kummerow, 2016). Studying Figure 3h, the shallowest cells are associated with very low reflectivity and have a broad reflectivity distribution, suggesting considerable variance in the intensity of these shallow cells. As CTH increases there is initially a rapid rise in reflectivity until approximately 30 dBZ. At this point, CTH is on average 5–7 km, in the region of the freezing level (dashed blue line) and the modal CTH over the season. This is also the approximate region of maximum atmospheric stability (as shown by the minima in the lapse rates in Figure 2c), and is associated with a narrowing of the reflectivity distribution (scanning horizontally between the black dotted lines in Figure 3). The stable layer near the freezing level forms physically as a result of latent heat release due to precipitation freezing and melting processes (Posselt et al., 2008). Above this level, there is a smaller increase in reflectivity with CTH. Furthermore, around 12 km the lapse rates in Figure 2c increase to quasi-dry adiabatic with remarkable consistency between all sites hinting at the regulatory and homogeneous influence of the large-scale monsoon pattern upon different regions in India. Around 10 km there is a small atmospheric layer of constant reflectivity with increased CTH for some of the sites (e.g. Bhopal, Lucknow, Mumbai). This is likely because clouds that penetrate to this level encounter significant atmospheric instability and thus can grow deeper without any significant increase in intensity.


Therefore, the convective modes identified from Figure 2a show distinct characteristics in terms of intensity and can be related to atmospheric stability. The first mode is shallow convection ( $\text{CTH} < 5 \text{ km}$ ), associated with highly variable, but on average low 2 km reflectivity (rapidly increasing with each 0.5 km CTH bin). The second mode represent cumulus congestus clouds (5–8 km), and is bounded by the maximum in atmospheric stability near the freezing level due to insubstantial 2 km reflectivity ( $< 30 \text{ dBZ}$ ). This is also visible as the modal peak in cells around this region for all sites (Figure 2a). The third mode penetrates above this level ( $\geq 8 \text{ km}$ ), and exhibits the most vigorous convection with 2 km reflectivity values of 30 dBZ or greater. These modes were also found by V. V. Kumar et al. (2013) for Darwin in Austral summer (in addition to an overshooting top mode which we are unable to investigate here due to 3-D grid restrictions). However, our results in contrast show more of a continuous increase in mean 2 km reflectivity as CTH increases above the freezing level. This may stem from the spreading of cloud tops by the TEJ above 8 km, except for the largest and most intense cells. To summarise, for all sites, the statically stable layer around the freezing level (Figure 2c) acts as a constraining factor upon CTH, and this manifests itself as a reduction in the gradient of 2 km reflectivity around this level (Figure 3h). When discussing the cell patterns over the diurnal cycle in section 5, we will consider the mean diurnal cycle of these separate convective modes as well. Similar analysis sub-sampled by intraseasonal variation is provided as supplementary material.

#### 4 Cell Patterns over Active and Break Periods

Active and break spells are the foremost manifestation of systematic intraseasonal variability during the ISM. These oscillations occur in most of the elements of the monsoon system, and are highly pronounced in monsoon cloudiness (Krishnamurti & Bhalmé, 1976). Here, we analyse how the active and break spells in 2016 (Table 2) affect the height of convective cells in each region. Figure 4 displays the probability of occurrence of each CTH bin for active and break periods. Bhopal, Lucknow and Mumbai form part of the central India gridbox that is used here to monitor anomalous rainfall and active and break spells. As a result, we expect to see more frequent deeper convection in active periods at these sites. Mumbai and Bhopal exhibit this most discernibly for all CTH. For example, Bhopal has  $53 \pm 2\%$  of cells with  $\text{CTH} > 8 \text{ km}$  during break periods, compared to  $75 \pm 2\%$  for active periods. Interestingly, Lucknow shows little difference between active and break CTH distributions, which we will explore further later in this section. There is little change between the distributions of Agartala and Kolkata in north-eastern India (Figure 4e–f). Chennai and Machilipatnam also show little change between active and break spells. However, Chennai does show a small signal in active spells having more frequent deep cells, with  $30 \pm 2\%$  of cells with  $\text{CTH} > 8 \text{ km}$  during break periods, compared to  $38 \pm 2\%$  for active periods. It is important to recall that only convective cells are considered here. For example, T. N. Rao et al. (2016) found a higher occurrence and fraction of stratiform rain in active spells, due to the prevalence of large, organised storms such as mesoscale convective systems. This suggests that rainfall will not only be more intense but more long-lived, consistent with the typical longer lifetimes of stratiform rains (Steiner et al., 1995).

Table 4 shows the numbers and proportions of cells in the different cumulus mode categories as defined in section 3 for active and break periods in 2016: shallow convection, cumulus congestus, and deep convection. Between active and break periods, similar proportions of cells in different modes are seen in Machilipatnam, Lucknow, Agartala, and Kolkata (consistent with Figure 4). An increase in the number of cells per radar volume from break to active periods is unanimous across all sites, but is most significant in Mumbai, with a doubling in the percentage of deep convection. In Bhopal, deeper convection is also more likely in active spells, but the overall amount of convection is only






**Figure 3.** PDF of 2 km cell average reflectivity using a bin size of 2 dB and as a function of CTH (not including CTH > 16 km). (a) Chennai, (b) Machilipatnam, (c) Bhopal, (d) Lucknow, (e) Agartala, (f) Kolkata, (g) Mumbai, and (h) a composite of all sites. The black solid line is the mean reflectivity at each CTH, with dotted lines denoting the 5th and 95th percentiles. The dashed horizontal blue line is the height of the average freezing level from local radiosonde profiles.

marginally higher. Chennai sees a reduction in the likelihood of the cumulus congestus mode during active periods, and a higher likelihood of shallow and deep cells.

It is useful to analyse these statistical patterns in the context of the large-scale circulation so as to understand what dynamic component of active and break spells drives convective variability in different locations. Anomalies in IMERG rainfall, ERA5 winds, and relative humidity between 2016 active and break periods are shown in Figure 5, along with contours of average mean sea-level pressure (MSLP). The rainfall anomalies (Figure 5a–b) provide the closest relationship to CTH, with a narrow band of increased rainfall down the Western Ghats region and a larger swathe over central and northern India during the active spells. The effect of convective enhancement over the Western Ghats means this positive rainfall anomaly remains confined to the coastal region south of 18°N,



**Figure 4.** Cumulative distribution function of CTH (0.5 km bins) for 2016 active (blue) and break (red) periods (see Table 2) for each site as in Figure 3, including CTH > 16 km. Vertical lines and their width represent the number of samples at each CTH bin. The uncertainty bounds (too small to be clearly seen on most subplots) are calculated from the Dvoretzky–Kiefer–Wolfowitz inequality ( $\sqrt{\frac{\ln 2\alpha}{2n}}$ ), where  $n$  is the sample size and  $1 - \alpha$  is the probability the true distribution lies within the uncertainty bounds. Here,  $1 - \alpha = 0.997$ .

despite anomalously strong westerly winds and high relative humidity extending further east (Figure 5c–d). This explains the lack of a definitive change in the distribution of CTH between active and break periods over south-eastern India (Figure 4). At 26.8°N, Lucknow falls just north of the region of increased rainfall during active spells. Over central India there are relatively much stronger westerlies during active periods. These winds then start to curve back towards the north-west over the Bay of Bengal, and so Lucknow experiences south-easterly winds in the lower troposphere, as it is positioned north of the monsoon trough. This results in less rainfall than the region of potent westerlies further south. In the transition to break periods the monsoon trough moves northwards to

**Table 4.** For each site, the number of cells in each cumulus mode category for 2016 active and break periods and the average number of cells per radar volume. The numbers in brackets show the percentage of cells in that cumulus mode. Includes cells with CTH > 16 km. Shallow mode (CTH < 5 km); congestus mode ( $5 \leq \text{CTH} < 8$  km); deep mode (CTH  $\geq 8$  km).

Site	Active/Break	Shallow	Congestus	Deep	Cells per Volume
Chennai (VOMM)	Break	807 (8)	5616 (57)	3474 (35)	6.4
	Active	1242 (12)	5067 (47)	4471 (41)	7.0
Machilipatnam (VOMP)	B	842 (12)	3192 (44)	3169 (44)	5.3
	A	2581 (14)	8151 (45)	7456 (41)	6.8
Bhopal (VABP)	B	478 (5)	3140 (32)	6130 (63)	8.5
	A	287 (2)	2197 (18)	9637 (80)	9.1
Lucknow (VILK)	B	241 (5)	1817 (37)	2899 (58)	5.9
	A	422 (7)	1841 (32)	3413 (60)	7.0
Agartala (VEAT)	B	4167 (16)	9964 (39)	11305 (44)	9.4
	A	3557 (16)	9153 (40)	10223 (45)	10.8
Kolkata (VECC)	B	1877 (16)	4836 (41)	5141 (43)	8.7
	A	2416 (17)	5581 (39)	6195 (44)	11.4
Mumbai (VABB)	B	711 (37)	846 (44)	354 (19)	5.8
	A	295 (13)	1082 (49)	836 (38)	17.7


the foot of the Himalayas (shown by the northward push in the MSLP pattern from Figure 5e to 5f.), bringing lower tropospheric westerlies further north to Lucknow. However, the background state is drier overall (Figure 5d), thus Lucknow exhibits no significant change in rainfall and the distribution of CTH. In north-eastern India, it is an even more complex picture, with small pockets of positive and negative rainfall anomalies in the region during 2016. To the east of Agartala (91.25°E) along the Bay of Bengal coast, there is a more definitive area of increased rainfall and higher relative humidity during break periods, associated with enhanced southerly winds from the Bay of Bengal.

## 5 Cell Patterns over the Diurnal Cycle

The diurnal cycle is intrinsic to convection, as well as the local meteorology and land surface features of the region, and so its monsoonal behaviour is important to understand. With a horizontal resolution of 1 km and a radar volume of 10 minutes, the diurnal cycle of convection is fully captured by the radar observations. Furthermore, Chennai, Machilipatnam and Mumbai radar domains include both coastal land and ocean areas (Figure 1), allowing an analysis of the diurnal cycle over these different surface types. Finally, we will then investigate how the diurnal cycle is modulated by the progression of the monsoon through the season, and the active and break periods discussed in section 4.

### 5.1 Diurnal Cycle Cell Statistics

The diurnal cycle of convective cells at every CTH bin is shown in Figure 6 with individual convective modes shown in Figure 7. Broadly speaking, observing the number of cells (at all CTH) through the day, we see distinct differences between the four different regions (and similarities within regions). Chennai and Machilipatnam display an overall night-time peak in the number of cells. Machilipatnam appears to show two peaks, one in the afternoon and a more prolonged peak overnight. We will see later how respective land and ocean convection contributes to this. IMERG rainfall (shown by the blue line in Figure 6 as a fraction of the daily total) is fairly well correlated for both sites. Assuming that convection dominates rainfall, we expect rainfall to be correlated with convective area (which is in itself correlated with the number of cells). More precisely,




**Figure 5.** Large-scale patterns for the 32 active and 32 break days in 2016. (a–b) IMERG rainfall anomalies, relative to the entire season (c–d) 850 hPa relative humidity (colours) and wind vector anomalies, (e–f) contours of average mean sea-level pressure in units of hPa. Grey shaded regions in the centre and right columns are areas of topography over 1.5 km.

the rainfall appears to peak alongside the maximum in the number of deep cells ( $\text{CTH} \geq 8$  km, see Figure 7). As discussed in section 3, deep convective cells are associated with higher 2 km reflectivity (and thus higher precipitation rates).


The effect of solar heating is clearly a larger source of influence for the inland sites Bhopal and Lucknow, with a convective peak in the afternoon and evening associated with maximum surface heating. Noticeably, IMERG rainfall is more constant through the day in comparison, although both sites exhibit an increase in rainfall into the evening a few hours after the peak in the number of cells, but well correlated with the peak in the fraction of deep convection (Figure 7c). Initially, there are a greater number of shallow and congestus cells associated with the afternoon peak in cell fraction. Some of these cells dissipate, and others grow and merge, meaning that as the number of cells decreases in the evening, the number of deep cells and the total area covered by convection nonetheless continues to increase for a few hours.

North-eastern India sees its peak in total cell number around noon and early afternoon, however, Agartala especially is prone to convection at all times of day. This explains the much larger number of cells per radar volume shown in Table 1, with no real convectively inactive time of day. Consistently high low-level moisture from the Bay of Bengal likely contributes to this. Interestingly, the peak in Agartala rainfall actually occurs at night, the opposite time of day to the peak in cell number. Figure 6e and Fig-



**Figure 6.** Number of cells as a function of the diurnal cycle (30-minute bins) and CTH, with histograms representing the probability density of cell occurrence across each axis for (a) Chennai (VOMM), (b) Machilipatnam (VOMP), (c) Bhopal (VABP), (d) Lucknow (VILK), (e) Agartala (VEAT), (f) Kolkata (VECC), and (g) Mumbai (VABB). Includes CTH > 16 km. IST = Indian Standard Time (UTC+5.30). Pixels with 0 cells are masked white. The blue dotted line is the normalised diurnal cycle in IMERG precipitation as a mean over each IMERG gridbox whose centroid lies in the 40–100 km radar domain.

ure 7e show a higher likelihood of deeper cells at night, whereas the daytime cells, whilst higher in number, are typically shallower. This appears to be a fairly localised effect however, as it is not seen further south and west in Kolkata, which instead shows one clear peak centred at 2pm along with the peak in rainfall. Figure 7 shows that cells in north-eastern India (and also Chennai) appear to deepen more quickly than those over northern India and Machilipatnam, meaning rainfall peaks more closely alongside the peak in the number of cells. Mumbai shows no diurnal cycle in cell number or rainfall, bar the



**Figure 7.** As Figure 6, but showing the normalised fraction of cells in different cumulus mode categories: shallow, congestus and deep. The black line is the total convective area normalised over the diurnal cycle, and IMERG rainfall is as in Figure 6.

slight suggestion of a night-time and an afternoon peak. The study of Romatschke and Houze (2011) supports this result. They noted using Tropical Rainfall Measurement Mission data that over the western Indian coast, small and medium sized systems are present throughout the day, as the prevailing south-westerly moisture laden flow is lifted over orography. The prevalence of shallow convection here also moderates the daytime temperatures relative to northern India, meaning there is less of a surface heating contribution upon convective initiation.

## 5.2 Diurnal Cycle over Land and Ocean

We look in more detail at Chennai, Machilipatnam and Mumbai by considering cells over land and ocean separately. These sites have the advantage of having nearly 50% of their domains over ocean, and 50% over land. For simplicity here, cells are judged to be land or ocean cells based off their centroids, so in reality some larger cells may straddle the coast. Firstly considering Chennai and Machilipatnam, Figure 8a–b shows clearly



that there is higher rainfall and a greater cell number at night and in the early morning over the ocean compared to land. This is also a consistent pattern for all three cumulus modes (Figure S1). Either cells are simply forming later into the night on average offshore, or there is advection of cells from onshore to offshore as the evening progresses. If we consider that the 1–6 km vertical wind shear in south-eastern India is typically quasi-westerly through the day, it largely supports the advection hypothesis. That is to say, cells that form over land in the evening move with the quasi-westerly offshore wind shear. This theory is further supported by the similar shapes and amplitudes of the cycles between land and ocean. This suggests the spreading of the cloud tops over the ocean from coastal regions in the evening. This was also found for the Bay of Bengal by Yang and Slingo (2001). They suggested gravity waves and complex land-sea breeze effects as drivers of this process. On further study of Figure 8, there is a clear second peak in the height of cells (at about 13 km) over ocean, but not over land. The fact that deeper cells are more likely over ocean does not contradict the advection hypothesis. As cells move offshore in the night and early morning, they mature and thus grow in vertical extent. However, there are also plenty of shallower cells over the ocean, suggesting that convection does also initiate over the ocean itself. Over land, both sites exhibit an initial afternoon peak in IMERG rainfall, followed by a slight dip, then a second night-time peak. Machilipatnam also displays this feature in the fraction of cells. This suggests in the average situation that cells continue to initiate over land for a period of time, with a slight lull around 9pm after the initial round of afternoon convection. After midnight, convection is hindered over land following large amounts of convection in the preceding hours, and so the diurnal cycle spreads out into the ocean (Yang & Slingo, 2001).


For Mumbai, the most discernible change in the diurnal cycle is that the afternoon peak in the fraction of land cells and rainfall is not seen over ocean, which instead has a peak at night or early morning, similar to south-eastern India. This suggests that land surface heating over Mumbai, whilst smaller than the other locations, still has a significant contribution to convection. Also similarly to south-eastern India, Mumbai has a slightly higher fraction of deep cells over ocean compared to land, although this is not significant enough for a clear second peak in the CTH histogram. This may partly occur due to a smaller amount of collision-coalescence of raindrops over ocean away from the Western Ghat mountains, giving clouds a chance to grow more in vertical extent before precipitating out.

### 5.3 Diurnal Cycle during Monsoon Progression

The diurnal cycle can vary over longer time scales, such as the progression of the monsoon itself. The normalised diurnal cycle in cell number is shown for each month individually (May–September) in Figure 9. Firstly, for all sites, the diurnal cycle is often least pronounced in May, as the monsoon regime is not yet established. Chennai shows a distinct night-time peak in the fraction of cells from June onwards. However, June also has the highest likelihood of convection in the hours around midday, with very few cells around this time later in the season. The length of this daytime period of relative inactivity extends into September, by which point convection is rare between 8am and 2pm local time. As with Figure 6, Machilipatnam’s diurnal cycle is more complex, exhibiting multiple peaks from June, most definitively one centred around 4am associated with ocean convection, and one around 4pm for land convection. However, a third peak around midnight seems to occur in June and September, although this is not very distinct in IMERG rainfall. The minimum occurs around noon from July through September, and in fact these latter three months match considerably. This suggests the progression of the monsoon at this stage has less effect on the diurnal cycle of south-eastern India.

Bhopal shows most clearly the arrival and retreat of the monsoon. In Bhopal and Lucknow the monsoon arrived on approximately 20 June (Ramesh, 2017). This is associated with a more pronounced diurnal cycle from June with a peak in the fraction of






**Figure 8.** Same as Figure 6 but with separate plots for cells with their respective centroids over land and ocean. (a) Chennai, (b) Machilipatnam, (c) Mumbai. A land-ocean mask is also applied to IMERG rainfall.

convection around 4pm and extending into the evening and night. However, during peak monsoon months (July and August), convection can occur any time of day, demonstrated by a shallower peak in the afternoon, and in fact the diurnal cycle in IMERG rainfall becomes hard to visually decipher. This suggests a moister background environment more conducive for convection and less reliance upon daytime surface heating. Into September, as the monsoon gradually retreats south again, convection is once again constrained into the afternoon hours, especially for Bhopal. The development of convection from shallow to deep before precipitating out is very defined in September (Figure S3a). This convection also seems to dissipate more quickly into the evening compared to June. This suggests a greater reliance upon surface heating for the initiation of convection. A longer convectively inactive period is visible from approximately 4 to 11am for Bhopal in September. Lucknow displays comparatively more morning convection in September compared to Bhopal, with a minimum around midnight.

Agartala shows a night-time and a day-time peak in convective fraction from June to August, with relative minima in between. We recall that the night-time peak in IMERG rainfall was found to stem from a higher proportion of deeper cells at night, with a tendency for cells during the day to be shallow and cumulus congestus type modes. The proportion of deep cells at night reduces into September, with all three convective modes then seeing their peak in the early afternoon (Figure S4). Kolkata has a peak in convec-

595 tive fraction centred around 2pm for all months from June, but with convective cells and  
 596 rainfall evident at all times of day, especially during June and July.



**Figure 9.** Diurnal cycle probability density histograms showing the number of convective cells subset by month for the same sites as in Figure 6, including CTH > 16 km. However, Mumbai is not shown here due to an insufficient number of cells from July onwards. The shaded region represents the variance in the sample as the bootstrap 10th and 90th percentiles. The equivalent average IMERG rainfall diurnal cycle for each month is shown by the blue dashed line. There must be at least 500 cells during the month or the diurnal cycle is not shown.

#### 5.4 Diurnal Cycle during Active and Break Periods

597

598 It is also of interest to consider whether active and break spells may modulate the  
 599 diurnal cycles shown earlier in this section, in terms of phase and/or amplitude. Figure  
 600 10 displays the phase of the diurnal cycles as in Figure 6, but now subsampled over the  
 601 active and break days in Table 2. Chennai shows no clear change in the amplitude or  
 602 phase of its diurnal cycle between active and break periods. Further up the coast in Machili-  
 603 patnam however, there are clearly more inactive periods of convection during a break

spell than an active spell, such as during late morning. In an active spell, convection can occur all times of day (although still preferentially during the afternoon and night-time). In fact, during active periods, deep cells are most common from 3pm all the way through until around 7am as a single peak. However, during break periods, these deep cells have a shallow peak at 4pm, followed by an evening lull, then a second, larger peak around midnight. Machilipatnam is systematically affected by active and break spells in this way, despite such a signal not being evident in the overall distribution of CTH in Figure 4. This suggests that cells in Machilipatnam that do occur are not on average less intense or deep during break periods, but are more dependent on the time of the day and thus less frequent, as shown in Table 4.


Turning our attention to northern India (Figure 10c–d), active periods display a higher fraction of convection during the night-time hours in Bhopal compared to break periods. Indeed, the diurnal maximum in the congestus mode during active periods is at night, whereas the peak in the deep mode is in the afternoon (Figure S5c). For Lucknow, break periods are associated with a higher fraction of cells in late evening and early morning, but active periods display a significantly higher fraction of late morning and afternoon convection. For both sites, there is a suggestion that the peak in normalised convection is slightly earlier in the afternoon for active periods, with more of an evening peak for break periods. This likely occurs as a result of the active spell environment being more conducive for convection, such that convection initiates, deepens and intensifies more quickly in the afternoon. This effect is also supported by the IMERG rainfall for Bhopal, although it is less clear for Lucknow.

Lastly observing the respective diurnal cycles for north-eastern India, it is evident that convection can occur at all times of day during both active and break periods. For Agartala, there is a more distinct peak in cell fraction in the early afternoon during break periods, with less of a diurnal cycle in active periods, and an almost constant proportion of IMERG rainfall through the day. In contrast, during break periods, rainfall exhibits two peaks, one in the afternoon and a more significant one at night. For Kolkata, there is less of a visual difference on first inspection, however convection appears quicker to initiate around midday during active spells, and this peak is more well defined than the afternoon peak during break periods.

## 6 Conclusions

An understanding of ISM cloud development on large spatiotemporal scales is an important aspect of monsoon forecasting. This study has used 2016 monsoon season data from 7 IMD Doppler radars in order to analyse CTH in relation to near-surface reflectivity, with a focus on active and break periods, the diurnal cycle, monsoon progression, and location. Our main findings are as follows:

1. A season-average peak in CTH was found between 6 and 8 km for all sites, just above the freezing level and region of maximum tropospheric stability.
2. The seasonal distribution of CTH suggests the presence of three distinct convective cumulus modes: shallow convection, cumulus congestus, and deep convection. Considering the season as a whole, cumulus congestus is the most dominant mode. 2 km reflectivity was found to increase rapidly with CTH below the melting layer, and slowly above it.
3. There is less convection per radar volume during break periods compared to active periods for all sites, but most significantly for Mumbai. CTH is also lower on average for the 2016 break periods for Bhopal and Mumbai, spatially matched with the positive rainfall anomalies. Other sites display no definitive change in the respective CTH distributions. Low-level wind circulation changes between active and break periods are an indicative driver of resultant variability in CTH, most no-



**Figure 10.** Same format as Figure 9 but shown for break (red) and active (blue) periods. Mumbai is again not shown owing to a lack of radar volumes during the active and break spells from July onwards. The total number of cells at a site over respective break and active spells is labelled.

4. All sites but Mumbai show a clear seasonally-averaged diurnal cycle in the number of cells, with differences between each climate region attributed to local and large-scale features of the atmosphere. The peak in rainfall tends to occur around or soon after the peak in the number of deep cells.

5. The phase of the diurnal cycle and average height of cells depends on the surface type being land or ocean for south-eastern India. Ocean cells exhibit more of an early morning peak in convection and have a higher proportion of deeper cells. Mumbai also shows more of an afternoon peak over land and an early morning peak over ocean, though the difference is more subtle.
6. The diurnal cycle is further modulated by the progression of the monsoon. During peak monsoon months (July and August), the diurnal cycle becomes less distinct for most sites, with longer spells of convective inactivity pre and post-monsoon. This is most definitive for northern India.
7. Active and break periods also modulate the diurnal cycle in rainfall and cell number for several sites. Break spells tend to see convection preferentially at the more convectively active times of day, whereas active spells are associated with convection through the day (but still more frequent at certain times).

This research naturally points towards further investigation of ISM monsoon clouds and their place in the large-scale monsoon circulation. An analysis of the vertical structure of convective cells (i.e. in terms of reflectivity) associated with different large-scale regimes would be a useful next step here. It would also be beneficial to expand aspects of this analysis to additional years if data were made available, especially regarding active and break periods which exhibit significant interannual variability. Similar analyses of the intricate relationship between cloud development and the wider monsoon circulation from a satellite or modelling standpoint could also further explore some of the patterns discussed here.

The high sampling rates and spatial resolution of radars, as well as an ever increasing network of reliable radars increasing coverage across India and other tropical regions, means they are valuable resources in the analysis of storm-scale and mesoscale phenomenon. The tools (e.g. Wradlib) and method used here should prove useful as an objective way of looking at clouds over India during the monsoon, making it simple to compare between different radars as well as looking at one radar over a period of time.

## Acknowledgments

The authors gratefully acknowledge the agreement of India's Ministry of Earth Sciences in providing the Doppler weather radar under the INCOMPASS project, funded in India under the Monsoon Mission. Atmospheric sounding data were provided by the University of Wyoming (<http://weather.uwyo.edu/upperair/sounding.html>). ERA5 data were downloaded from <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels>. GPM IMERG data were downloaded from <https://gpm.nasa.gov/data/directory>. This work is supported by SCENARIO, grant number NE/L002566/1.

## References

- Das, S. K., Konwar, M., Chakravarty, K., & Deshpande, S. M. (2017). Raindrop size distribution of different cloud types over the Western Ghats using simultaneous measurements from Micro-Rain Radar and disdrometer. *Atmospheric Research*, 186, 72–82. doi: 10.1016/j.atmosres.2016.11.003
- Durai, V. R., & Roy Bhowmik, S. K. (2014). Prediction of Indian summer monsoon in short to medium range time scale with high resolution global forecast system (GFS) T574 and T382. *Climate Dynamics*, 42(5-6), 1527–1551. doi: 10.1007/s00382-013-1895-5
- Harris Jr, G. N., Bowman, K. P., & Shin, D.-B. (2000). Comparison of freezing-level altitudes from the NCEP reanalysis with TRMM precipitation radar brightband data. *Journal of Climate*, 13(23), 4137–4148. doi: 10.1175/1520-0442(2000)013<4137:COFLAF>2.0.CO;2
- Heistermann, M., Jacobi, S., & Pfaff, T. (2013). An open source library for process-



- ing weather radar data (wradlib). *Hydrology and Earth System Sciences*, 17, 863–871. doi: 10.5194/hess-17-863-2013
- Hersbach, H., De Rosnay, P., Bell, B., Schepers, D., Simmons, A., Soci, C., ... Zuo, H. (2018). Operational global reanalysis: progress, future directions and synergies with NWP including updates on the ERA5 production status. *ERA Report Series*, 27(8th October), 1–63. doi: 10.21957/tkic6g3wm
- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., & Tan, J. (2019). *Integrated Multi-satellitE Retrievals for GPM (IMERG) Technical Documentation*. Retrieved 2020-05-27, from [https://docserver.gesdisc.eosdis.nasa.gov/public/project/GPM/IMERG\\_doc.06.pdf](https://docserver.gesdisc.eosdis.nasa.gov/public/project/GPM/IMERG_doc.06.pdf)
- Jiang, X., Li, T., & Wang, B. (2004). Structures and mechanisms of the northward propagating boreal summer intraseasonal oscillation. *Journal of Climate*, 17(5), 1022–1039. doi: 10.1175/1520-0442(2004)017<1022:SAMOTN>2.0.CO;2
- Johnson, R. H., Ciesielski, P. E., & Hart, K. A. (1996). Tropical Inversions near the 0°C Level. *Journal of the Atmospheric Sciences*. doi: 10.1175/1520-0469(1996)053<1838:TINTL>2.0.CO;2
- Krishnamurti, T. N., & Bhalme, H. (1976). Oscillations of a monsoon system. Part I. Observational aspects. *Journal of the Atmospheric Sciences*, 33(10), 1937–1954. doi: 10.1175/1520-0469(1976)033<1937:OOAMSP>2.0.CO;2
- Kumar, S. (2017). A 10-year climatology of vertical properties of most active convective clouds over the Indian regions using TRMM PR. *Theoretical and applied climatology*, 127(1-2), 429–440. doi: 10.1007/s00704-015-1641-5
- Kumar, V. V., Jakob, C., Protat, A., May, P. T., & Davies, L. (2013). The four cumulus cloud modes and their progression during rainfall events: A C-band polarimetric radar perspective. *Journal of Geophysical Research: Atmospheres*, 118(15), 8375–8389. doi: 10.1002/jgrd.50640
- Louf, V., Protat, A., Warren, R. A., Collis, S. M., Wolff, D. B., Raunyar, S., ... Petersen, W. A. (2019). An Integrated Approach to Weather Radar Calibration and Monitoring Using Ground Clutter and Satellite Comparisons. *Journal of Atmospheric and Oceanic Technology*, 36(1), 17–39. doi: 10.1175/JTECH-D-18-0007.1
- Martin, G. M., Klingaman, N. P., & Moise, A. F. (2017). Connecting spatial and temporal scales of tropical precipitation in observations and the MetUM-GA6. *Geoscientific Model Development*, 10(1), 105–126. doi: 10.5194/gmd-10-105-2017
- Martin, G. M., Milton, S. F., Senior, C. A., Brooks, M. E., Ineson, S., Reichler, T., & Kim, J. (2010). Analysis and reduction of systematic errors through a seamless approach to modeling weather and climate. *Journal of Climate*, 23(22), 5933–5957. doi: 10.1175/2010JCLI3541.1
- Menon, A., Turner, A. G., Martin, G. M., & MacLachlan, C. (2018). Modelling the moistening of the free troposphere during the northwestward progression of Indian monsoon onset. *Quarterly Journal of the Royal Meteorological Society*, 144(713), 1152–1168. doi: 10.1002/qj.3281
- Parker, D. J., Willetts, P., Birch, C., Turner, A. G., Marsham, J. H., Taylor, C. M., ... Martin, G. M. (2016). The interaction of moist convection and mid-level dry air in the advance of the onset of the Indian monsoon. *Quarterly Journal of the Royal Meteorological Society*, 142(699), 2256–2272. doi: 10.1002/qj.2815
- Posselt, D. J., van den Heever, S. C., & Stephens, G. L. (2008). Trimodal cloudiness and tropical stable layers in simulations of radiative convective equilibrium. *Geophysical Research Letters*, 35(8), L08802. doi: 10.1029/2007GL033029
- Rajeevan, M., Bhate, J., Kale, J., & Lal, B. (2006). High resolution daily gridded rainfall data for the Indian region: analysis of break and active monsoon spells. *Current Science*, 91(3), 296–306.
- Rajeevan, M., Gadgil, S., & Bhate, J. (2010). Active and break spells of the Indian

- summer monsoon. *Journal of Earth System Science*, 119(3), 229–247. doi: 10.1007/s12040-010-0019-4
- Ramesh, K. J. (2017). *Annual Report 2016*. Retrieved 2020-07-03, from <https://metnet.ind.gov.in/indnews/ar2016.pdf>
- Rao, S. A., Goswami, B. N., Sahai, A. K., Rajagopal, E. N., Mukhopadhyay, P., Rajeevan, M., ... Maini, P. (2019). Monsoon mission a targeted activity to improve monsoon prediction across scales. *Bulletin of the American Meteorological Society*, 100(12), 2509–2532. doi: 10.1175/BAMS-D-17-0330.1
- Rao, T. N., Saikranthi, K., Radhakrishna, B., & Bhaskara Rao, S. V. (2016). Differences in the climatological characteristics of precipitation between active and break spells of the Indian summer monsoon. *Journal of Climate*, 29(21), 7797–7814. doi: 10.1175/JCLI-D-16-0028.1
- Ravi Kiran, V., Rajeevan, M., Vijaya Bhaskara Rao, S., & Prabhakara Rao, N. (2009). Analysis of variations of cloud and aerosol properties associated with active and break spells of Indian summer monsoon using MODIS data. *Geophysical Research Letters*, 36(9), L09706. doi: 10.1029/2008GL037135
- Romatschke, U., & Houze, R. A. (2011). Characteristics of Precipitating Convective Systems in the South Asian Monsoon. *Journal of Hydrometeorology*, 12(1), 3–26. doi: 10.1175/2010JHM1289.1
- Roy Bhowmik, S. K., Sen Roy, S., Srivastava, K., Mukhopadhyay, B., Thampi, S. B., Reddy, Y. K., ... Adhikary, S. (2011). Processing of Indian Doppler Weather Radar data for mesoscale applications. *Meteorology and Atmospheric Physics*, 111(3–4), 133–147. doi: 10.1007/s00703-010-0120-x
- Sathiyamoorthy, V., Pal, P., & Joshi, P. (2004). Influence of the upper-tropospheric wind shear upon cloud radiative forcing in the Asian monsoon region. *Journal of Climate*, 17(14), 2725–2735. doi: 10.1175/1520-0442(2004)017<2725:IOTUWS>2.0.CO;2
- Shige, S., & Kummerow, C. (2016). Precipitation-top heights of heavy orographic rainfall in the Asian monsoon region. *Journal of the Atmospheric Sciences*, 73(8), 3009–3024. doi: 10.1175/JAS-D-15-0271.1
- Sindhu, K. D., & Bhat, G. S. (2018). Characteristics of monsoonal precipitating cloud systems over the Indian subcontinent derived from weather radar data. *Quarterly Journal of the Royal Meteorological Society*. doi: 10.1002/qj.3328
- Sindhu, K. D., & Bhat, G. S. (2019). Storm characteristics and precipitation estimates of monsoonal clouds using C-band polarimetric radar over Northwest India. *Theoretical and Applied Climatology*, 1–12. doi: 10.1007/s00704-019-02828-6
- Steiner, M., Houze, R. A., Yuter, S. E., Steiner, M., Jr., R. A. H., & Yuter, S. E. (1995). Climatological Characterization of Three-Dimensional Storm Structure from Operational Radar and Rain Gauge Data. *Journal of Applied Meteorology*, 34(9), 1978–2007. doi: 10.1175/1520-0450(1995)034<1978:CCOTDS>2.0.CO;2
- Testud, J., Bouar, E. L., Obligis, E., & Ali-Mehenni, M. (2000). The Rain Profiling Algorithm Applied to Polarimetric Weather Radar. *Journal of Atmospheric and Oceanic Technology*. doi: 10.1175/1520-0426(2000)017<0332:TRPAAT>2.0.CO;2
- Turner, A. G., Bhat, G. S., Martin, G. M., Parker, D. J., Taylor, C. M., Mitra, A. K., ... New, B. (2019). Interaction of Convective Organisation with Monsoon Precipitation, Atmosphere, Surface and Sea: the 2016 INCOMPASS field campaign in India. *Quarterly Journal of the Royal Meteorological Society*, qj.3633. doi: 10.1002/qj.3633
- Turner, A. G., Sperber, K. R., Slingo, J., Meehl, G., Mechoso, C. R., Kimoto, M., & Giannini, A. (2011). Modelling Monsoons: Understanding and Predicting Current and Future Behaviour. In *The Global Monsoon System: Research and Forecast (2nd Edition)* (World Scientific Series on Asia-Pacific Weather and



- 820 *Climate*) (pp. 421–454). doi: 10.1142/9789814343411\_0025
- 821 Volonté, A., Turner, A., & Menon, A. (2019). Air mass analysis of the processes driv-  
 822 ing the progression of the Indian summer monsoon. *Quarterly Journal of the*  
 823 *Royal Meteorological Society*, qj.3700. doi: 10.1002/qj.3700
- 824 Warren, R. A., & Protat, A. (2019). Should interpolation of radar reflectivity be  
 825 performed in Z or dBZ? *Journal of Atmospheric and Oceanic Technology*,  
 826 JTECH-D-18-0183.1. doi: 10.1175/JTECH-D-18-0183.1
- 827 Warren, R. A., Protat, A., Siems, S. T., Ramsay, H. A., Louf, V., Manton, M. J.,  
 828 & Kane, T. A. (2018). Calibrating ground-based radars against TRMM and  
 829 GPM. *Journal of Atmospheric and Oceanic Technology*, 35(2), 323–346. doi:  
 830 10.1175/JTECH-D-17-0128.1
- 831 Yang, G.-Y., & Slingo, J. (2001). The Diurnal Cycle in the Tropics. *Monthly*  
 832 *Weather Review*, 129(4), 784–801. doi: 10.1175/1520-0493(2001)129<0784:  
 833 TDCITT>2.0.CO;2