

Board W-35: Parametric study of prompt methane release impacts on global mean temperature using GISS ModelE

PattiMichelle Sheaffer pattimichelle000@gmail.com
Research Scientist; Aerospace Corporation (retired)

There have been important criticisms of the Intergovernmental Panel on Climate Change (IPCC) recent reports for failing to communicate the *dire* nature of the current predicament facing civilization – so-called “scientific reticence” – as well as for *assuming* functional, planetary-effective scale biomass carbon capture and storage in its survivable scenarios [1-3]. In the light of major reports released in 2018 [4,5] which underscore the discrepancy between the current climate trajectory and best-case requirements to maintain global civilization, the current predicament is often described as an “existential” crisis [6]. Part of the confusion appears to stem from the lack of discussion of specific scenarios, such as rapid arctic methane release [7,8], which are not discussed by the IPCC in proportion to their catastrophic potential. This scenario is briefly examined using the Goddard Institute for Space Studies (GISS) ModelE v2 7.50.05 [9]. It is suggested that the results presented here represent a lower bound to climate disruption since in this set-up, neither the oceans nor arctic sea ice (a significant and ongoing runaway feedback [10]) respond to the changes modelled; namely, a sudden release of stored methane gas.

Goddard Institute ModelE

- Standard “Non-Interactive” (NINT) atmospheric model template with prescribed ocean (*Hadley 1975 – 1984 annual SST*) and sea ice (*Hadley 1996 – 2005 annual Sea Ice*) observational datasets. [11]
- Model spin-up 100 years 1920 – 2020 (transient simulation) starting from default global atmosphere observations
- *Likely these are conservative results as well-observed recent Arctic ice/snow changes are not modeled.*






Prompt Land Surface Temperature Response
Only land surface air temperature values were calculated. Temperatures were calculated by:

$$\sum_C (LF_C A_C P_C) / \sum_C (LF_C A_C)$$

...summing over all grid cells, LF_C = in-cell land fraction, A_C = cell area, P_C = in-cell model variable being averaged, for annual or seasonal averages of the monthly model outputs. CH_4 is assumed uniformly dispersed.

Mean grid cell temperatures were converted to *temperature change* by subtracting mean of 2020 – 2022 temperatures.

The temperature-change signal is noisy, but boxcar smoothing shows that it rises concomitantly with CH_4 loading, suggesting a unit impulse response for the system as modeled.






1. Anderson, K., *Nature*, 24/31 DECEMBER | VOL 528 | 437 [2015]; Watson, M., Bristol U., *Geoengineering*, Roy. Soc. London, November 2014
2. Anderson, K., and Peters, G., “The trouble with negative emissions,” *Science*, Vol. 354, Issue 6309, pp. 182-183 [2016]
3. Muratori, et al., “Global economic consequences of deploying bioenergy with carbon capture and storage (BECCS),” *Environ. Res. Lett.* 11 [2016]
4. Fourth National Climate Assessment, United States [2018]
5. IPCC Special Report SR-15, Figure SPM3.b [2018]
6. Spratt, Dunlop; Schellnhuber, “What Lies Beneath,” *Natnl. Ctr Climate Rest.*, Melbourne, Australia [2017]
7. Stolaroff, et al., “Review of Methane Mitigation Technologies with Application to Rapid Release of Methane from the Arctic,” *Environ. Sci. Technol.*, 46, 6455–6469 [2012]
8. Natalia Shakhova, et al., “Extensive Methane Venting to the Atmosphere from Sediments of the East Siberian Arctic Shelf,” *Science* 327, 1246 [2010]
9. Schmidt, G.A., et al., “Configuration and assessment of the GISS ModelE2 contributions to the CMIP5 archive,” *J. Adv. Model. Earth Syst.*, 6, no. 1, 141-184 [2014]
10. Guiles, et al., “Circumpolar thinning of Arctic sea ice following the 2007 record ice extent minimum,” *Geophys. Res. Lett.*, Vol 35, Issue 22 [2008]
11. National Center for Atmospheric Research Staff (Eds). “The Climate Data Guide: SST data: HadISST v1.1.” <https://climatedataguide.ucar.edu/climate-data/sst-data-hadisst-v1.1>.

Model Effects of 20% Decrease in Mean Arctic Insolation

(cf., Latham, et al. *Phil.Trans.R.Soc.A* 372 [2014])

Insolation Difference



- Unit impulse response (*fully-dispersed CH_4*)
- Temperature lags may be related primarily to CH_4 dispersion rates
- Larger warming over Arctic land surface (*oceans not investigated*)
- Decreased Arctic insolation measurably decreases mean Arctic land temperature