

Insights on Calving Processes from Fragmentation Theory Applied to Iceberg Size Distributions

Ellyn M Enderlin¹, Jukes Liu^{1*}, Michal Kopera¹, Caitlin Oliver¹, Timothy C Bartholomaus²

¹Boise State University Department of Geosciences, ²University of Idaho Department of Geological Sciences



BOISE STATE UNIVERSITY

C35A-0869



Iceberg Calving

Iceberg calving and associated terminus position change directly and indirectly control marine glacier and ice sheet mass.

The calving process is relatively poorly understood because it is incredibly difficult to collect observations of calving events.

- calving is a stochastic process & therefore difficult to predict
- the near-terminus region is a dangerous location to install & retrieve instruments

There is a broad spectrum of calving modes:

- low-energy very large tabular iceberg detachment
- high-energy full-thickness iceberg capsizing
- moderate-energy submarine or subaerial growler or bergy bit detachment

Fragmentation Theory

Iceberg calving requires either (a) the connection of branching fractures or (b) full-thickness penetration of isolated fractures.

Different fracture patterns produce mathematically-distinct iceberg size distributions:

- branching fractures = power-law distribution with a size cut-off
- isolated fractures = exponential distribution with a size cut-off

$$n(v) = c_1 v^{-\alpha} \exp(-v/c_2) + c_3 \exp(-v/c_4)$$

branching
isolated

Equation 1: Fragmentation curves used to describe iceberg size distributions. From Åström et al. (2021). v is iceberg size, $n(v)$ is iceberg count divided by iceberg size, and c_{1-4} are constants tuned to observations.




Figure 1: Idealized fragmentation curves. Modified from Åström et al. (2021).

Application of fragmentation theory to ice mélange size distributions can yield insights into controls on calving.





Figure 1: Iceberg size distribution study sites. Study sites are delineated by colored boxes in the large overview map produced in QGreenland. Subplots contain geographically-organized example digital elevation models (DEMs) for the study sites. The colored lines delineate the boundaries of the regions searched for ice mélange, with different colors marking the interior boundary (i.e., terminus) in each available DEM. The same elevation terminus colormaps are used in all subplots. Greenland Polar Stereo coordinates are used in all subplots but the scales for the subplots differ depending on extent of each region.



Figures 3 & 4: Observed iceberg size distributions. (a) Digital elevation models, (b) pixel elevation distributions, and (c) iceberg size distributions for Alison Glacier on May 7 and July 27, 2020, respectively.




Figure 5: Iceberg size distributions for Alison Glacier for 2011-2020. Colors indicate months. The same axes scales are used for all subplots.

Methods Example: Alison Glacier (Figs. 3-5)

1. Extract iceberg surface elevations from very high-resolution digital elevations of near-terminus ice mélange.
2. Convert ice mélange elevations to iceberg size distributions using width-thickness (i.e., aspect) ratios ranging from 1.7-2.1. See Enderlin et al. (2016) for more details.
3. Fit fragmentation theory curves to normalized iceberg size distributions:
 - automated inversion
 - manual adjustment as necessary

Results

- Uncertainty in iceberg aspect ratios has minimal influence on iceberg size distributions (Figs. 3&4: Blue shading in panel c).
- Power-law can over-predict small iceberg abundance due to rapid submarine melting of growlers & bergy bits.
- No clear seasonal patterns in the shape of the distributions (Fig. 6).




Figure 6 (left): Seasonal fragmentation theory-modeled iceberg size distributions from all cloud-free DEMs available from 2011 through winter 2019. Colors correspond to the different regions. Warmer colors generally correspond with warmer subsurface water temperatures.

Figure 7 (right): The iceberg size that is equally driven by branching and isolated fractures. Larger sizes are representative of tabular icebergs that form as the result of isolated fractures.

- Some seasonality apparent in the iceberg size above which isolated fracture dominates and tabular icebergs form (Fig.7).

Acknowledgements & References

This research is funded by NSF 1933105 "Collaborative Research: What controls calving? A Greenland-wide test of terminus change mechanisms" awarded to Boise State University. DEMs provided by the Polar Geospatial Center under NSF-OPP awards 1043681, 1559691, and 1542736.

- Åström J, Cook S, Enderlin EM, Sutherland DA, Mazur A, Glasser N (2021). Fragmentation theory reveals processes controlling iceberg size distributions. *Journal of Glaciology* 1–10. <https://doi.org/10.1017/jog.2021.14>.
- Enderlin, E. M., G. S. Hamilton, F. Straneo, and D. A. Sutherland (2016), Iceberg meltwater fluxes dominate the fresh-water budget in Greenland's iceberg-congested glacial fjords, *Geophys. Res. Lett.*, 43, 11,287–11,294, doi:10.1002/2016GL070718.