

Zooplankton fecal pellets as the primary driver of settling particles in reef systems

Jaclyn Torkelson (jftorkel@syr.edu) and Mark A. Teece
Department of Chemistry, State University of New York College of Environmental Science and Forestry

Background

Settling particles and their associated organic matter are a potentially vital source of nutrients for corals. While settling particles can block out light and reduce the ability of zooxanthellae (the symbiotic algae that live inside coral tissues) to photosynthesize, this primarily occurs in highly turbid environments (Jones et al. 2019; Nugroho et al. 2018; Risk 2014). In areas with low turbidity, settling particles can be ingested by several species of Scleractinia (reef building coral) (e.g. Anthony, 1999; Krueger et al., 2018). Corals must acquire essential fatty acids (EFAs) from their zooxanthellae (Papina et al., 2003) or through their diet (Ferrier-Pagès & Gattuso, 1998). Through both their daily vertical migration and fecal pellets, zooplankton help to deliver organic matter from the upper water column to coral reefs (Heidelberg et al., 2010). The presence of both phytoplankton and zooplankton are vital to the organic matter input to coral reefs as the average daily mass of phytoplankton alone in the Florida Keys can reach 21.4 mg C m⁻³ (Heidelberg et al., 2010). Since Scleractinia have been shown to ingest settling particles, it could be an important source of nutrients, such as essential lipids, fatty acids, and nitrogen, to supplement their diet.

The primary objective of our study was to evaluate the role of zooplankton as a conduit of organic matter to benthic corals through settling particles. We measured the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of values of settling particles collected from reefs of the Florida Reef Tract. Additionally, fatty acid and lipid biomarkers were measured to determine the origin of organic matter. Through their diurnal vertical migration, zooplankton and their production of fecal pellets are vital to the transferal of organic matter from the water column to the benthos and coral reefs. Our study provides crucial information on the origin of nutrients on coral reefs.

Methods




Figure 1. Location of the Florida Keys within the US



Figure 2. Sampling Locations in the Upper Florida Keys



Figure 3. Sediment trap at Offshore South



Results




Figure 4. Composition of fatty acids in settling particles (mean of n=5 bars represent standard error)




Figure 5. Composition of sterols in settling particles (mean of n=5, bars represent standard error)





Figure 6. Total sedimenting flux of settling particles (mean of n=5, bars represent standard error)

- Saturated fatty acids (SAFA; 52-58%) were the dominant compounds in all samples with lower amounts of monounsaturated (MUFA; 25-29%) and polyunsaturated (PUFA; 11-20%) fatty acids.
- $\delta^{13}\text{C}$ values of individual fatty acids in settling particles spanned a considerable range from -18.8 to -27‰.
- Cholesterol accounted for up to 48% of all sterols.
- All stanols were present in low concentrations (< 3%).
- $\delta^{13}\text{C}$ values of all sterols and stanols were lower at the most offshore sites (Mid Reef and Offshore North) compared with the nearshore sites.
- Organic matter and organic nitrogen flux mirrored total flux.
- No significant difference in organic matter and inorganic carbon between sites.

Discussion



- Zooplankton are the primary component of settling particles
- Compound specific isotope analysis indicates zooplankton as $\delta^{13}\text{C}$ values were similar to zooplankton, not phytoplankton.
- Little to no presence of bacteria as there were small concentrations of odd-chain fatty acids and stanols.
- Sterols representing diatoms were constituted 13-23% of total sterols.
- Biomarkers related to vascular plants and macroalgae were more common near shore and constituted up to 18% of the total lipids.
- Gorgosterol, the primary indicator of coral and zooxanthellae constituted 0.2-2.2% of the total sterols, demonstrating little contribution to settling particles.
- Of the phytoplankton, known biomarkers of diatoms and dinoflagellates were the most common.
- As the climate continues to change organic matter will be remineralized at shallower depths, reducing the quantity of fecal pellets reaching the benthos.

References

- Anthony, K. R. N. (1999). Coral suspension feeding on fine particulate matter. *Journal of Experimental Marine Biology and Ecology*, 232(1), 85–106. [https://doi.org/10.1016/S0022-0981\(98\)00099-9](https://doi.org/10.1016/S0022-0981(98)00099-9)
- Ferrier-Pagès, C., & Gattuso, J. P. (1998). Biomass, Production and Grazing Rates of Pico- and Nanoplankton in Coral Reef Waters (Miyako Island, Japan). *Microbial Ecology*, 35(1), 46–57. <https://doi.org/10.1007/s002489900059>
- Heidelberg, K. B., O'Neil, K. L., Bythell, J. C., & Sebens, K. P. (2010). Vertical distribution and diel patterns of zooplankton abundance and biomass at Conch Reef, Florida Keys (USA). *Journal of Plankton Research*, 32(1), 75–91. <https://doi.org/10.1093/plankt/fbp101>
- Krueger, T., Bodin, J., Horwitz, N., Lousset-Fonta, C., Sake, A., Escrig, S., Fine, M., & Meibom, A. (2018). Temperature and feeding induce tissue level changes in autotrophic and heterotrophic nutrient allocation in the coral symbiosis – A NanoSIMS study. *Scientific Reports*, 8(1), 12710. <https://doi.org/10.1038/s41598-018-31094-1>
- Papina, M., Meziane, T., & van Woerk, R. (2003). Symbiotic zooxanthellae provide the host-coral *Montipora digitata* with polyunsaturated fatty acids. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 145(3), 333–337. [https://doi.org/10.1016/S1366-4995\(03\)00118-1](https://doi.org/10.1016/S1366-4995(03)00118-1)
- Storlazzi, C. D., Elias, E., Field, M. E., & Presto, M. K. (2011). Numerical modeling of the impact of sea-level rise on fringing coral reef hydrodynamics and sediment transport. *Coral Reefs*, 30(5), 83–96. <https://doi.org/10.1007/s00338-011-0723-9>
- Teece, M. A., Estes, B., Gelseichter, E., & Lirman, D. (2011). Heterotrophic and autotrophic assimilation of fatty acids by two scleractinian corals, *Montastraea faveolata* and *Porites astreoides*. *Limnology and Oceanography*, 56(4), 1285–1296. <https://doi.org/10.4319/lno.2011.56.4.1285>

Acknowledgements

Thank you to Jesse Crandall, Diego Lierman, and Jess Ciesla for their work and analysis.

This material is based on work supported by Mote Marine Laboratory Protect Our Reefs Program, and in part by a grant from the National Undersea Research Program of the National Oceanic and Atmospheric Administration (NOAA) and the NOAA Coral Reef Conservation Program.