

# Subseasonal Great Plains Rainfall via Remote Extratropical Teleconnections: Regional Application of Theory-guided Causal Networks

Kelsey Malloy<sup>1</sup>, Ben P. Kirtman<sup>2</sup>

<sup>1</sup>Department of Applied Physics and Applied Mathematics, Columbia University, New York, New York  
<sup>2</sup>Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Florida

## Key Points:

- Subseasonal monsoon variability via the boreal summer intraseasonal oscillation is linked to rainfall signals over U.S. Great Plains and its associated dynamical drivers.
- An algorithm that specializes in identifying cause-and-effect relationships verified a pathway from regional monsoon rainfall to Great Plains rainfall, which takes approximately 2 weeks.
- Weekly rainfall over the East Asian monsoon region is causally linked to the triggering of a Rossby wave pattern, Great Plains low-level jet strengthening, and active Great Plains convection about one week later.

## Abstract

Long-range U.S. summer rainfall prediction skill is low. Monsoon variability, especially over the West North Pacific Monsoon (WNPM) and/or East Asian Monsoon (EAM) region, can influence U.S. Great Plains hydroclimate variability via a forced Rossby wave response. Here we explored subseasonal monsoon variability as a source of predictability for Great Plains rainfall. The boreal summer intraseasonal oscillation is related to Great Plains convection and Great Plains low-level jet (LLJ) anomalies as well as a cross-Pacific wave train. Using a causal effect network, we found that the time between BSISO-related geopotential height anomalies and Great Plains rainfall anomalies is about 2 weeks; therefore, BSISO convection may be a valuable forecast of opportunity for subseasonal prediction of Great Plains convection anomalies. More specifically, causal link patterns/maps revealed that the above-normal weekly EAM rainfall, rather than WNPM rainfall or general geopotential height activity over the East Asia, was causally linked to Great Plains LLJ strengthening and active Great Plains convection the following week.

## 1 Introduction

Understanding summertime continental U.S. (CONUS) hydroclimate predictability on the subseasonal-to-seasonal (S2S) timescale has been challenging, and relationships between tropical remote forcing and mid-latitude circulation are difficult to assess due to the overall weak signals of the summer season (Trenberth et al., 1998; S. Zhou et al., 2012). Many studies suggest that Asian summer monsoon (ASM) variability on the seasonal-to-interannual timescale, especially over the West North Pacific (WNPM) and/or East Asian Monsoon (EAM) region, can influence CONUS hydroclimate via a quasi-stationary Rossby wave response (Di Capua, Runge, et al., 2020; Lopez et al., 2019; Kornhuber et al., 2019; Malloy & Kirtman, 2022a, 2022b, manuscript submitted; Yang et al., 2020; Zhu & Li, 2016, 2018). The Great Plains low-level jet (LLJ) is the prominent transporter of moisture into that region, and large-scale LLJ anomalies are typically associated with rainfall events (Arritt et al., 1997; Higgins et al., 1997; Cook et al., 2008; Weaver & Nigam, 2008; Weaver et al., 2009; Nayak & Villarini, 2017; Algarra et al., 2019; Malloy & Kirtman, 2020). The upper-level pattern associated with the monsoon-forced Rossby wave response can often align (constructively interfere) with the Great Plains LLJ to amplify Great Plains rainfall signals (Malloy & Kirtman, 2022b, manuscript submitted).

The ASM also exhibits subseasonal variability, typically called the boreal summer intraseasonal oscillation (BSISO), and it is the dominating mode of tropical convection over ASM region and western Pacific (Yasunari, 1979, 1980; S. S. Lee & Wang, 2016). Moon et al. (2013) and Krishnamurthy et al. (2021) identified the monsoon intraseasonal oscillation as a source of subseasonal predictability over CONUS in the summer in observations and/or climate forecast models. Few studies have explored the dynamical pathway between BSISO-related anomalies and Great Plains rainfall anomalies, such as understanding the timescale of Rossby wave initiation and propagation to influence North American features, such as the Great Plains LLJ.

In many of these studies, climate models were used to quantify the monsoon responses, usually by prescribed heating, and were compared to observations (Lopez et al., 2019; Malloy & Kirtman, 2022a, 2022b, manuscript submitted; Yang et al., 2020). In this case, causality is implied (amongst natural variability or chaos). For example, the EAM heating causes the elongated anomalous ridge over the North Pacific, anomalous trough over western North America, and anomalous ridge over eastern North America from the set of experiments in Malloy and Kirtman (2022a). However, there are also ways to quantify causal links via data-driven methods i.e. using observations alone. Causal discovery methods, such as causal effect networks (CENs), are becoming popular as a way to map physical links in the climate system within an inputted time series of data (Runge

et al., 2014; Runge, 2018; Runge et al., 2019; Runge, 2020; Kretschmer et al., 2016). Using CENs, Di Capua, Kretschmer, et al. (2020) found that there was a link between the North Atlantic Oscillation (NAO), circumglobal teleconnection (CGT), and ASM variability, as well as between the BSISO and ASM variability. Di Capua, Runge, et al. (2020) suggested that the WNPM may force the North Pacific circulation that subsequently influences temperature and rainfall anomalies over North America.

CENs effectively determine causal links while removing the effects from autocorrelation, indirect (spurious) links, or common drivers, which maintaining a high detection power over other techniques, such as Granger causality model (Runge, 2018; Runge et al., 2019; Runge, 2020). There are many assumptions in using CENs, including that causality can only be determined among the given drivers. Adding or removing drivers can change the conditional (in)dependence and hence change the linkages. Therefore, knowledge of the physical system beforehand, including relevant variables and timescales, is essential for interpreting the output of the algorithm.

The objective of this study is to apply CENs to identify the remote drivers of the Great Plains LLJ and rainfall anomalies on the subseasonal timescale. This extends upon the methodology from Di Capua, Kretschmer, et al. (2020) and Di Capua, Runge, et al. (2020) by applying it to understand more regional-scale mechanisms. We also successfully isolate the impacts from interrelated drivers in the CEN, shedding light on the source of U.S. Great Plains hydroclimate predictability.

## 2 Data and Methods

### 2.1 Observational Datasets

This study focuses on the extended summer season (May through September), though April was considered for the lead-lag correlation analysis and CEN. Pressure-level meridional wind, zonal wind, geopotential height were taken from the European Centre for Medium-Range Weather Forecasts (ECMWF) fifth-generation reanalysis (ERA5). ERA5 atmospheric data is provided hourly on a  $0.25^\circ$  latitude/longitude grid (Hersbach et al., 2020), and it is recalculated to daily averages. U.S. precipitation data were taken from the CPC Global Unified Gauge-based Analysis, provided on a  $0.5^\circ$  latitude/longitude grid over land (Chen et al., 2008; Xie et al., 2007). Outgoing Longwave Radiation (OLR) data, used as a proxy for convection, were taken from the interpolated daily OLR version 1.2 from National Oceanic and Atmospheric Administration (NOAA) Climate Data Record (CDR), accessed from <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00875>.

Because this study is focused on intraseasonal variability, for every variable, we removed the centered 120-day moving mean at every grid point in addition to removing the annual cycle. Then we took the centered 10-day running mean of the variables to focus on large-scale, low-frequency features. For the CEN analysis, we used the filtered data and resampled the data as weekly averages; using weekly data is a practical approach for subseasonal predictability analysis in order to filter short-term temporal fluctuations (Di Capua, Kretschmer, et al., 2020; Krishnamurthy et al., 2021).

### 2.2 Potential Drivers

The Great Plains precipitation index is defined by averaged precipitation anomalies within the  $35\text{--}50^\circ\text{N}$ ,  $85\text{--}105^\circ\text{W}$  domain, and the Great Plains LLJ index is defined by the averaged V850 anomalies within the  $25\text{--}35^\circ\text{N}$ ,  $90\text{--}102^\circ\text{W}$  domain. These domains are slightly larger than in previous literature (Weaver & Nigam, 2008; Malloy & Kirtman, 2020) to account for shifts eastward, which may be more important for EAM-forced Great Plains LLJ variability that is coupled to the upper levels (Malloy & Kirtman, 2022b,

**Table 1.** Potential Drivers to Great Plains Rainfall<sup>a</sup>

Name	Identifier	Index Calculation
<i>Great Plains precipitation</i>	<i>GP rainfall</i>	<i>Precipitation*[35-50°N, 85-105°W]</i>
Great Plains Low-level Jet	GPLLJ	V850*[25-35°N, 90-102°W]
Pacific-North America High-Low dipole	PNA-HL	(Z200*[35-60°N, 135-165°W] - Z200*[35-60°N, 100-130°W])
North Pacific Low	NPac-L	Z200*[35-60°N, 160°E-170°W]
East Asian Monsoon Low	EAM-L	Z200*[25-50°N, 90-130°E]
East Asian Monsoon precipitation	EAM rainfall	Precipitation*[20-30°N, 100-125°E]
West North Pacific Monsoon precipitation	WNPM rainfall	Precipitation*[0-20°N, 90-120°E]

<sup>a</sup>The Great Plains precipitation index (italicized) included here as predictand. Index is calculated by taking the domain-averaged anomalies of the variable.

manuscript submitted). In addition, we defined various other indices to input into the CEN as potential drivers based on the lagged correlation analysis. All inputs, or potential drivers, to the Great Plains LLJ and Great Plains rainfall are outlined in Table 1 and can be visualized in Figure 3 boxes.

### 2.3 Causal Effect Networks

The CEN is constructed by first applying the Peter and Clark Momentary Conditional Independence (PC-MCI) algorithm (Spirtes et al., 2000; Runge et al., 2014, 2019). This is a two-step procedure: (1) the PC step finds the relevant drivers, or “parents”, of each variable via an iterative independence testing, and (2) the MCI step removes spurious or common parents by conditioning the partial correlations between parents and variables on the parents of the parents.

Start with a set  $X$  of  $n$  variables that are timeseries of anomalies. The PC algorithm first calculates the correlation between the  $i$ th variable in  $X$  and the rest of the variables in  $X$  at time lag  $\tau$ . The significant correlations with the  $i$ th variable form a set of potential parents  $X_i^0$  at time lag  $\tau$ . Then, it calculates the partial correlation between the  $i$ th variable and each potential parent in  $X_i^0$ , but with a condition that the first variable in  $X_i^0$  that has the strongest correlation with the  $i$ th variable. If  $a$ ,  $b$ , and  $c$  are variables in  $X$ , the partial correlation between  $a$  and  $b$  conditioned on  $c$  is calculated by performing a linear regression of  $a$  on  $b$  and  $b$  on  $c$ , then correlating the residuals. Variables  $a$  and  $b$  are conditionally dependent given  $c$ , i.e. their correlation cannot be explained by the influence of  $c$  (not spurious link) if the resulting partial correlation is significant at threshold  $\alpha$ . This may reduce the set of parents for the next iteration  $X_i^1$ . The process is repeated for this set of parents but with now two conditions, leading to a next (possibly reduced) set of parents  $X_i^2$ . When the number of parents is equal to or greater than the number of conditions needed to calculate partial correlation, the algorithm converges.

The MCI step calculates the partial correlation between each variable and its parents at different time lags conditioned on both the set of parents and the parents of the parents, essentially removing common driver effects and reducing to a final set of causal parents.

The CEN calculates these causal relationships by performing a standardized multiple regression of each variable with its parents. The final link is represented as the change in standard deviation ( $\sigma$ ) of variable at time  $t$  if the parent was raised to  $1\sigma$  at time  $t - \tau$ . More detail of this algorithm and its comparison to other causality methods can be found in Runge et al. (2019); the PC-MCI algorithm is freely available at <https://github.com/jakobrunge/tigramite>.

There are many assumptions to using the CEN, including that causal links are determined *relative to the chosen set of variables*. Removing or adding variables may change the CEN, and therefore, it is important for the user to understand the physical system. Other assumptions include stationarity of relationships and near-linear interactions.

In this study, the CEN visualizes the causal links with a time lag of one week. Contemporaneous links are also visualized with no causality direction inferred. The winter season is masked, which means that timescales of variables are restricted to MJJAS season, but the parent (and conditional) timeseries may include April. We set  $\alpha = 0.05$ , which is the significance threshold as explained above, and  $\tau_{max} = 3$  weeks, which is maximum time delay, though we find that the results are not sensitive to the choice of  $\tau_{max}$  between 2 and 5.

## 2.4 Causal Maps

Finally, we experiment with causal maps, which plots the link coefficient from the CEN spatially (Di Capua, Runge, et al., 2020). Two one-dimensional timeseries are chosen that have a theoretical relationship with a three-dimensional field. The CEN determines the causal link between one of the one-dimensional timeseries and a timeseries of a gridpoint from the three-dimensional field, conditioned on the other one-dimensional timeseries. To distinguish between the WNPM- and EAM-forced patterns, as well as the EAM- and EAM-L-forced patterns, we use the weekly WNPM, EAM, and EAM-L timeseries, and the three-dimensional fields of interest are weekly Z200, V850, and OLR. Time lags of 1 and 2 weeks are explored, but, because the 2-week lagged patterns lack statistical significance over CONUS, only the 1-week lagged patterns are presented here.

## 3 Results

### 3.1 Link Between BSISO and Great Plains Rainfall

Before constructing the CEN, we first establish the potential influence of subseasonal monsoon variability on CONUS hydroclimate. Composites of OLR and U200 anomalies for the combined BSISO phases (phases 8+1, 2+3, 4+5, and 6+7) are depicted in Figure 1, highlighting the northeastward propagation of the regions of active and inactive convection as well as its influence on jet stream anomalies over the North Pacific. In particular, phases 2+3 are associated with active convection (negative OLR) over the equatorial Indian Ocean and weak wet anomalies over the EAM region, which corresponds with strengthening or northward displacement of the jet stream over East Asia at  $55^\circ\text{N}$  (Figure 1b). During phases 4+5, active convection over East Asia strengthens slightly and the positive U200 anomalies are extended over the North Pacific (Figure 1c). Composites of phases 6+7 and 8+1 present opposite patterns to 2+3 and 4+5, respectively.

The location of above-normal convection in the ASM region, indicated by BSISO phase, is related to the anomalous probability of Great Plains rainfall events, Great Plains LLJ events, and height patterns over the northeastern Pacific-western North America regions (Figure 2). For example, there is an increased (decreased) probability of a below-normal (above-normal) rainfall event  $\sim 3$  weeks after BSISO phase 3 (Figure 2a,b). The anomalous probabilities for the rainfall events coincide with the expected anomalous probabilities for the Great Plains LLJ and PNA events (Figure 2b-f). For example, days with

increased probability for the below-normal rainfall event are generally days with an increased probability for strong Great Plains LLJ event and PNA+ pattern. This is also true for the above-normal Great Plains rainfall events. Overall, there is an inferred propagation of the signal from the BSISO on these timescales, as seen by the diagonal stripes of increased or decreased anomalous probability.

This is further analyzed by investigating the lagged spatial correlation between Great Plains rainfall at  $T = 0$  and the OLR, V850, and Z200 anomaly fields at 0, 10, and 20 days before. The correlation between the Great Plains precipitation index and OLR anomalies at  $T = 0$  demonstrates the active convection, and hence the precipitation, over the northern Plains (Figure 3a). This corresponds with the strong anomalous southerly flow over the region (Figure 3d) and anomalous low pressure over western North America (Figure 3g). Over the monsoon region and North Pacific, there is a negative correlation with OLR anomalies at  $30^\circ\text{N}$  between  $90^\circ\text{E}$  and  $150^\circ\text{E}$  (Figure 3a, magenta domain) in addition to a positive correlation with EAM-related southerly flow (Figure 3e, black contour outline). A wave train is correlated with the Great Plains precipitation, including our PNA-HL pattern and NPac-L feature (Figure 3g, orange boxes). The PNA-HL pattern has been identified before as an important precursor for Plains rainfall events (Rogers & Coleman, 2003; Harding & Snyder, 2015; Patricola et al., 2015; Mallakpour & Villarini, 2016; Nayak & Villarini, 2017; Malloy & Kirtman, 2020).

The correlation between the Great Plains precipitation index and these field anomalies at  $T = -10$  (10 days before) reveals that some of the variability of Great Plains rainfall can be due to this cross-Pacific wave train that can be forced/modulated by EAM rainfall (Figure 3b,e,h). The correlation with negative OLR and positive V850 anomalies over the EAM region is -0.1 to -0.2, and the wave train pattern is present, including a  $\sim 0.2$  correlation with the EAM-L and NPac-L features (Figure 3h, left and right orange domains, respectively). There is also a correlation with positive OLR over the WNPM region 10 days before Great Plains precipitation events, showing an OLR pattern similar to that of combined phases 8+1 of the BSISO (cf. Figure 1a). In general, the correlations at  $T = -20$  are somewhat opposite to  $T = 0$  and  $T = -10$ , respectively.

These results suggest that the BSISO influences Great Plains rainfall on subseasonal timescales via a cross-Pacific Rossby wave train. We are motivated to test these linkages with a causal discovery algorithm and to confirm if the pathway from EAM rainfall to Great Plains rainfall exists and is considered causal. This method will also approximate the timeframe on which the BSISO-related rainfall anomalies lead to Great Plains anomalies (e.g. within  $\sim 2$  weeks, cf. Figure 3).

### 3.2 Causal Effect Network for Great Plains Rainfall

Because of the relatively large number of potential drivers, we simplify the discussion of the causal network by separating it into three spatial domains: over East Asia and western North Pacific, over the mid-latitude North Pacific, and over central-eastern North Pacific and North America. First, we test for a causal pathway between the WNPM rainfall, EAM rainfall, EAM-L feature, and NPac-L feature (Figure 4). There is a contemporaneous negative link between weekly-averaged WNPM rainfall and EAM rainfall, and a positive contemporaneous link between weekly-averaged EAM rainfall and the EAM-L feature. A strengthening of the weekly-averaged EAM-L by  $1\sigma$  leads to a  $0.38\sigma$  strengthening in the NPac-L feature the following week. The EAM-L and NPac-L features also have a contemporaneous link. This CEN indicates that the excitation of the Rossby wave activity over the North Pacific often depends on the presence of the EAM-L feature.

Next we consider the pathway between the different geopotential height features over the North Pacific (Figure 5). A strengthening of the NPac-L by  $1\sigma$  leads to a  $0.1\sigma$  strengthening in the PNA-HL pattern the following week, suggesting that the full cross-



Pacific Rossby wave train pathway may take up to 2 weeks. The contemporaneous links between these features demonstrate persistence which makes these relationships complex. For example, the contemporaneous negative link between the EAM-L and PNA-HL is also found in Figure 3i: by the time the Rossby wave reaches North America, the geopotential heights over EAM region are reversed.

Finally, the pathway between NPac-L and the Great Plains LLJ and Great Plains rainfall are visualized in the CEN (Figure 6). A strengthening of the PNA-HL by  $1\sigma$  leads to a  $0.16\sigma$  increase in the Great Plains rainfall and  $0.09\sigma$  strengthening of the Great Plains LLJ the following week. However, a strengthening of the NPac-L by  $1\sigma$  leads to a  $0.2\sigma$  increase in the Great Plains rainfall and  $0.14\sigma$  strengthening of the Great Plains LLJ the following week, which are greater causal links. Considering the contemporaneous link between PNA-HL and Great Plains rainfall and the PNA-HL and Great Plains LLJ are strongly positive, it is likely that the PNA-HL influence on these indices operates on sub-weekly timescales. However, this demonstrates that the NPac-L may assist in longer-lead prediction of the Great Plains LLJ and rainfall.

The CEN captured the intricacies of the relationships between these indices and their influence on the Great Plains LLJ and Great Plains rainfall. In addition, it demonstrated that the rainfall over the BSISO region is linked to a EAM-L feature that can generate a wave train response over the North Pacific that influences rainfall anomalies over the Great Plains. This can occur on a  $\sim 2$ -week timescale, potentially advantageous for understanding prediction on the subseasonal timescale.

Because of the contemporaneous relationship between the WNPM rainfall, EAM rainfall, and the EAM-L, it is difficult to assess the true causality between these features and downstream impacts. This motivates the use of causal maps to separate the patterns between WNPM and EAM rainfall as well as between EAM rainfall and EAM-L.

### 3.3 Causal Maps

By comparing the causal maps for Z200, V850, and OLR (Figures 7-9) with the lag correlation patterns from Figure 3, it is evident that EAM rainfall variability contributes to Great Plains rainfall variability on weekly timescales. First, we consider the causal linkages between the WNPM rainfall, EAM rainfall, and EAM-L feature at  $T = -1$  (one week before) and the Z200 field at  $T = 0$ . After removing the signal from EAM rainfall, WNPM rainfall affects the Z200 field mostly in the tropics and subtropics, with a  $1\sigma$  increase in WNPM rainfall causally linked to  $\sim 0.2\sigma$  anomalous ridging across the central-eastern tropical Pacific (Figure 7a). After removing the signal from the WNPM, EAM rainfall affects the mid-latitude Z200 field, with a  $1\sigma$  increase in EAM rainfall causally linked to a  $\sim 0.15$ - $0.2\sigma$  anomalous troughing over the WNPM region and North Pacific as well as  $\sim 0.15\sigma$  anomalous ridging at higher latitudes in East Asia (Figure 7b). This pattern is similar to the EAM rainfall causal map with the EAM-L signal removed (Figure 7c), but the removal of the EAM-L feature reduces the magnitude of the links. Finally, after removing the signal from EAM rainfall, the EAM-L impact on the Z200 field is prominent, with a  $1\sigma$  strengthening of the EAM-L causally linked to  $\sim 0.2$  anomalous ridging over high-latitude East Asia and  $0.3$ - $0.4\sigma$  anomalous troughing over the mid-latitude Pacific - including the NPac-L feature - and the Pacific Northwest (Figure 7d). The map of causal links between the EAM rainfall and the Z200 field help explain the Z200 patterns at  $T = 0$ , and perhaps  $T = -10$ , from the lag correlations (cf. Fig 3g,h) over the EAM region and North Pacific. The wave train over North America is not explained causally by the WNPM, EAM or EAM-L, suggesting that maybe this wave train is forced locally, perhaps by feedbacks from Great Plains rainfall. Nevertheless, EAM rainfall can generate the upstream Rossby wave activity that affects this region.

Next, we consider causal linkages with the V850 field. WNPM impacts to V850 are most evident in the WNPM and EAM regions as well as the central tropical Pacific (Fig-

ure 8a). After removing either the WNPM or EAM-L signal, the causal linkages between EAM rainfall and V850 field are relevant to North America, with a  $1\sigma$  increase in EAM rainfall leading to a  $\sim 0.15\sigma$  strengthening of the Great Plains LLJ (Figure 8b,c). EAM rainfall also impacts flow over the high-latitude Bering Sea/Alaska region. The causal linkages between the EAM-L and V850 field are prevalent over the EAM region and North Pacific. In particular, a  $1\sigma$  strengthening in the EAM-L is causally linked to a  $0.1\text{--}0.2\sigma$  strengthening of the low-level EAM flow. Once again, the EAM rainfall causal links explain more of the lag correlation patterns over North America (cf. Figure 3d,e), though the EAM-L is likely playing a role in amplifying the EAM or its signals.

The causal maps for the OLR field further demonstrate the influence of EAM rainfall. While WNPM rainfall impacts to OLR are mostly constrained to the subtropics and tropics (Figure 9a), the EAM rainfall links to OLR are most evident over the North Pacific and North America (Figure 9b). A  $1\sigma$  increase in EAM rainfall is causally linked to a  $0.1\text{--}0.2\sigma$  decrease in OLR (active convection) over the Great Plains. Patterns and link magnitudes are similar for the EAM rainfall impacts with the EAM-L signal removed (Figure 9c). Interestingly, the EAM-L is causally linked to OLR over the EAM region and Pacific Northwest (Figure 9d). A  $1\sigma$  strengthening of the EAM-L may lead to a  $0.1\text{--}0.2\sigma$  increase in EAM rainfall in addition to  $0.1\text{--}0.2\sigma$  increase in Pacific Northwest rainfall. These OLR patterns agree with the Z200 patterns from the EAM-L forcing (cf. Fig 7d), i.e. active convection is expected in these regions with that upper-level geopotential height pattern. The map of causal links from the EAM rainfall is helpful to explain the OLR patterns over the Great Plains region from the  $T = 0$  lag correlations (cf. Fig 3a), whereas the map of causal links from the EAM-L feature is helpful for describing the OLR patterns over North Pacific and Pacific Northwest at  $T = -10$  days (cf. Fig 3b).

In brief, EAM rainfall and EAM-L feature contribute to Z200 and OLR patterns over the mid-latitude Pacific and/or North America with a one week lag. However, EAM rainfall is more directly linked to Great Plains rainfall variability on this timescale, while EAM-L may modulate or amplify EAM-forced activity (or vice versa). Strong upper-level circulation anomalies over North America from Figure 3i were not explained by the monsoons nor the EAM-L feature, suggesting that localized feedbacks by the Great Plains rainfall itself might be forcing or amplifying that pattern.

### 3.4 Rossby Wave Source Anomalies

To further interpret the causal maps and contextualize these results with respect to potential dynamical mechanisms, we consider the composited 200-hPa Rossby wave source (RWS) anomalies during upper tercile WNPM or EAM days, which we calculated using the filtered daily data. The RWS term describes vorticity advection by the divergent wind and vortex stretching by the divergent wind (Sardeshmukh & Hoskins, 1988). RWS anomalies provide information about the production and origin of teleconnection wave patterns, which has been useful for explaining summertime circulation variability (Fuentes-Franco et al., 2022; Lopez et al., 2019; O'Reilly et al., 2018). We hypothesize that the magnitude and/or location of the diabatic heating from EAM generates a RWS with greater magnitude than from the WNPM heating, explaining the greater mid-latitude response and resulting teleconnection from EAM rainfall (cf. Figs. 7-9).

Strong WNPM days are associated with weak Rossby wave forcing over the West North Pacific region (Figure 10a). In contrast, for EAM days, there is a relatively strong negative 200-hPa RWS anomaly (shaded) collocated with EAM-related divergence (gray contours; Figure 10b). U200 anomalies suggest jet stream perturbations due to the divergence being close to the East Asian jet. The Z200 response from the lagged correlation analysis (cf. Fig. 3g) and causal maps (cf. Fig. 7b,c) can be explained by this teleconnection excitation from EAM-related divergence, likely due to release of diabatic heating.



## 4 Summary and Discussion

Here we explored the subseasonal predictability of Great Plains rainfall with a theory-guided application of CENs. Using a traditional lead-lag analysis approach, we found that the BSISO is related to Great Plains rainfall, the Great Plains LLJ, and PNA-HL pattern via a cross-Pacific wave train. The time between EAM-L anomalies potentially influencing Great Plains rainfall anomalies is  $\sim 2$  weeks; therefore, BSISO forcing or modulation of the EAM-L may be valuable forecast of opportunity for subseasonal prediction of Great Plains rainfall. Causal link patterns and associated RWS anomalies from the EAM rainfall revealed that the EAM is causally linked to excitation of Rossby wave patterns, leading to downstream Great Plains LLJ and rainfall anomalies. Anomalous geopotential height activity over EAM region (e.g. EAM-L pattern) may have a role in modulating the EAM-related patterns.

We applied similar techniques to Di Capua, Kretschmer, et al. (2020) and Di Capua, Runge, et al. (2020) to understand subseasonal North American hydroclimate variability, and we focus on the EAM as a regionally significant branch of the ASM based on results from Malloy and Kirtman (2022a) and Malloy and Kirtman (2022b, manuscript submitted). The subseasonal patterns related to WNPM and EAM convection in Figure 3 are different from the seasonal EAM-forced patterns from Malloy and Kirtman (2022a) and Malloy and Kirtman (2022b, manuscript submitted), demonstrating the importance of timescale for quantifying impacts (Yang et al., 2020). In addition, the definition/index and spatial scale of the drivers may affect interpretation of results; for instance, the WNPM in Di Capua, Runge, et al. (2020) was defined by maximum covariance analysis between tropical OLR and mid-latitude upper-level heights, highlighting their different approach in defining this region of active convection and its remote impacts. Nevertheless, our causal map results generally agree with the patterns from their study.

There are limitations to using the CEN, such as the causal links are only determined based on the set of drivers here. Adding other known influences of Great Plains rainfall, such as the NASH (W. Li et al., 2011; L. Li et al., 2012; Wei et al., 2019; Nieto Ferreira & Rickenbach, 2020; Malloy & Kirtman, 2022b, manuscript submitted), may change the CEN. In addition, despite the ease of using weekly-averaged indices for the CEN, there are drawbacks. Linkages considered contemporaneous on this weekly timescale may actually be causal on a sub-weekly timescale. For instance, the contemporaneous link between EAM rainfall and EAM-L (Figure 4) and the contemporaneous link between the Great Plains LLJ and rainfall (Figure 6) may be considered causal on daily timescales.

Interestingly, the EAM-L feature was important for modulating Rossby wave activity over the North Pacific, even when removing the influence of EAM. This suggests that the EAM-L feature can be forced by non-EAM activity. The EAM is only a regional branch of the ASM system. Other sub-monsoonal systems via the CGT might be impacting the variability of geopotential height activity over the EAM region (Di Capua, Kretschmer, et al., 2020; Ding & Wang, 2005; Ding et al., 2011; Zhao et al., 2018; Kornhuber et al., 2019; F. Zhou et al., 2020), and on different timescales, which should be explored further. F. Zhou et al. (2020) suggested that the EAM might maintain the CGT through latent heat release, which is supported in our causal map results. In addition, other aspects of subseasonal variability unrelated to the monsoon might be involved. For example, the NAO has been shown to modulate upper-level circulation over Eurasia (Di Capua, Kretschmer, et al., 2020; Syed et al., 2012; Wang et al., 2018).

Future work should address the subseasonal predictability of summer Great Plains rainfall via the BSISO or, more generally, wave activity over the EAM region, in climate forecast models. The CEN and causal maps with model data may reveal dissimilar causal linkages from observations, which would be valuable for understanding model biases of these teleconnections. Additionally, noting the influence of El Niño-Southern Oscillation on monsoon variability (Ding et al., 2011; F. Liu et al., 2016; Malloy & Kirt-

man, 2020) and general summertime predictability over CONUS (J. Y. Lee et al., 2011; Y. Liu et al., 2019; F. Zhou et al., 2020; Krishnamurthy et al., 2021), it would be advantageous to investigate the potential impacts warm or cool phases have on the causal pathways and link magnitudes.

## 5 Open Research

### 5.1 Data Availability Statement

All data in this study is available online. ERA5 data can be accessed through their website <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5> (Hersbach et al., 2020). The CPC Global Unified Gauge-based Analysis data was provided by the NOAA PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov> (Chen et al., 2008; Xie et al., 2007). OLR data was taken from the National Oceanic and Atmospheric Administration (NOAA) Climate Data Record from <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00875> (H.-T. Lee & Program, 2011).

The PC-MCI algorithm is publicly available and can be found by going to <https://github.com/jakobrunge/tigramite>.

### Acknowledgments

The authors would like to acknowledge that this work was supported through NOAA Grants NA15OAR4320064, NA16OAR4310141, N16OAR4310149 and NA20OAR430472 and DOE Grant DE-SC0019433.

## References

- Algarra, I., Eiras-Barca, J., Miguez-Macho, G., Nieto, R., & Gimeno, L. (2019, 2). On the assessment of the moisture transport by the great plains low-level jet. *Earth System Dynamics*, 10, 107-119. doi: 10.5194/esd-10-107-2019
- Arritt, R. W., Rink, T. D., Segal, M., Todey, D. P., Clark, C. A., Mitchell, M. J., & Labas, K. M. (1997). The great plains low-level jet during the warm season of 1993. *Monthly Weather Review*, 2176-2192.
- Chen, M., Shi, W., Xie, P., Silva, V. B., Kousky, V. E., Higgins, R. W., & Janowiak, J. E. (2008, 2). Assessing objective techniques for gauge-based analyses of global daily precipitation. *Journal of Geophysical Research Atmospheres*, 113. doi: 10.1029/2007JD009132
- Cook, K. H., Vizy, E. K., Launer, Z. S., & Patricola, C. M. (2008). Springtime intensification of the great plains low-level jet and midwest precipitation in gcm simulations of the twenty-first century. *Journal of Climate*, 21, 6321-6340. doi: 10.1175/2008JCLI2355.1
- Di Capua, G., Kretschmer, M., Donner, R. V., Van Den Hurk, B., Vellore, R., Krishnan, R., & Coumou, D. (2020). Tropical and mid-latitude teleconnections interacting with the indian summer monsoon rainfall: a theory-guided causal effect network approach. *Earth System Dynamics*, 11(1), 17-34.
- Di Capua, G., Runge, J., Donner, R. V., van den Hurk, B., Turner, A. G., Vellore, R., ... Coumou, D. (2020). Dominant patterns of interaction between the tropics and mid-latitudes in boreal summer: causal relationships and the role of timescales. *Weather and Climate Dynamics*, 1(2), 519-539.
- Ding, Q., & Wang, B. (2005). Circumglobal teleconnection in the northern hemisphere summer. *Journal of Climate*, 18(17), 3483-3505.
- Ding, Q., Wang, B., Wallace, J. M., & Branstator, G. (2011, 4). Tropical-extratropical teleconnections in boreal summer: Observed interannual variability. *Journal of Climate*, 24, 1878-1896. doi: 10.1175/2011JCLI3621.1

- Fuentes-Franco, R., Koenigk, T., Docquier, D., Graef, F., & Wyser, K. (2022). Exploring the influence of the north pacific rossby wave sources on the variability of summer atmospheric circulation and precipitation over the northern hemisphere. *Climate Dynamics*, 1–15.
- Harding, K., & Snyder, P. (2015). The relationship between the pacific–north american teleconnection pattern, the great plains low-level jet, and north central u.s. heavy rainfall events. *Journal of Climate*, 28, 6729–6742.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... Thépaut, J. N. (2020, 7). The era5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049. doi: 10.1002/qj.3803
- Higgins, R. W., Yao, Y., Yarosh, E. S., Janowiak, J. E., & Mo, K. C. (1997). Influence of the great plains low-level jet on summertime precipitation and moisture transport over the central united states. *Journal of Climate*, 481–507.
- Kornhuber, K., Osprey, S., Coumou, D., Petri, S., Petoukhov, V., Rahmstorf, S., & Gray, L. (2019). Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern. *Environmental Research Letters*, 14(5), 054002.
- Kretschmer, M., Coumou, D., Donges, J. F., & Runge, J. (2016). Using causal effect networks to analyze different arctic drivers of midlatitude winter circulation. *Journal of Climate*, 29(11), 4069–4081.
- Krishnamurthy, V., Meixner, J., Stefanova, L., Wang, J., Worthen, D., Moorthi, S., ... Stan, C. (2021). Sources of subseasonal predictability over conus during boreal summer. *Journal of Climate*, 34(9), 3273–3294.
- Lee, H.-T., & Program, N. C. (2011). *Noaa climate data record (cdr) of daily outgoing longwave radiation (olr), version 1.2*. [dataset]. National Centers for Environmental Information, NESDIS, NOAA, U.S. Department of Commerce. Retrieved from <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00875> doi: 10.7289/V5SJ1HH2
- Lee, J. Y., Wang, B., Ding, Q., Ha, K. J., Ahn, J. B., Kumar, A., ... Alves, O. (2011, 9). How predictable is the northern hemisphere summer upper-tropospheric circulation? *Climate Dynamics*, 37, 1189–1203. doi: 10.1007/s00382-010-0909-9
- Lee, S. S., & Wang, B. (2016, 7). Regional boreal summer intraseasonal oscillation over indian ocean and western pacific: Comparison and predictability study. *Climate Dynamics*, 46, 2213–2229. doi: 10.1007/s00382-015-2698-7
- Li, L., Li, W., & Kushnir, Y. (2012, 9). Variation of the north atlantic subtropical high western ridge and its implication to southeastern us summer precipitation. *Climate Dynamics*, 39, 1401–1412. doi: 10.1007/s00382-011-1214-y
- Li, W., Li, L., Fu, R., Deng, Y., & Wang, H. (2011). Changes to the north atlantic subtropical high and its role in the intensification of summer rainfall variability in the southeastern united states. *Journal of Climate*, 24(5), 1499–1506.
- Liu, F., Li, T., Wang, H., Deng, L., & Zhang, Y. (2016). Modulation of boreal summer intraseasonal oscillations over the western north pacific by enso. *Journal of Climate*, 29, 7189–7201. doi: 10.1175/JCLI-D-15-0831.1
- Liu, Y., Ke, Z., & Ding, Y. (2019, 12). Predictability of east asian summer monsoon in seasonal climate forecast models. *International Journal of Climatology*, 39, 5688–5701. doi: 10.1002/joc.6180
- Lopez, H., Lee, S. K., Dong, S., Goni, G., Kirtman, B., Atlas, R., & Kumar, A. (2019, 6). East asian monsoon as a modulator of u.s. great plains heat waves. *Journal of Geophysical Research: Atmospheres*, 124, 6342–6358. doi: 10.1029/2018JD030151
- Mallakpour, I., & Villarini, G. (2016, 6). Investigating the relationship between the frequency of flooding over the central united states and large-scale climate. *Advances in Water Resources*, 92, 159–171. doi: 10.1016/j.advwatres.2016.04.008

- Malloy, K. M., & Kirtman, B. P. (2020, 2). Predictability of midsummer great plains low-level jet and associated precipitation. *Weather and Forecasting*, *35*, 215–235. doi: 10.1175/WAF-D-19-0103.1
- Malloy, K. M., & Kirtman, B. P. (2022a, 2). The summer asia-north america teleconnection and its modulation by enso in community atmosphere model, version 5 (cam5). *Climate Dynamics*.
- Malloy, K. M., & Kirtman, B. P. (2022b, manuscript submitted, 7). East asian monsoon forcing and north atlantic subtropical high modulation of summer great plains low-level jet. *Climate Dynamics, under review*, 1–18.
- Moon, J. Y., Wang, B., Ha, K. J., & Lee, J. Y. (2013, 6). Teleconnections associated with northern hemisphere summer monsoon intraseasonal oscillation. *Climate Dynamics*, *40*, 2761–2774. doi: 10.1007/s00382-012-1394-0
- Nayak, M. A., & Villarini, G. (2017, 2). A long-term perspective of the hydroclimatological impacts of atmospheric rivers over the central united states. *Water Resources Research*, *53*, 1144–1166. doi: 10.1002/2016WR019033
- Nieto Ferreira, R., & Rickenbach, T. M. (2020). Effects of the north atlantic subtropical high on summertime precipitation organization in the southeast united states. *International Journal of Climatology*, *40*(14), 5987–6001.
- O'Reilly, C. H., Woollings, T., Zanna, L., & Weisheimer, A. (2018, 8). The impact of tropical precipitation on summertime euro-atlantic circulation via a circumglobal wave train. *Journal of Climate*, *31*, 6481–6504. doi: 10.1175/JCLI-D-17-0451.1
- Patricola, C. M., Chang, P., & Saravanan, R. (2015, 4). Impact of atlantic sst and high frequency atmospheric variability on the 1993 and 2008 midwest floods: Regional climate model simulations of extreme climate events. *Climatic Change*, *129*, 397–411. doi: 10.1007/s10584-013-0886-1
- Rogers, J. C., & Coleman, J. S. (2003). Interactions between the atlantic multi-decadal oscillation, el nino/la nina, and the pna in winter mississippi valley stream flow. *Geophysical Research Letters*, *30*(10), 1518.
- Runge, J. (2018). Causal network reconstruction from time series: From theoretical assumptions to practical estimation. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *28*(7), 075310.
- Runge, J. (2020, 3). Discovering contemporaneous and lagged causal relations in autocorrelated nonlinear time series datasets. In (pp. 1388–1397).
- Runge, J., Nowack, P., Kretschmer, M., Flaxman, S., & Sejdinovic, D. (2019). Detecting and quantifying causal associations in large nonlinear time series datasets. *Science Advances*, *5*(11), eaau4996.
- Runge, J., Petoukhov, V., & Kurths, J. (2014). Quantifying the strength and delay of climatic interactions: The ambiguities of cross correlation and a novel measure based on graphical models. *Journal of climate*, *27*(2), 720–739.
- Sardeshmukh, P. D., & Hoskins, B. J. (1988). The generation of global rotational flow by steady idealized tropical divergence. *Journal of the Atmospheric Sciences*, *45*(7), 1228–1251.
- Spirtes, P., Glymour, C. N., Scheines, R., & Heckerman, D. (2000). *Causation, prediction, and search*. MIT press.
- Syed, F. S., Yoo, J. H., Körnich, H., & Kucharski, F. (2012). Extratropical influences on the inter-annual variability of south-asian monsoon. *Climate Dynamics*, *38*(7), 1661–1674.
- Trenberth, K. E., Branstator, G. W., Karoly, D., Kumar, A., Lau, N. C., & Ropelewski, C. (1998, 6). Progress during toga in understanding and modeling global teleconnections associated with tropical sea surface temperatures. *Journal of Geophysical Research: Oceans*, *103*, 14291–14324. doi: 10.1029/97jc01444
- Wang, Z., Yang, S., Lau, N.-C., & Duan, A. (2018). Teleconnection between summer nao and east china rainfall variations: a bridge effect of the tibetan plateau.

- 557 *Journal of Climate*, 31(16), 6433–6444.
- 558 Weaver, S. J., & Nigam, S. (2008, 4). Variability of the great plains low-level jet:  
559 Large-scale circulation context and hydroclimate impacts. *Journal of Climate*,  
560 21, 1532–1551. doi: 10.1175/2007JCLI1586.1
- 561 Weaver, S. J., Ruiz-Barradas, A., & Nigam, S. (2009, 10). Pentad evolution of the  
562 1988 drought and 1993 flood over the great plains: An narr perspective on the  
563 atmospheric and terrestrial water balance. *Journal of Climate*, 22, 5366–5384.  
564 doi: 10.1175/2009JCLI2684.1
- 565 Wei, W., Li, W., Deng, Y., & Yang, S. (2019, 7). Intraseasonal variation of the sum-  
566 mer rainfall over the southeastern united states. *Climate Dynamics*, 53, 1171–  
567 1183. doi: 10.1007/s00382-018-4345-6
- 568 Xie, P., Yatagai, A., Chen, M., Hayasaka, T., Fukushima, Y., Liu, C., & Yang, S.  
569 (2007, 6). A gauge-based analysis of daily precipitation over east asia. *Journal*  
570 *of Hydrometeorology*, 8, 607–626. doi: 10.1175/JHM583.1
- 571 Yang, Y., Zhu, Z., Li, T., & Yao, M. (2020, 5). Effects of western pacific intrasea-  
572 sonal convection on surface air temperature anomalies over north america. *In-*  
573 *ternational Journal of Climatology*, 40, 2913–2923. doi: 10.1002/joc.6373
- 574 Yasunari, T. (1979). Cloudiness fluctuations associated with the northern hemi-  
575 sphere summer monsoon. *Journal of the Meteorological Society of Japan. Ser.*  
576 *II*, 57(3), 227–242.
- 577 Yasunari, T. (1980). A quasi-stationary appearance of 30 to 40 day period in the  
578 cloudiness fluctuations during the summer monsoon over india. *Journal of the*  
579 *Meteorological Society of Japan. Ser. II*, 58(3), 225–229.
- 580 Zhao, S., Deng, Y., & Black, R. X. (2018, 12). An intraseasonal mode of atmo-  
581 spheric variability relevant to the u.s. hydroclimate in boreal summer: Dy-  
582 namic origin and east asia connection. *Journal of Climate*, 31, 9855–9868. doi:  
583 10.1175/JCLI-D-18-0206.1
- 584 Zhou, F., Ren, H. L., Hu, Z. Z., Liu, M. H., Wu, J., & Liu, C. Z. (2020, 1). Sea-  
585 sonal predictability of primary east asian summer circulation patterns by three  
586 operational climate prediction models. *Quarterly Journal of the Royal Meteo-*  
587 *rological Society*, 146, 629–646. doi: 10.1002/qj.3697
- 588 Zhou, S., L’Heureux, M., Weaver, S., & Kumar, A. (2012, 4). A composite study  
589 of the mjo influence on the surface air temperature and precipitation over  
590 the continental united states. *Climate Dynamics*, 38, 1459–1471. doi:  
591 10.1007/s00382-011-1001-9
- 592 Zhu, Z., & Li, T. (2016). A new paradigm for continental u.s. summer rainfall vari-  
593 ability: Asia-north america teleconnection. *Journal of Climate*, 29, 7313–7327.  
594 doi: 10.1175/JCLI-D-16-0137.1
- 595 Zhu, Z., & Li, T. (2018, 5). Amplified contiguous united states summer rainfall vari-  
596 ability induced by east asian monsoon interdecadal change. *Climate Dynamics*,  
597 50, 3523–3536. doi: 10.1007/s00382-017-3821-8

**Figure 1.** Composited anomalies of OLR (shaded) and U200 (purple contours) anomalies for BSISO combined phases (a) 8+1, (b) 2+3, (c) 4+5, and (d) 6+7. U200 anomalies are contoured every  $1 \text{ m s}^{-1}$  between  $-5$  and  $5 \text{ m s}^{-1}$ .

**Figure 2.** Anomalous probability of the following events for days after a BSISO phase: (a) below-normal Great Plains rainfall, (b) weak Great Plains LLJ, (c) PNA+ pattern, (d) above-normal Great Plains rainfall, (e) strong Great Plains LLJ, and (f) PNA- pattern. White dots denote statistical significance at the 90% confidence level determined by bootstrapping method with 1000 iterations.

**Figure 3.** Lag correlation between Great Plains precipitation index at  $T = 0$  and (a-c) OLR anomalies at  $T = 0, -10$ , and  $-20$  days, (d-f) V850 anomalies at  $T = 0, -10$ , and  $-20$  days, and (g-i) Z200 anomalies at  $T = 0, -10$ , and  $-20$  days. Pink domains indicate the Great Plains precipitation, EAM rainfall, and WNPM rainfall indices. Green domain indicates the Great Plains LLJ index, and black contour outline highlight the southerly flow over the relevant BSISO region. Orange domains indicate the EAM-L, NPac-L, and PNA-HL indices. Stippling indicates statistical significance at the 90% confidence level and a correlation value  $> 0.05$  or  $< -0.05$ . See Table 1 for more information about indices.

**Figure 4.** Causal effect network between WNPM rainfall, EAM rainfall, EAM-L, and NPac-L. Color of individual nodes indicates autocorrelated  $\sigma$  change from one week to the next. Color of lines or arrows indicate the  $\sigma$  change. Arrows indicate the direction of causality, with strength of  $\sigma$  change annotated on arrow, with lag of one week. Dashed lines are contemporaneous links, which, by themselves, do not imply causality.

**Figure 5.** Same as Fig. 4, but for the causal effect network between EAM-L, NPac-L, and PNA-HL.

**Figure 6.** Same as Fig. 4, but for the causal effect network between NPac-L, PNA-HL, Great Plains LLJ, and Great Plains rainfall.



**Figure 7.** Causal maps showing the causal link value between (a) WNPM rainfall at  $T = -1$  (one week before) and Z200 at every grid point, with EAM rainfall conditioned out, (b) EAM rainfall at  $T = -1$  and Z200 at every grid point, with WNPM rainfall conditioned out, (c) EAM rainfall at  $T = -1$  and Z200 at every grid point, with EAM-L conditioned out, and (d) EAM-L at  $T = -1$  and Z200 at every grid point, with EAM rainfall conditioned out. Causal link value is interpreted the same as arrows in Figs 4-6. Only values with significance at 95% confidence level and a magnitude  $> 0.05$  are shown.

**Figure 8.** Same as Fig. 7, but for links with V850 at every grid point.

**Figure 9.** Same as Fig. 7, but for links with OLR at every grid point.

**Figure 10.** 200-hPa Rossby Wave Source anomaly (shaded), with 200-hPa divergence anomaly (gray contours) and U200 anomaly (black contours) overlaid, for (a) upper tercile WNPM days and (b) upper tercile EAM days. Divergence anomalies are contoured every  $2 \times 10^{-6} \text{ s}^{-1}$  between 0 and  $4 \times 10^{-6} \text{ s}^{-1}$ , and U200 anomalies are contoured every  $1 \text{ m s}^{-1}$  between -3 and  $3 \text{ m s}^{-1}$ .