


Deep brittle-ductile transition, water percolation into the mantle and reactivation of weakness zone: new insights of the seismicity on St. Paul Transform System, Equatorial Atlantic

Guilherme de Melo^{1*}, Aderson do Nascimento¹, Marcia Maia², Julie Perrot², Alexey Sukhovich², Jean-Yves Royer², Thomas Campos³

1-Department of Geophysics, Federal University of Rio Grande do Norte, Natal, RN, Brazil; 2-Institut Universitaire Européen de La Mer, UBO, Plouzané, France, 1-Department of Geology, Federal University of Rio Grande do Norte, Natal, RN, Brazil;

*gws Melo@ufrn.edu.br



Introduction


In Equatorial Atlantic, a complex multifault system of four transform faults, bounding three short intra-transform segments form the St. Paul Transform System (SPTS) (Maia et al., 2016). SPTS is a complex group of four transform faults, bounding three short intra-transform segments located at the Equatorial Atlantic. In the northern transform fault of SPTS, the 200 km long and 30 km wide Atoba Ridge is a major topographic feature that reaches the sea level at the St. Peter and St. Paul Archipelago (SPSPA). We have determined the hypocentral location of 62 minor-moderate earthquake (>ML 1.6), using records of three hydrophones and one seismograph.

Geological Setting

In northern of SPTS, the transform present an offset with the two adjacent ridges segments in 630 km with a slow-spreading rate of 32 mm/year (DeMets et al., 2010; Maia et al., 2016). Along this segment is located the Atobá Ridge, a 200 km long and 30 km major topographic formed for a push-up due to transpressive stress that started 10 million years ago (Maia et al., 2016). The structure is formed for exposed peridotite rocks, that reach until the sea level 3500 m above seafloor, in the SPSPA islets. The set of islets is composed for peridotite mylonite and alkaline ultramafic with different serpentinization degree (Campos et al., 2010).

Objectives


Our objective is to investigate the estimates the seismicity that occur along the St. Paul Transform System, in Equatorial Atlantic. Using earthquake data, we estimate the brittle-ductile transition depth along this transform fault, limit depth of serpentinization, and also show reactivation on weakness zones in the SPTS (de Melo et al., in review).



2


Data Acquisition

Earthquakes were analyzed using recorded data during six months (January-July) by a single seismographic station located on St. Peter and St. Paul Archipelago (ASPSP) and three hydrophones deployed in 2013 as part of COLMEIA cruise (Maia et al., 2016) that obtained records of P waves.



Analysis

SEISAN package was used in the analysis (Havskov and Ottemöller., 1999). P and S seismic waves were analyzed in ASPSP station, together the P waves recorded by the hydrophone array. Waveform data were bandpass filtered at 4-18 Hz in ASPSP station, and 2-12 Hz to the hydrophone. Back-azimuth and angle of incidence method (Havskov and Ottemöller., 2010) was used to estimated the hypocenter catalogue in ASPSP station. Angle of incidence were measured using Seismic Analysis Software (Helffrich et al., 2013). P waves recorded by the hydrophones were analyzed applying the travel time correction because the change solid-to-acoustic that occur in wave propagation. This delay was added to account for the length of the mooring cable, using a constant water velocity from the global ocean sound speed model (GDEM).




3


Earthquake catalog

62 earthquakes were recorded by the ASPSP station with magnitudes between 1.9 and 5.3 M. Among these, 15 events with $M > 2.9$ were recorded also by the hydrophone array. Example of these events are shown in (a), (b) and (c) figures.



Event A - St. Paul Transform System ML=2.1 05/03/2013 15:36:52 1.023 -29.305



Event B - St. Paul Transform System ML=3.1 05/04/2013 05:29:52 0.986 -29.179



Event C - St. Paul Transform System Mw=5.3 04/03/2013 05:30:58 1.041 -27.201




Geodynamic

- 50 earthquakes with 0.2 km being the minimum depth and the maximum of 17.8 km were located in the Atobá Ridge Zone (ARZ) between longitudes 29.6° and 29.1° W. This high number is probably due to the high transpressive stress zone and the presence of a large number of faults that concentrate the deformation near the core of the push-up ridge.
- In the East Shear Zone (ESZ) were recorded only three earthquakes, with depths of 10.2 until 15.5 km. ESZ section is a deep basin crossed by the principal transform displacement zone, which is here more linear and less segmented than to the West of SPTS (Maia et al., 2016);
- Nine earthquakes were located at Central Fracture Zone (CFZ). They are distributed between longitudes 29.1° and 28.4°. These earthquakes possibly are linked to the reactivation of this segment of the fracture zone between the North and Central segments of the St. Paul Transform System, associated with possible zone of weakness reactivated due to the regional the transpressive stresses uplifting the Atoba Ridge..



Hypocenter location

Earthquakes located by the ASPSP single station present error latitude (ERLT) from 1.5 until 4.5 km, and error longitude of 1.5 until 4.6 km (ERLN). The depth error (ERDP) was estimated with an uncertainty of $\pm 3^\circ$ to incidence angle, that presented error range 3-7 km. 15 events using hydrophone array together presented ERLT from 2.5 until 13.9 km, ERLN from 1.4 until 9.7 and ERDP range 1.3-9.7 km. Then, the final catalogue is defined by the 15 events located together hydrophone and another 47 using only ASPSP station.



Serpentinization

The thickness of the low density layer in the East Atoba (near 29°25'W) is highly variable, reaching 17 km. We identify a correlation between the crustal thickness of low density layer and the depths of the earthquakes at the Atoba Ridge suggesting that the deeper events may be linked to a fluid interaction with the mantelic rocks caused by water percolation down the thrust faults that border the push up ridge and the occasion a fluid interaction with the mantelic rocks and after the faults develops into the mantle. (a), (Bonatti., 1976; Escartin et al., 1997; Roland et al., 2010; Maia et al., 2016). Maia et al (2016) showed that the density pattern correlates with a deepening of the 500-600°C isotherms and that the thick low density layer probably corresponds to serpentinized mantle rocks found in the Atobá Ridge and SPSP islets.



Brittle-Ductile Transition

The hypocenter depth can be controlled by the $\sim 600^\circ\text{C}$ isotherm, defining the brittle-ductile transition. This was discussed by several authors (Abercrombie and Ekström., 2001; Schlindwein and Schmid., 2016). We compare the 600°C isotherm depth (in a plot of offset age against depth) reported by Boettcher and Jordan (2004), and the curve of the maximum observed hypocentral depth of St. Paul transform system. Considering the thermal lithosphere structure changes which occur when the isothermal limit depth (600°C) increases as the fault length, we view a deep brittle-ductile limitation in the St. Paul Transform System (15-18 km).

c

Transform	Latitude(°)	Longitude(°)	Length(km)	Age Offset(y)	Hypocenter_max(km)	600°C isotherm(km)	References
Oceanographer	35.150	-35.960	105	10	9	11.5	Cesaro and Haxel (1986), Bergman and Solomon (1986)
Hayes	33.630	-38.530	75	6	8	12.2	Bergman and Solomon (1986)
Kane	23.760	-45.860	150	11	8	11.4	Bergman and Solomon (1986)
Fifteen-Twenty	15.310	-45.960	180	13	12	13.5	Cleveland et al (2016)
Mauritron	12.650	-44.660	80	5	8.5	12	Smith et al (2008), Cleveland et al (2016)
St Paul(North Segment)	0.950	-29.000	310	14	15.5	20.1	Maia et al (2016), Hypocenter depth of this study
Chall	-1.240	-14.645	300	20	11	15.1	Abercrombie and Ekström (2001), Harmon et al (2016)

Conclusions

- Earthquakes occur regularly in the St. Paul Transform System with hypocenters reaching 17.8 km below the seafloor
- A thicker lithosphere in SPTS with a 600°C isotherm limit situated in 15-18 km depth.
- The correlation between the low density layer and the hypocenter depths present an extensive serpentinization beneath the Atobá Ridge reaching depths of 18 km.
- Earthquakes at the CFZ possibly is caused by a reactivated zone of weakness, due to the regional transpressive stresses uplifting the Atoba Ridge that contributed to the reactivation of an otherwise inactive fracture zone.

Selected References

Abercrombie and Ekström. (2001), *Nature*, **410**, 74-77
de Melo and do Nascimento. (2018), *Pure and Applied Geophysics*, **178**, 741-756
de Melo et al., *Journal of Geophysical Research: Solid Earth*, in review
Boettcher and Jordan. (2004), *Journal of Geophysical Research: Solid Earth*, **109**, 1-21
Maia et al. (2016), *Nature Geoscience*, **9**, 619-624
Roland et al. (2010), *Geochemistry, Geophysics and Geosystems*, **11**, 1525-2027

Acknowledgements

G.W.S. de Melo thanks PETROBRAS for a MSc via the PHR-229 Capacity Building Programme. G.W.S. de Melo also grateful LabexMER ANR-10-LABX-19-01 and PPG 110/2018 by the financial support to his708 internship at the France. We authors thank the Brazilian Navy and CNPq (Brazilian Funding Agency) for logistical and financial support. The COLMEIA experiment and cruise were funded through the grants to the Flotte Océanographique Française and grants from CNRS-INSU Géosciences Marines and LabexMER Actions à la Mer.



de Melo and do Nascimento (2018)