

Resistance, recovery, and resilience:

rethinking the three Rs of survival in the Anthropocene

Benjamin W. Abbott¹, Kristen L. Underwood², Erin C. Seybold³, Dustin W. Kincaid⁴, Scott D. Hamshaw⁴, Raymond M. Lee¹, Donna M. Rizzo⁴, Brian Brown⁵, Regina Toolin⁶, Jon Chorover⁷, Li Li⁸, Sayedeh Sara Sayedi¹, Samuel St. Clair¹, Gabriel Lewis⁹, Rachel L. Buck¹⁰, Zachary T. Aanderud¹, Janice Brahney¹¹, Ryan S. Nixon¹², Weihong Wang¹³, Cally Flox¹⁴, Julia Perdrial⁴

¹Brigham Young University, Department of Plant and Wildlife Sciences, Provo, USA

²University of Vermont, Department of Civil & Environmental Engineering, Burlington, VT, USA

³Kansas Geological Survey, University of Kansas, Lawrence, KS

⁴University of Vermont, Department of Civil & Environmental Engineering, Burlington, VT, USA

⁵Brigham Young University, Department of Computer Science, Provo UT, USA

⁶University of Vermont, Department of Education, Burlington, VT, USA

⁷University of Arizona, Department of Environmental Science, Tucson, AZ, USA

⁸Penn State University, Department of Civil and Environmental Engineering, University Park, USA

⁹University of Nevada, Reno, Natural Resources and Environmental Sciences, Reno, NV, USA

¹⁰Brigham Young University, Department of Biology, Provo, USA

¹¹Utah State University, Department of Watershed Sciences and Ecology Center, UT, USA

¹²Brigham Young University, Department of Teacher Education, Provo, UT, USA

¹³Utah Valley University, Department of Earth Science, Orem, Utah, USA

¹⁴Brigham Young University, McKay School of Education, CITES Department, Provo, Utah USA

Key words: Ecosystem, Critical Zone, Resilience, Earth Stewardship, Sustainability, Traditional Ecological Knowledge, State Factors, Dynamical Systems, Nature Positivity, Anthropocene

Abstract:

The concepts of resistance, recovery, and resilience are in diverse fields from behavioral psychology to planetary ecology. These “three Rs” describe some of the most important properties allowing complex systems to survive in dynamic environments. However, in many fields—including ecology—our ability to predict resistance, recovery and resilience remains limited. Here, we propose new disturbance terminology and describe a unifying definition of resistance, recovery, and resilience. We distinguish *functional disturbances* that affect short-term ecosystem processes from *structural disturbances* that alter the state factors of ecosystem development. We define resilience as the combination of resistance and recovery—i.e., the ability of a system to maintain its state by withstanding disturbance or rapidly recovering from it. In the Anthropocene, humans have become dominant drivers of many ecosystem processes and nearly all the state factors influencing ecosystem development. Consequently, the resilience of an individual ecological parameter is not an inherent

attribute but a function of linkages with other biological, chemical, physical, and especially social parameters. Because every ecosystem experiences multiple, overlapping disturbances, a *multidimensional resilience* approach is needed that considers both ecosystem structure (configuration of linkages) and disturbance regime. We explore these concepts with a few case studies and recommend analytical tools and community-based approaches to strengthen ecosystem resilience. Disregarding cultural and social dimensions of disturbance regimes and ecosystem structures leads to undesirable outcomes, particularly in our current context of intensifying socioecological crises. Consequently, cultivating reciprocal relationships with natural disturbance regimes and ecosystem structures is crucial to Earth stewardship in the Anthropocene.

Introduction

The paradox, in a nutshell, is this: humans have grown so powerful that they have become a force of nature - and forces of nature are those things which, by definition, are beyond the power of humans to control.

-Oliver Morton, *The Planet Remade*, 2015

The history of the Earth system is a remarkable story of life causing, responding to, and adapting to catastrophic changes (Schlesinger & Bernhardt 2020). In the dynamic environment of our planetary home, the organisms and ecosystems not suited to disturbance are rare or nonexistent. From individual cells to human societies to the entire biosphere, every aspect of the Earth system is shaped by change.

In the Anthropocene, humans have emerged as a force of nature in a way that perhaps no vertebrate organism ever has (Lewis & Maslin 2015; Keys *et al.* 2019; Folke *et al.* 2021). Humans have influenced much of Earth's terrestrial surface for more than ten thousand years (Ellis 2021), but in the past few centuries, we have become the primary force structuring Earth's habitats, biogeochemical cycles, and disturbance regimes (Steffen *et al.* 2015a; Watson *et al.* 2018; Schlesinger & Bernhardt 2020). Humans are now the largest driver of the extinction and evolution of species, and we have shifted patterns of sediment transport, nutrient cycling, carbon balance, climate, water cycling, and wildfire at global scales (Wilkinson 2005; Benson 2012; Steffen *et al.* 2015b; Cooper *et al.* 2018; Abbott *et al.* 2019b; Hurteau *et al.* 2019). Our physical creations outweigh all life

on Earth (Elhacham *et al.* 2020), our bodies and livestock account for ~93% of total vertebrate biomass (Bar-On *et al.* 2018), and we have created novel planetary material cycles, including plastics and persistent organic pollutants, with largely unknown impacts on human health and ecosystem functioning (Nizzetto *et al.* 2010; Bank & Hansson 2019; Hannah *et al.* 2022). From changing the structure of the thermosphere to triggering tectonic tremors (Manney *et al.* 2011; Wilson *et al.* 2017; Mlynczak *et al.* 2022), our direct and indirect footprints have altered all the Earth's aquatic, terrestrial, marine, and subsurface environments (Watson *et al.* 2018; Díaz *et al.* 2019; Kolbe *et al.* 2019; Bochet *et al.* 2020; Ellis *et al.* 2021). The land-cover transformation, amplification of biogeochemical flows, and climate disruption that characterize the Anthropocene are triggering transformations that are likely unprecedented in our planet's past (Diffenbaugh & Field 2013; Kemp *et al.* 2015; Ceballos *et al.* 2020; Armstrong McKay *et al.* 2022; Fricke *et al.* 2022).

The combined effects of these Earth system alterations have caused catastrophic global consequences, including diminished quality of life for humankind (Fig. 1). There has been a pervasive decline of species on Earth in aquatic, terrestrial, and marine environments (Vörösmarty *et al.* 2010; Díaz *et al.* 2019; Fricke *et al.* 2022). Environmental pollution, primarily from burning fossil fuels, causes more than 15 million premature human deaths annually—one in four deaths each year (Errigo *et al.* 2020; Vohra *et al.* 2021). This means that our unhealthy relationship with the Earth directly causes more deaths than all violence, malnutrition, and communicable diseases combined (Landrigan *et al.* 2017; Errigo *et al.* 2020; Fuller *et al.* 2022). Ongoing ecosystem state changes threaten the future of billions of people across every country and socioeconomic condition (Abatzoglou & Williams 2016; Van Loon *et al.* 2016; Dupas *et al.* 2019; Mu *et al.* 2020; Cheng *et al.* 2022; Hannah *et al.* 2022). Our individual and communal survival depends on restoring positive and reciprocal relationships between human societies and the ecosystems we have come to dominate (Kimmerer 2002; Sandifer *et al.* 2015; Bradshaw *et al.* 2021; Chapin *et al.* 2022). In this context of accelerating planetary disruption, understanding how ecosystems respond to change is more critical than ever.




Figure 1. Signs and symptoms of planetary vulnerability in the Anthropocene. Data for specific claims drawn from (Watson et al. 2018; Abbott et al. 2019a; Díaz et al. 2019; Errigo et al. 2020; Bradshaw et al. 2021; Ritchie et al. 2021; Vobra et al. 2021; Armstrong McKay et al. 2022; Fuller et al. 2022).

Disturbance, succession, and equilibrium have been central themes of ecology since it emerged as a quantitative science in the 20th century (Tansley 1935; Lindeman 1942; Turner *et al.* 1989; Chapin *et al.* 1994). Across multiple natural and social sciences, a wealth of terminology has developed describing the characteristics of disturbance and system response to ecological and evolutionary change (Callicott & Mumford 1997; Carpenter *et al.* 2001; Redman 2014; Larsson & Abbott 2018; Elmqvist *et al.* 2019; Fuller *et al.* 2019; Barbe *et al.* 2020; Frei *et al.* 2020). However, our ability to predict ecological state changes, such as the collapse of a population or loss of an important ecosystem process, remains limited (Jasinski & Payette 2005; Scheffer *et al.* 2009; Marlon 2020; Schoolmaster Jr. *et al.* 2020; Gouveia *et al.* 2021; Ritchie *et al.* 2021). While deterministic modeling of stochastic events in complex Earth systems has long been out of reach, advances in monitoring and analysis now allow deeper characterization and better prediction of emergent changes and nonlinearities (Loehle 2006; Beven & Alcock 2012; Lum *et al.* 2013; Brunton *et al.* 2016). The development and simplification of multiple sensing technologies have significantly expanded our ability to measure individual and composite vital signs of global ecosystems, including traditional ecological data and near-real-

time indices of how information and emotions are moving through human communication networks (Abbott *et al.* 2016; Rode *et al.* 2016; Newman 2017; Zhang *et al.* 2022). At the same time, the development of an extraordinary range of complex systems tools has dramatically enhanced our ability to interpret multivariate data (Barbe *et al.* 2020; Underwood *et al.* 2021; Brunton *et al.* 2022; Heddam *et al.* 2022).

In this context, we convened a group of interdisciplinary researchers and educators to explore how human perception and management of ecosystems affect ecological resilience and vulnerability in the Anthropocene. We begin by presenting new terminology for describing disturbance and then propose a unified framework around what we call the three Rs of survival in the Anthropocene: resistance, recovery, and resilience. Based on definitions from the fields of sustainable development and fluvial geomorphology (Meerow *et al.* 2016; Fuller *et al.* 2019), we define resilience as the combination of resistance and recovery—i.e., the ability of an ecosystem to maintain its state by withstanding disturbance or rapidly recovering from it. We hypothesized that resilience measured in an individual ecological variable is not an inherent attribute but a function of linkages with other social, biological, chemical, and physical parameters, including the disturbance regime (Turner *et al.* 2003; Chapin *et al.* 2022). We present ecological case studies and assess the potential of analytical tools to characterize multidimensional resilience and inform applied solutions. We conclude that successful ecological restoration and planetary sustainability depend on cultivating an ethic of Earth stewardship that recognizes and rehabilitates humanity's unique roles in the Earth system (Steffen *et al.* 2011; Palmer & Stewart 2020; Locke *et al.* 2021; Rockström *et al.* 2021; Chapin *et al.* 2022).

Resilience vocabulary

An advantage and challenge of resilience terminology is its familiarity. Resistance, resilience, and recovery are commonly used to describe a wide range of technical and nontechnical phenomena (Carpenter *et al.* 2001; Allison 2004; Rogers *et al.* 2012; Shade *et al.* 2012; Anderies *et al.* 2013; Elmqvist *et al.* 2019). We recognize the utility and origin of multiple definitions and do not seek to invalidate their use. For the purposes of this paper, we propose the most intuitive and direct meanings based on our opinion and recent scholarship (Chapin *et al.* 2012; Meerow *et al.* 2016; Fuller *et al.* 2019). We note that while some sustainability researchers use a version of the term social-ecological systems (SES) to emphasize human-environment

interactions (Anderies *et al.* 2013; Chapin *et al.* 2013; Folke *et al.* 2016), we use the terms ecosystem and ecological as fully inclusive of human dimensions of the Earth system. This is in line with the original definition of the ecosystem concept, and we use these terms deliberately to erode what we see as an unhelpful distinction between society and ecosystems (Tansley 1935; Chapin *et al.* 2012; Abbott *et al.* 2019b). While sustainable development frames economy, environment, and society as competing interests, an Earth stewardship or nature-positive approach sees economy as a nested component of society and society as an embedded and intertwined part of the Earth system (Folke *et al.* 2016; Locke *et al.* 2021; Chapin *et al.* 2022). Human society only exists within ecosystems, and it is impossible to meaningfully study ecosystems in the Anthropocene without considering society.

An *ecological threshold* describes the boundary between two ecological states or sets of conditions, and a *state change* describes an ecosystem crossing such a threshold, e.g., forest to grassland or clear-water to turbid (Carpenter *et al.* 2020; Cassidy *et al.* 2022). *Ecological resistance* is the capacity to avoid crossing a threshold during or immediately after disturbance. *Ecological recovery* describes the tendency, degree, and rate of return to pre-disturbance conditions after perturbation. *Ecological resilience* is the combination of resistance and recovery, which therefore describes the likelihood of an ecosystem or ecological variable to be found in a particular state throughout time. *Ecological vulnerability* is the inverse of ecological resilience, describing a system's tendency to transition and stay in a different state. These concepts are summarized visually in Figure 2.

Disturbance is often characterized by intensity, duration, timing, frequency, rate of change, extent, and patchiness. These terms are already quite intuitive, though highly dependent on the observed spatiotemporal scale and resolution (Glasby & Underwood 1996; Poff *et al.* 1997; Kemp *et al.* 2015; Collins *et al.* 2018; Meerow & Newell 2019). For example, a disturbance could be characterized as either a press or a pulse, where the former comes on slowly but potentially lasts longer (low rate of change, long duration), and the latter comes on fast but does not last as long, relative to the timescale of interest (Bergstrom *et al.* 2021). Multiple characteristics of a single disturbance type are often described as the disturbance regime (Mack & D'Antonio 1998; Turner *et al.* 2003; North & Keeton 2008). However, for our purposes, we distinguish between *disturbance characteristics* of an individual disturbance type (e.g., wildfire frequency, extent, severity etc.)

and the *disturbance regime* of an ecosystem, which always includes multiple interacting disturbance types (e.g., wildfire, acidification, logging, climate change, invasive species, etc.) (Atkins *et al.* 2020).




Figure 2. Diagrams of the disturbance and resilience concepts described in this paper. a) Depictions of Ecosystem states (yellow circles), thresholds (orange lines), disturbance types, and response surfaces representing resistance and recovery to disturbance. Functional disturbances change the current ecosystem state, while structural disturbances affect the interacting state factors that regulate the response of the ecosystem to disturbance. b) Top-down view of multiple dimensions of ecosystem state on their respective response surfaces, including feedbacks and thresholds, with thresholds near the center of the diagram representing more vulnerable dimensions. Exceeding a threshold in one dimension is likely to modify the condition and response surface of others, i.e., create a structural disturbance.

We think it is helpful to introduce new terminology for both individual disturbances and disturbance regimes. The state factor concept was originally developed for predicting soil formation (Jenny 1941; Florinsky 2012), and through time it has been applied to ecosystem development and structure (Chapin *et al.* 2012; Tank *et al.* 2020). This concept predicts that a set of initial ecological conditions or *state factors* strongly constrain the development of an ecosystem (Fig. 3). Useful predictions about ecosystem type and processes are possible with knowledge of these state factors: parent material, potential biota, climate, topography, and

time since the last major disturbance (Jenny 1941). Human activity has been proposed as an additional state factor, given the extent of anthropogenic influence in the Anthropocene (Chapin *et al.* 2012). We distinguish *functional disturbances* that affect short-term ecosystem processes from *structural disturbances* that alter the state factors of ecosystem development (Jenny 1941; Florinsky 2012; Tank *et al.* 2020). Conversely, disturbances that primarily affect current ecosystem processes would be described as *functional disturbances* (Figs. 2 and 3). This distinction might be informative because it indicates whether a disturbance is likely to affect the short-term status of an ecosystem (e.g., does the functional disturbance exceed the ecological resistance for a given parameter) or the long-term recovery trajectory (e.g., is the structural disturbance severe enough to alter the multidimensional response surface guiding recovery).

We recognize that many disturbances—and especially those controlled by humans—have both functional and structural dimensions. Indeed, there is a continuum between ecosystem processes and state factors depending on the severity of the disturbance and the successional timescale of interest. For example, what might seem like an ephemeral ecosystem process to the geomorphological evolution of a watershed could be an effectively permanent state factor from the perspective of a microbial community (Fisher *et al.* 1998; Shade *et al.* 2012).

Combining these concepts, we define *multidimensional resilience* as an ecosystem's ability to maintain its state under a current or future disturbance regime through a combination of resistance and recovery. We hypothesize that multidimensional resilience depends on ecosystem structure—the configuration of linkages among the state factors and ecosystem processes—which itself is influenced by the disturbance regime (Adler 2019; Frei *et al.* 2020; Mo *et al.* 2020). More specifically, we hypothesize that the physical, chemical, and biological structures of the critical zone—the portion of the ecosystem from unweathered bedrock to the vegetation canopy (National Research Council 2001; Chorover *et al.* 2007)—strongly influence its resilience and vulnerability (Figs. 1-3). For this concept of multidimensional resilience to be relevant for research or management, human participation in the physical and biological structure of the critical zone must be integrated (Chapin *et al.* 2022).

Because many of these terms have emotional connotations in nontechnical usage, we point out that the disturbance and resilience terminology presented above does not connote desirability or ecological value (Elmqvist *et al.* 2019). For example, resilience of can be negative (i.e., unhelpful) when present in undesirable aspects of the system, such as antisocial trends of disregard for the environment or fellow humans. Likewise, a specific disturbance can be positive or negative depending on the ecosystem structure (including human needs and goals) and broader disturbance regime.




Figure 3. Conceptual diagram of ecosystem development adapted from Chapin et al., (2012). We have added the distinction between structural and functional disturbances as well as the effect of human activity on state factors.

In the following paragraphs, we elaborate these concepts with examples from catchment hydrology and freshwater biogeochemistry to evaluate how ecosystem structure (i.e., the configuration of social, biological, chemical, and physical attributes in the critical zone) influences the timing, direction, and intensity of linkages among multiple responses and consequently multidimensional resilience.

Case Study 1: Artificial resistance through erosion control

Because humans have long congregated along river networks, flood control and fluvial erosion have been areas of focus in ecosystem management for centuries (Allaire 2016; Fang & Jawitz 2019; Tate *et al.* 2021; Sanders *et al.* 2022; Syvitski *et al.* 2022). Human efforts to control rivers and floodplains have yielded both benefits and major problems, including environmental injustice and substantial loss of life (Reisner 1993; Tate *et al.* 2021; Sanders *et al.* 2022; Sowby & Hotchkiss 2022). This highlights the need to consider human culture and infrastructure as integrated components of ecosystems, with similar unanticipated behaviors (Leavitt & Kiefer 2006; South *et al.* 2018; Wohl 2019; Wang & He 2022).

The northeastern United States provides well-documented examples of multiple agrarian and industrial disturbances of river networks (Wolock 1995; Armfield *et al.* 2019). In this region and many areas globally, the provisioning of clean water for drinking, agriculture, and aquatic ecosystems is threatened by low geomorphological resistance to changes in river flow (Davis *et al.* 2009; Abbott *et al.* 2018a; Zarnetske *et al.* 2018; NASEM 2020). Two examples of vulnerability are 1) headwater stream networks with susceptibility to hillslope and channel erosion due to glacial history, and 2) valley and piedmont river corridors with large legacy sediment stores that are coupled closely with receiving waters (Pinay *et al.* 2018; Dearman & James 2019). The legacy of glacial and ice sheet retreat has created bouldery tills and fine glacio-lacustrine clays. This combination of high energy streams that can come into contact with glaci-lacustrine clays through streambank or bed erosion (Davis *et al.* 2009) creates significant stream management challenges. The postglacial context creates low resistance but high recovery regarding sediment transport. Exceeding modest thresholds of stream movement during high streamflows can trigger multiple problems including mass movement (landslides) and persistent high turbidity levels in downstream drinking water reservoirs.

While these environments regularly transported large amounts of sediment naturally during the Holocene, European settlement altered sediment sources and sinks. Forest clearing for agriculture and the construction of many small dams resulted in the accumulation of sediments along the river corridors (Dearman & James 2019; Johnson *et al.* 2019; Jiang *et al.* 2020; Noe *et al.* 2020). This combination of land use

and vulnerable critical zone structure is now threatening the provisioning of drinking water for millions of people in the New York metropolitan area.

Over the past several decades, stream and watershed management efforts have been designed to counterbalance the overabundance of sediment sources in river systems to maintain. Engineering-oriented techniques initially focused on increasing resistance to erosion (Fig. 4), including armoring streambanks and hillslopes and dredging to temporarily increase flood conveyance (Bernhardt & Palmer 2007; Wohl *et al.* 2015). However, these techniques have proven very short lived given the artificial disequilibrium (e.g., legacy sediments) and natural characteristics of the critical zone structure (e.g., high sediment availability in postglacial landscapes). Management interventions have so far largely treated symptoms rather than causes while also creating greater problems upstream and downstream of the hard-part interventions. However, because of the high societal value of the drinking water provisioned by this ecosystem, the inefficient management approach has been acceptable (Davis *et al.* 2009; NASEM 2020). The question is whether society will continue to support this kind of active river management or call for a change. The underlying hypothesis has been that sufficient resources (financial and human capital) are available to respond to the shifting disturbance regime (greater magnitude and intensity of storms, increased persistence and magnitude of precipitation) with ecohydrological expertise continually nudging the system back towards a more “natural” equilibrium in an effort to create a more resistant critical zone structure. As such, watershed and drinking water managers have prioritized extensive mapping of glacial tills and clays and initiated an active management program including streambank and hillslope stabilization, floodplain reconnection, and full-channel restoration (NASEM 2020).

Seeking to enhance river system resilience by maximizing resistance can create rigidity that results in continual or ever increasing management costs and decreasing ecosystem function and safety (Fig. 4). Seeking to preserve or restore local disturbance regimes—including sustainable human land use and other activities—is a much more robust approach with many more co-benefits (Bishop *et al.* 2009; Christianson 2015; Houlton *et al.* 2019). However, overlying regional disturbance regimes that include increasing flow magnitudes and changes in precipitation patterns may require more frequent stabilizing feedbacks from active watershed management in order to maintain clean water provisioning. This highlights the importance of cultivating

more meaningful and multidimensional relationship between local societies and the ecosystems they depend on. This avoids undue focus on a single ecosystem service, such as seeing a watershed primarily or exclusively as a drinking water provisioning device.




Figure 4. Conceptual examples of how changing disturbance regimes and intentional modification of ecosystem structure can lead to greater vulnerability. Managing for resistance (i.e., modifying structure to impose physical constraints on the system and its dynamic ecosystem states) often leads to rigidity that can result in catastrophic transformations when the system is subjected to a new disturbance regime with increased amplitude of disturbance (e.g., higher flood magnitude).

This case study showcases a broader shift toward *naturalness* as a more resilient and cost-effective management strategy in dynamic environments (Bishop *et al.* 2009; Palmer & Stewart 2020). Recent stream restoration practices recommend restoring naturalness to disturbance regimes by removing obstructions (dams, berms, levees) and buying out flood damaged homes to allow the river system more room to dynamically adjust to increased flows (Fig. 5). This management shift is informed by observations that the more altered and artificial a system is, the more rigid and high maintenance it tends to be (Bishop *et al.* 2009). Additionally, more extreme modifications of critical zone structure and disturbance regime create more severe tradeoffs and compromises (Palmer & Stewart 2020; Abbott *et al.* 2021a).

Case study 2: Coastal forests and sea level rise

Human-caused sea level rise from ice melt and thermal expansion has progressed much faster than expected and is currently tracking the most extreme model projections (King *et al.* 2020; Slater *et al.* 2020; Boers & Rypdal 2021; Heinze *et al.* 2021). This quintessential press disturbance is interacting with the pulse disturbances of extreme storms (Crandall *et al.* 2021; Fowler *et al.* 2021; IPCC 2021). Coastal forests have been categorized into two bands based on proximity to the ocean (Fagherazzi *et al.* 2019; Kearney *et al.* 2019; Mo *et al.* 2020). Stands of mature trees that established before major sea level rise and storm intensification can be found within a meter above the normal high tide. These stands are resistant to storm surges because the adult trees can survive temporary inundation by salt water, partly by accessing fresh groundwater. However, they are not resilient because recruitment cannot occur in salinized soil. As windfall and old age kills adult trees, the mature stands are overtaken by marshes that are more able to survive frequent seawater inundations and take advantage of the increased light availability (Fagherazzi *et al.* 2019).

Above the mature, resistant zone near the ocean, there is an area described as the Regenerative Zone because tree recruitment is still occurring (Kearney *et al.* 2019; Paldor *et al.* 2022). This zone is more distal and higher in elevation, meaning the storm surges less frequently introduce ocean waters and the degree of salinity in soils is less and within the tolerances of germination and seedling recruitment.

This case study shows the interaction between anthropogenic structural disturbances and a relatively unmanaged ecosystem. Sea level rise and storms are interacting structural disturbances that have altered the state factors of coastal vegetation development. The change in hydraulic gradient associated with sea level rise and the increased risk of windfall in saturated soils are precluding the persistence of the near-shore community while also accelerating its decline (Paldor *et al.* 2022). These structural disturbances would change the management options, precluding reestablishment of ecological communities in their former locations, but allowing community shifts were adjacent environments conserved and left dynamic.

Case Study 3: Paleo and present climate change effects on the permafrost zone.

The permafrost zone in polar regions provides a useful example of response to perturbation because of the dramatic climatic changes it has experienced over the past 30,000 years and its importance to Earth's climate over the next several centuries (Lindgren *et al.* 2018; Finger & Rekvig 2022; Schuur *et al.* 2022). The terrestrial and subsea permafrost regions contain nearly 3,000 Gt of organic carbon, more than the sum of all other soil, the atmosphere, living biomass, and cumulative human emissions since the Industrial Revolution (Bar-On *et al.* 2018; Abbott *et al.* 2019a; Abbott 2022). These massive stocks of organic matter have been described as climate-protected, as they have been stabilized by persistent cold and wet conditions, which limit microbial and abiotic decomposition (Ernakovich *et al.* 2022; Schuur *et al.* 2022). Gradual climate warming after the Last Glacial Maximum (LGM), some 26,500 years ago, resulted in over 100 meters of sea level rise, retreat of ice sheets, and widespread development of lakes and peatlands (Lindgren *et al.* 2018; Sayedi *et al.* 2020). These enormous reorganizations were archetypal structural disturbances that altered land-water linkages, long-term carbon and nutrient balance, and distribution of vegetation. These changes created a state of net carbon uptake over large portions of Arctic Tundra and Boreal Forest, which has only recently been forced into carbon release because of anthropogenic climate change (Hayes *et al.* 2011; Turetsky *et al.* 2020; Schuur *et al.* 2022).

Across high-latitude and high-elevation ecosystems, local ecosystem structure modulated to effects of the gradual climate press that caused the transition from the Pleistocene to the Holocene. Organic soil

horizons and vegetation strongly influence the exchange of heat between the atmosphere and the soil, creating up to 12°C of difference between mean annual soil temperature relative to the overlying air (Shur & Jorgenson 2007). The development of soil and vegetation protected many Pleistocene permafrost deposits, imparting thermal resistance that effectively arrested—or at least delayed—the deglaciation process (Shur & Jorgenson 2007; Kokelj *et al.* 2017; Loranty *et al.* 2018; Strauss *et al.* 2022).

Ongoing anthropogenic warming is much more abrupt than the relatively gradual glacial-interglacial transition (Bova *et al.* 2021; Cheng *et al.* 2022), particularly in the permafrost zone, which is warming 3- to 6-times faster than the global mean (Abbott 2022; Abbott *et al.* 2022). This increased amplitude of climatic disturbance (Fig. 4) has surpassed the protective resistance of Holocene-aged soils and vegetation, triggering abrupt thaw and surface collapse in many of the regions with highest carbon densities (Olefeldt *et al.* 2016; Turetsky *et al.* 2020). Additionally, rapid warming is altering permafrost disturbance regimes. Functional disturbances such as wildfire are becoming more common and widespread (Mack *et al.* 2011), accelerating the structural disturbance of permafrost collapse, which together affect long-term carbon, nutrient, and water balance (Larouche *et al.* 2015; Moskovchenko *et al.* 2020; Rodríguez-Cardona *et al.* 2020; Abbott *et al.* 2021b). More acutely, the destabilization of permafrost soils, coastlines, and shorelines is profoundly impacting marine and terrestrial wildlife and the diverse human cultures of the permafrost zone (Chapin *et al.* 2013; Bronen *et al.* 2020; Abbott *et al.* 2022).

This case study demonstrates the interactions between the local structure of the critical zone and global climate change. Perhaps more importantly, it highlights some of the difficulties of creating Earth stewardship when the causes and consequences of environmental degradation are highly separated in space and time. Greenhouse gas emissions from outside of the permafrost zone are eroding resistance and recovery of permafrost ecosystems, including human villages and transportation infrastructure at circumpolar scales (ICC 2022). Communities in the permafrost zone have been innovative in adaptation and local mitigation (Chapin *et al.* 2013; Bronen *et al.* 2020; Abbott *et al.* 2022). At the same time, many community members are using intergovernmental forums such as the Arctic Council and Inuit Circumpolar Council to increase climate mitigation commitments to address the source of the problem: burning of fossil fuels (Johnson 2010; Kristoffersen & Langhelle 2017; Arctic Council 2022; ICCI 2022). This shows the intersection of local

community stewardship and global environmental governance, both of which are needed to resolve environmental injustice in the Anthropocene (Errigo *et al.* 2020; Webber *et al.* 2021; Chapin *et al.* 2022).

Case study 4: Hydrochemical recovery from acidification in the stormier present

Critical zone structure in watersheds in eastern North America and central Europe has been impacted by multiple changes to disturbance regimes over the past century. Terrestrial and aquatic ecosystems were subjected to decades of atmospheric acid deposition, which led to reduced soil pH and base cation loss from soils (Likens & Bormann 1974; Wettstad 2018). Environmental legislation on both continents reduced acid deposition starting in the 1980s, creating a natural experiment of recovery for watersheds with diverse critical zone structures (Likens 2013; Daniels *et al.* 2020; Hannah *et al.* 2022). In the decades since, many watersheds have seen streamwater dissolved organic carbon (DOC) and phosphorus concentrations increase (Evans *et al.* 2005; Kopáček *et al.* 2015), while streamwater inorganic nitrogen concentrations have decreased (Driscoll *et al.* 2003). Many studies have explored the mechanisms that may explain these temporal patterns, invoking various explanations including reduced mineralization under low soil pH, stabilization of soil aggregates at high ionic strengths/low soil pHs, and reduced vegetation uptake as a result of base cation limitation (Rosi-Marshall *et al.* 2016; Armfield *et al.* 2019; Cincotta *et al.* 2019).

Concurrently, these regions have been experiencing an increasing frequency of extreme hydrologic events. Large precipitation events have been linked to substantial flushing and export of carbon and nitrogen, thus comprising the majority of annual export in some watersheds (Raymond *et al.* 2016; Zarnetske *et al.* 2018; Kincaid *et al.* 2020). A recent study at Hubbard Brook Experimental Forest suggested that recovery from acidification and increasing frequency of extreme precipitation events interact in important ways, with greater stormflow nitrate export in an experimental watershed recovering from acidification (Marinos & Bernhardt 2018). This suggests that a multidimensional resilience approach is needed to understand the complex biogeochemical responses to acidification, recovery, and changing hydrologic regimes.



Figure 5. Examples of vulnerable and resilient approaches to human development in dynamic ecosystems. Each row shows how a different ecosystem structure responds to the functional disturbance of a flood. The first two rows were inspired by Delgado (2020).

While there are broad regional trends in these responses to reduced acid loading, there is a considerable degree of variability across individual catchments, likely associated with critical zone structure. For example, variability in DOC trends across catchments in New England depended on soil characteristics

and depth (Adler *et al.* 2021). Well-buffered, calcite-dominated watersheds are recovering faster than granitic watersheds with limited ability to buffer changes in soil pH. Differences in watershed topography and slope may lead to variability among watersheds in their hydrologic responsiveness to extreme events.

Rethinking the R's in the age of Big Data:

These case studies show how ecosystem structure and disturbance regimes interact to determine multidimensional resilience. To predict and prevent dangerous ecological state changes in the Anthropocene, we now need to dramatically advance our understanding of the nature of these interactions at global scales (Jiang *et al.* 2018; Turner *et al.* 2019). In many ecological contexts, resilience and resistance are viewed as mono-dimensional properties—e.g. collapse in a biological population or breakdown in an atmospheric or oceanic current (Liu *et al.* 2019; Steffen *et al.* 2018)—rather than as a nested, interacting system that intrinsically depends on the structure and state of the ecosystem. If resilience does indeed emerge from the ecosystem structure—the linkages across physical, biological, and social systems—this adds complexity but could also substantially increase predictive power (Gouveia *et al.* 2021). Indeed, we could be on the cusp of major breakthroughs in humanity's ability to quantitatively monitor and manage ecosystems for resilience. The availability of data from multiple observatories and monitoring networks at site to global scales (Leon *et al.* 2019; Brown *et al.* 2021; Ebeling *et al.* 2021; Heiner *et al.* 2022; Shogren *et al.* 2022) and the emergence of techniques that can analyze such voluminous and intricate data streams (Bergen *et al.* 2019) create an unprecedented opportunity to identify individual and interactive controls on ecosystem response to disturbance.

Until recently, characterizing multidimensional interactions at necessary spatiotemporal scales has been beyond the scope of disciplinary three- to five-year ecological projects (Abbott *et al.* 2016; Kolbe *et al.* 2019; Thomas *et al.* 2019). With the advent of new technology such as in situ sensors and remote sensing (e.g., lidar), we are amassing high volumes and a wide variety of observational data that can be used to test hypotheses about ecosystem response to disturbance regimes and associated water, carbon, and nutrient dynamics (Demchenko 2013). This big data revolution has had revolutionary effects across disciplines

(Alexander et al. 2015; Li et al. 2012) and is poised to transform ecosystem science as well (Reichstein et al. 2019). The recent emergence of new statistical and machine-learning algorithms has been driven, in part, by the advances in distributed computing and storage that accompany long-term monitoring, but more importantly, by the challenges in mining and analyzing these large, multi-scale, data-rich complex systems (Loehle 2006; Beven & Alcock 2012; Lum *et al.* 2013; Brunton *et al.* 2016).

Collectively, complex-systems tools comprise a variety of approaches including machine-learning algorithms, nonparametric statistics, network analysis, Bayesian inference, stochastic models, and evolutionary computation (Marçais & de Dreuzay 2017; Underwood *et al.* 2017; Shen *et al.* 2018; Frei *et al.* 2021). They can be used for classification, regression, and prediction tasks in the analysis of ecological dynamics across scales. A subset of machine-learning algorithms called ‘deep learning’ shows promise for advances in classification, anomaly detection, regression and prediction, where state variables are spatiotemporally dependent (Reichstein et al. 2019)—the default assumption for coevolving ecosystem structures and disturbance regimes (Thomas *et al.* 2016; Abbott *et al.* 2018a; Adler 2019). Deep learning models have gained rapid adoption in certain fields such as hydrology where long short-term memory (LSTM) models have eclipsed the performance of existing physics-based models in certain tasks (e.g., rainfall-runoff modeling) and are now being explored for their ability to capture hydrological concepts (Kratzert *et al.* 2019; Jiang *et al.* 2022; Lees *et al.* 2022). Three-dimensional convolutional neural networks have enhanced lidar-based forest inventories by spatially resolving individual tree crowns and distinguishing needle-leaf trees from deciduous (Ayrey & Hayes 2018). Image-based deep learning models have also been used for classification and interpretation of water quality dynamics such as with storm event suspended sediment transport (Hamshaw *et al.* 2018).

These tools are simultaneously revolutionizing the acquisition, cleaning, and analysis of multivariate ecological data (Hamshaw *et al.* 2018; Underwood *et al.* 2021; Wu *et al.* 2022). We can apply complex-systems tools to draw inferences from both terrestrial and aquatic signals of high temporal and spatial resolution (e.g., lidar first returns, time series of rainfall-runoff patterns or concentration discharge monitoring data) that serve as integrators of ecosystem dynamics, and have the potential to reflect the large-scale impacts of disturbances on the Earth system as a whole. For example, machine-learning algorithms are increasingly being used to

learn patterns from data for both clustering (i.e., unsupervised) and classification (i.e., supervised) tasks (Bergen et al. 2019). Unsupervised neural networks such as Self-Organizing Maps have been used to cluster catchments with similar combinations of multi-variate catchment attributes (Underwood 2017). Supervised methods, including nearest-neighbor and ‘random forests’ imputation methods, have been applied to model forest structural parameters including biomass and total timber volume using predictor variables generated from lidar data or orthoimagery (Latifi et al. 2010). Supervised methods are especially useful for cases such as this where manual classification would be too time-intensive, but can also be used to learn something about the multivariate feature interactions that manifest in an outward class or condition (Underwood et al. 2021).

In addition to the technical advances, this complex data revolution is accelerating conceptual crosspollination and opening doors to new collaborations among traditional ecological knowledge holders, researchers, and managers (Kimmerer 2002; Shen *et al.* 2018; Sayedi *et al.* 2020). Even terminology from the study of dynamical systems is helpful when describing ecosystem state and development. Attractors or basins of attraction are self-organizing or favored system configurations, and alternative stable states or multistability is the existence of multiple possible resilient ecosystem configurations (Dudkowski *et al.* 2016). Structural disturbances can erode resilience by creating alternative attractors that alter the recovery trajectory or reducing the resistance of the original ecosystem state (Fig. 2). The flexibility and power of complex system tools have only begun to be tapped. We think that major breakthroughs will occur as collaborations increase among Earth system scientists and local knowledge holders with deep intuitive and quantitative understanding of their systems, managers who know the pressing ecological questions and challenges, geospatial analysts who can collect massive amounts of remotely-sensed data, scientific instrument engineers who can facilitate direct measurements, and data scientists who can manage and implement data workflows, and finally control theorists and complex systems scientists who can help with interpretation and application.

People as a positive part of the ecosystem concept

Reminding researchers and readers not to forget people may sound ludicrous. Most of us are working on global environmental change, constantly engrossed in the causes and consequences of human

alteration of the Earth system. However, ecosystem ecology, hydrogeology, and many fields central to critical zone science tend to exclude humans implicitly and explicitly, often focusing on reference watersheds with no direct human influence or using “natural” conditions prior to the Anthropocene as a baseline (Chorover *et al.* 2007; Fandel *et al.* 2018; Abbott *et al.* 2019b; Ellis *et al.* 2021). Indeed, our focus on problems created by humanity can lead to bias against modified ecosystems despite their prevalence and indispensability in creating a sustainable global community (Hagerhall *et al.* 2004; Abbott *et al.* 2019b; Blaszcak *et al.* 2019; Elmqvist *et al.* 2019; Hill *et al.* 2022). Likewise, academic researchers and natural resource managers sometimes view environmental solutions as technical interventions to be imposed on communities rather than a tool for cultivating long-term relationship and cultural change (Chapin *et al.* 2022). In an ideal world, we would think in terms of communities and watersheds rather than administrative management units and environmental policies. There are compelling practical and ethical reasons for including human dimensions of ecosystems on both sides of resilience, i.e., when characterizing disturbance and considering the response. The social solidarity and respect we need to face intensifying ecological crises in the Anthropocene are unlikely in an environment of disciplinary dismissal and divisiveness (Allaire 2016; Abbott *et al.* 2018b; Webber *et al.* 2021).

Meaningful predictions and successful management depend on fully integrating human cultural and social dynamics into our conceptualization of ecosystems (Budds *et al.* 2014; Linton 2014; Abbott *et al.* 2021a; Chapin *et al.* 2022). While consideration of the human dimensions of ecosystems is necessary from a harm reduction perspective, it is arguably more important for the establishment of pro-environmental norms, policies, and individual behaviors (Behailu *et al.* 2016; Schuster *et al.* 2019). Examples of positive human-environment interactions are needed as models and motivators to accelerate cultural change (Kimmerer 2002; Palmer & Stewart 2020; Locke *et al.* 2021; Ansari & Landin 2022; Chapin *et al.* 2022).

Conclusions

We conclude that conceptual and practical rapprochement of human culture and the ecosystems we are a part of can enhance ecological resilience. Specifically, meaningful relationships with and affection for

our local environment can lead to sustainable norms, policies, and behaviors that humanity and the Earth system as a whole need urgently. We conclude that resilience emerges from the ecosystem structure—the linkages across physical and biological systems, especially human society. Finally, we recommend modeling human infrastructure and development patterns on natural disturbance regimes. Maximizing resistance is not a reliable strategy for maintaining ecosystem function, including ecosystem services, in the Anthropocene. Instead, we need connected and expansive habitat, disturbance regimes that are as natural and unregulated as possible, and complete and redundant biological communities, including all dimensions of human diversity. While creating and sharing an ethic of Earth stewardship is a multi-generational project, thankfully, we are not starting from zero. There are threads of stewardship and sustainability in every human culture and our species likely has an evolutionary penchant for environmental connection and care. It is our task to emphasize and cultivate these precious legacies.

Acknowledgments

This research was funded by the US National Science Foundation (grant numbers EAR-2012123, EAR-2011439, 2012188, 2011346, and 2012080). We thank Terry Chapin for input on an early version of the manuscript.

Data Availability Statement

This manuscript did not use any new data.

References

- Abatzoglou, J.T. & Williams, A.P. (2016). Impact of anthropogenic climate change on wildfire across western US forests. *PNAS*, 113, 11770–11775.
- Abbott, B.W. (2022). Permafrost Climate Feedbacks. In: *Global Arctic: An Introduction to the Multifaceted Dynamics of the Arctic* (eds. Finger, M. & Rekvig, G.). Springer International Publishing, Cham, pp. 189–209.

Abbott, B.W., Baranov, V., Mendoza-Lera, C., Nikolakopoulou, M., Harjung, A., Kolbe, T., *et al.* (2016). Using multi-tracer inference to move beyond single-catchment ecohydrology. *Earth-Science Reviews*, 160, 19–42.

Abbott, B.W., Bishop, K., Zarnetske, J.P., Hannah, D.M., Frei, R.J., Minaudo, C., *et al.* (2019a). A water cycle for the Anthropocene. *Hydrological Processes*, 33, 3046–3052.

Abbott, B.W., Bishop, K., Zarnetske, J.P., Minaudo, C., Chapin, F.S., Krause, S., *et al.* (2019b). Human domination of the global water cycle absent from depictions and perceptions. *Nature Geoscience*, 12, 533–540.

Abbott, B.W., Brown, M., Carey, J.C., Ernakovich, J., Frederick, J.M., Guo, L., *et al.* (2022). We Must Stop Fossil Fuel Emissions to Protect Permafrost Ecosystems. *Frontiers in Environmental Science*, 10.

Abbott, B.W., Errigo, I.M., Follett, A., Lawson, G., Meyer, M.M., Moon, H., *et al.* (2021a). *Getting to know the Utah Lake ecosystem*. Provo, Utah.

Abbott, B.W., Gruau, G., Zarnetske, J.P., Moatar, F., Barbe, L., Thomas, Z., *et al.* (2018a). Unexpected spatial stability of water chemistry in headwater stream networks. *Ecology Letters*, 21, 296–308.

Abbott, B.W., Moatar, F., Gauthier, O., Fovet, O., Antoine, V. & Ragueneau, O. (2018b). Trends and seasonality of river nutrients in agricultural catchments: 18 years of weekly citizen science in France. *Science of The Total Environment*, 624, 845–858.

Abbott, B.W., Rocha, A.V., Shogren, A., Zarnetske, J.P., Iannucci, F., Bowden, W.B., *et al.* (2021b). Tundra wildfire triggers sustained lateral nutrient loss in Alaskan Arctic. *Global Change Biology*, 27, 1408–1430.

Adler, R.W. (2019). Coevolution of Law and Science. *1*, 44, 1–66.

Adler, T., Underwood, K.L., Rizzo, D.M., Harpold, A., Sterle, G., Li, L., *et al.* (2021). Drivers of Dissolved Organic Carbon Mobilization From Forested Headwater Catchments: A Multi Scaled Approach. *Frontiers in Water*, 3.

Allaire, M.C. (2016). Using practical and social information to influence flood adaptation behavior: USING INFORMATION TO INFLUENCE FLOOD ADAPTATION BEHAVIOR. *Water Resour. Res.*, 52, 6078–6093.

Allison, G. (2004). The Influence of Species Diversity and Stress Intensity on Community Resistance and Resilience. *Ecological Monographs*, 74, 117–134.

Anderies, J., Folke, C., Walker, B. & Ostrom, E. (2013). Aligning Key Concepts for Global Change Policy: Robustness, Resilience, and Sustainability. *Ecology and Society*, 18.

Ansari, R.A. & Landin, J.M. (2022). Coverage of climate change in introductory biology textbooks, 1970–2019. *PLOS ONE*, 17, e0278532.

Arctic Council. (2022). *The Arctic Council*. Arctic Council. Available at: <https://arctic-council.org/>. Last accessed 20 February 2022.

Armfield, J.R., Perdrial, J.N., Gagnon, A., Ehrenkranz, J., Perdrial, N., Cincotta, M., *et al.* (2019). Does Stream Water Composition at Sleepers River in Vermont Reflect Dynamic Changes in Soils During Recovery From Acidification? *Front. Earth Sci.*, 6.

Armstrong McKay, D.I., Staal, A., Abrams, J.F., Winkelmann, R., Sakschewski, B., Loriani, S., *et al.* (2022). Exceeding 1.5°C global warming could trigger multiple climate tipping points. *Science*, 377, eabn7950.

Atkins, J.W., Bond-Lamberty, B., Fahey, R.T., Haber, L.T., Stuart-Haëntjens, E., Hardiman, B.S., *et al.* (2020). Application of multidimensional structural characterization to detect and describe moderate forest disturbance. *Ecosphere*, 11, e03156.

Ayrey, E. & Hayes, D.J. (2018). The Use of Three-Dimensional Convolutional Neural Networks to Interpret LiDAR for Forest Inventory. *Remote Sensing*, 10, 649.

Bank, M.S. & Hansson, S.V. (2019). The Plastic Cycle: A Novel and Holistic Paradigm for the Anthropocene. *Environ. Sci. Technol.*, 53, 7177–7179.

Barbe, L., Mony, C. & Abbott, B.W. (2020). Artificial Intelligence Accidentally Learned Ecology through Video Games. *Trends in Ecology & Evolution*, S0169534720301105.

Bar-On, Y.M., Phillips, R. & Milo, R. (2018). The biomass distribution on Earth. *Proc Natl Acad Sci USA*, 115, 6506–6511.

Behailu, B.M., Pietilä, P.E. & Katko, T.S. (2016). Indigenous Practices of Water Management for Sustainable Services: Case of Borana and Konso, Ethiopia. *SAGE Open*, 6, 2158244016682292.

Benson, M. (2012). Intelligent Tinkering: the Endangered Species Act and Resilience. *Ecology and Society*, 17.

570 Bergstrom, D.M., Wienecke, B.C., van den Hoff, J., Hughes, L., Lindenmayer, D.B., Ainsworth, T.D., *et al.*
 571 (2021). Combating ecosystem collapse from the tropics to the Antarctic. *Global Change Biology*, 27,
 572 1692–1703.
 573 Bernhardt, E.S. & Palmer, M.A. (2007). Restoring streams in an urbanizing world. *Freshwater Biology*, 52, 738–
 574 751.
 575 Beven, K.J. & Alcock, R.E. (2012). Modelling everything everywhere: a new approach to decision-making for
 576 water management under uncertainty. *Freshwater Biology*, 57, 124–132.
 577 Bishop, K., Beven, K., Destouni, G., Abrahamsson, K., Andersson, L., Johnson, R.K., *et al.* (2009). Nature as
 578 the “Natural” Goal for Water Management: A Conversation. *Ambio*, 38, 209–214.
 579 Blaszcak, J.R., Delesantro, J.M., Urban, D.L., Doyle, M.W. & Bernhardt, E.S. (2019). Scoured or suffocated:
 580 Urban stream ecosystems oscillate between hydrologic and dissolved oxygen extremes. *Limnol*
 581 *Oceanogr.*, 64, 877–894.
 582 Bochet, O., Bethencourt, L., Dufresne, A., Farasin, J., Pédrot, M., Labasque, T., *et al.* (2020). Iron-oxidizer
 583 hotspots formed by intermittent oxic–anoxic fluid mixing in fractured rocks. *Nat. Geosci.*, 1–7.
 584 Boers, N. & Rypdal, M. (2021). Critical slowing down suggests that the western Greenland Ice Sheet is close
 585 to a tipping point. *PNAS*, 118.
 586 Bova, S., Rosenthal, Y., Liu, Z., Godad, S.P. & Yan, M. (2021). Seasonal origin of the thermal maxima at the
 587 Holocene and the last interglacial. *Nature*, 589, 548–553.
 588 Bradshaw, C.J.A., Ehrlich, P.R., Beattie, A., Ceballos, G., Crist, E., Diamond, J., *et al.* (2021). Underestimating
 589 the Challenges of Avoiding a Ghastly Future. *Front. Conserv. Sci.*, 1.
 590 Bronen, R., Pollock, D., Overbeck, J., Stevens, D., Natali, S. & Maio, C. (2020). Usteq: integrating indigenous
 591 knowledge and social and physical sciences to coproduce knowledge and support community-based
 592 adaptation. *Polar Geography*, 43, 188–205.
 593 Brown, B., Fulterton, A., Kopp, D., Tromboni, F., Shogren, A., Webb, J., *et al.* (2021). Streamflow Metrics
 594 and Catchment Characteristics for Global Streamflow Dataset.
 595 Brunton, S.L., Budišić, M., Kaiser, E. & Kutz, J.N. (2022). Modern Koopman Theory for Dynamical Systems.
 596 *SIAM Rev.*, 64, 229–340.
 597 Brunton, S.L., Proctor, J.L. & Kutz, J.N. (2016). Discovering governing equations from data by sparse
 598 identification of nonlinear dynamical systems. *Proceedings of the National Academy of Sciences*, 113, 3932–
 599 3937.
 600 Budds, J., Linton, J. & McDonnell, R. (2014). The hydrosocial cycle. *Geoforum*, 167–169.
 601 Callicott, J.B. & Mumford, K. (1997). Ecological Sustainability as a Conservation Concept. *Conservation Biology*,
 602 11, 32–40.
 603 Carpenter, S., Walker, B., Anderies, J.M. & Abel, N. (2001). From Metaphor to Measurement: Resilience of
 604 What to What? *Ecosystems*, 4, 765–781.
 605 Carpenter, S.R., Arani, B.M.S., Hanson, P.C., Scheffer, M., Stanley, E.H. & Nes, E.V. (2020). Stochastic
 606 dynamics of Cyanobacteria in long-term high-frequency observations of a eutrophic lake. *Limnology*
 607 *and Oceanography Letters*.
 608 Cassidy, L., Perkins, J. & Bradley, J. (2022). Too much, too late: fires and reactive wildfire management in
 609 northern Botswana’s forests and woodland savannas. *African Journal of Range & Forage Science*, 39, 160–
 610 174.
 611 Ceballos, G., Ehrlich, P.R. & Raven, P.H. (2020). Vertebrates on the brink as indicators of biological
 612 annihilation and the sixth mass extinction. *PNAS*, 117, 13596–13602.
 613 Chapin, F.S., Matson, P.A. & Vitousek, P.M. (2012). *Principles of Terrestrial Ecosystem Ecology*. Springer New
 614 York, New York, NY.
 615 Chapin, F.S., Robards, M.D., Johnstone, J.F., Lantz, T.C. & Kokelj, S.V. (2013). Case Study: Novel Socio-
 616 Ecological Systems in the North: Potential Pathways Toward Ecological and Societal Resilience. In:
 617 *Novel Ecosystems*. John Wiley & Sons, Ltd, pp. 334–344.
 618 Chapin, F.S., Walker, L.R., Fastie, C.L. & Sharman, L.C. (1994). Mechanisms of Primary Succession
 619 Following Deglaciation at Glacier Bay, Alaska. *Ecological Monographs*, 64, 149–175.
 620 Chapin, F.S., Weber, E.U., Bennett, E.M., Biggs, R., van den Bergh, J., Adger, W.N., *et al.* (2022). Earth
 621 stewardship: Shaping a sustainable future through interacting policy and norm shifts. *Ambio*.

- Cheng, F., Garzione, C., Li, X., Salzmann, U., Schwarz, F., Haywood, A.M., *et al.* (2022). Alpine permafrost could account for a quarter of thawed carbon based on Plio-Pleistocene paleoclimate analogue. *Nat Commun*, 13, 1329.
- Chorover, J., Kretzschmar, R., Garcia-Pichel, F. & Sparks, D.L. (2007). Soil Biogeochemical Processes within the Critical Zone. *Elements*, 3, 321–326.
- Christianson, A. (2015). Social science research on Indigenous wildfire management in the 21st century and future research needs. *Int. J. Wildland Fire*, 24, 190–200.
- Cincotta, M.M., Perdrial, J.N., Shavitz, A., Libenson, A., Landsman-Gerjoi, M., Perdrial, N., *et al.* (2019). Soil Aggregates as a Source of Dissolved Organic Carbon to Streams: An Experimental Study on the Effect of Solution Chemistry on Water Extractable Carbon. *Frontiers in Environmental Science*, 7.
- Collins, S.E., Matter, S.F., Buffam, I. & Flotemersch, J.E. (2018). A patchy continuum? Stream processes show varied responses to patch- and continuum-based analyses. *Ecosphere*, 9, e02481.
- Cooper, A.H., Brown, T.J., Price, S.J., Ford, J.R. & Waters, C.N. (2018). Humans are the most significant global geomorphological driving force of the 21st century. *The Anthropocene Review*, 5, 222–229.
- Crandall, T., Jones, E., Greenhalgh, M., Frei, R.J., Griffin, N., Severe, E., *et al.* (2021). Megafire affects stream sediment flux and dissolved organic matter reactivity, but land use dominates nutrient dynamics in semiarid watersheds. *PLOS ONE*, 16, e0257733.
- Daniels, B., Follett, A. & Davis, J. (2020). *The Making of the Clean Air Act*. Hastings L. J.
- Davis, D., Knuepfer, P.L.K., Miller, N. & Vian, M. (2009). *Fluvial Geomorphology of the Upper Esopus Creek Watershed and Implications for Stream Management*. New York State Geological Society, New York, N.Y., U.S.A.
- Dearman, T.L. & James, L.A. (2019). Patterns of legacy sediment deposits in a small South Carolina Piedmont catchment, USA. *Geomorphology*, 343, 1–14.
- Delgado, J.A.S. (2020). A várzea e as enchentes. *Confins. Revue franco-brésilienne de géographie / Revista franco-brasileira de geografia*.
- Díaz, S., Settele, J., Brondízio, E.S., Ngo, H.T., Agard, J., Arneth, A., *et al.* (2019). Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science*, 366.
- Diffenbaugh, N.S. & Field, C.B. (2013). Changes in Ecologically Critical Terrestrial Climate Conditions. *Science*, 341, 486–492.
- Driscoll, C.T., Whitall, D., Aber, J., Boyer, E., Castro, M., Cronan, C., *et al.* (2003). Nitrogen Pollution in the Northeastern United States: Sources, Effects, and Management Options. *BioScience*, 53, 357.
- Dudkowski, D., Jafari, S., Kapitaniak, T., Kuznetsov, N.V., Leonov, G.A. & Prasad, A. (2016). Hidden attractors in dynamical systems. *Physics Reports*, Hidden Attractors in Dynamical Systems, 637, 1–50.
- Dupas, R., Minaudo, C. & Abbott, B.W. (2019). Stability of spatial patterns in water chemistry across temperate ecoregions. *Environ. Res. Lett.*, 14, 074015.
- Ebeling, P., Dupas, R., Abbott, B.W., Kumar, R., Ehrhardt, S., Fleckenstein, J.H., *et al.* (2021). Long-Term Nitrate Trajectories Vary by Season in Western European Catchments. *Global Biogeochemical Cycles*, 35, e2021GB007050.
- Elhacham, E., Ben-Uri, L., Grozovski, J., Bar-On, Y.M. & Milo, R. (2020). Global human-made mass exceeds all living biomass. *Nature*, 588, 442–444.
- Ellis, E.C. (2021). Land Use and Ecological Change: A 12,000-Year History. *Annual Review of Environment and Resources*, 46, 1–33.
- Ellis, E.C., Gauthier, N., Goldewijk, K.K., Bird, R.B., Boivin, N., Díaz, S., *et al.* (2021). People have shaped most of terrestrial nature for at least 12,000 years. *PNAS*, 118.
- Elmqvist, T., Andersson, E., Frantzeskaki, N., McPhearson, T., Olsson, P., Gaffney, O., *et al.* (2019). Sustainability and resilience for transformation in the urban century. *Nat Sustain*, 2, 267–273.
- Ernakovich, J.G., Barbato, R.A., Rich, V.I., Schädel, C., Hewitt, R.E., Doherty, S.J., *et al.* (2022). Microbiome assembly in thawing permafrost and its feedbacks to climate. *Global Change Biology*, 28, 5007–5026.
- Errigo, I.M., Abbott, B.W., Mendoza, D.L., Mitchell, L., Sayedi, S.S., Glenn, J., *et al.* (2020). Human Health and Economic Costs of Air Pollution in Utah: An Expert Assessment. *Atmosphere*, 11, 1238.
- Evans, C.D., Monteith, D.T. & Cooper, D.M. (2005). Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution*, Recovery

- from acidification in the UK: Evidence from 15 years of acid waters monitoring Recovery from acidification in the UK: Evidence from 15 years of acid waters monitoring, 137, 55–71.
- Fagherazzi, S., Nordio, G., Munz, K., Catucci, D. & Kearney, W.S. (2019). Variations in Persistence and Regenerative Zones in Coastal Forests Triggered by Sea Level Rise and Storms. *Remote Sensing*, 11, 2019.
- Fandel, C.A., Breshears, D.D. & McMahon, E.E. (2018). Implicit assumptions of conceptual diagrams in environmental science and best practices for their illustration. *Ecosphere*, 9, 1–15.
- Fang, Y. & Jawitz, J.W. (2019). The evolution of human population distance to water in the USA from 1790 to 2010. *Nature Communications*, 10, 430.
- Finger, M. & Rekvig, G. (2022). *Global Arctic: An Introduction to the Multifaceted Dynamics of the Arctic*. 1st edn. Springer International Publishing.
- Fisher, S.G., Grimm, N.B., Martí, E., Holmes, R.M. & Jones Jr, J.B. (1998). Material spiraling in stream corridors: a telescoping ecosystem model. *Ecosystems*, 1, 19–34.
- Florinsky, I.V. (2012). The Dokuchaev hypothesis as a basis for predictive digital soil mapping (on the 125th anniversary of its publication). *Eurasian Soil Sc.*, 45, 445–451.
- Folke, C., Biggs, R., Norström, A.V., Reyers, B. & Rockström, J. (2016). Social-ecological resilience and biosphere-based sustainability science. *Ecology and Society*, 21.
- Folke, C., Polasky, S., Rockström, J., Galaz, V., Westley, F., Lamont, M., *et al.* (2021). Our future in the Anthropocene biosphere. *Ambio*, 50, 834–869.
- Fowler, H.J., Lenderink, G., Prein, A.F., Westra, S., Allan, R.P., Ban, N., *et al.* (2021). Anthropogenic intensification of short-duration rainfall extremes. *Nature Reviews Earth & Environment*, 2, 107–122.
- Frei, R.J., Abbott, B.W., Dupas, R., Gu, S., Gruau, G., Thomas, Z., *et al.* (2020). Predicting Nutrient Incontinence in the Anthropocene at Watershed Scales. *Front. Environ. Sci.*, 7.
- Frei, R.J., Lawson, G.M., Norris, A.J., Cano, G., Vargas, M.C., Kujanpää, E., *et al.* (2021). Limited progress in nutrient pollution in the U.S. caused by spatially persistent nutrient sources. *PLOS ONE*, 16, e0258952.
- Fricke, E.C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C.D., Sanisidro, O., *et al.* (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene. *Science*, 377, 1008–1011.
- Fuller, I.C., Gilvear, D.J., Thoms, M.C. & Death, R.G. (2019). Framing resilience for river geomorphology: Reinventing the wheel? *River Research and Applications*, 35, 91–106.
- Fuller, R., Landrigan, P.J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., *et al.* (2022). Pollution and health: a progress update. *The Lancet Planetary Health*, 0.
- Glasby, T.M. & Underwood, A.J. (1996). Sampling to differentiate between pulse and press perturbations. *Environ Monit Assess*, 42, 241–252.
- Gouveia, C., Mórèh, Á. & Jordán, F. (2021). Combining centrality indices: Maximizing the predictability of keystone species in food webs. *Ecological Indicators*, 126, 107617.
- Hagerhall, C.M., Purcell, T. & Taylor, R. (2004). Fractal dimension of landscape silhouette outlines as a predictor of landscape preference. *Journal of Environmental Psychology*, 24, 247–255.
- Hamshaw, S.D., Dewoolkar, M.M., Schroth, A.W., Wemple, B.C. & Rizzo, D.M. (2018). A New Machine-Learning Approach for Classifying Hysteresis in Suspended-Sediment Discharge Relationships Using High-Frequency Monitoring Data. *Water Resources Research*, 54, 4040–4058.
- Hannah, D.M., Abbott, B.W., Khamis, K., Kelleher, C., Lynch, I., Krause, S., *et al.* (2022). Illuminating the ‘invisible water crisis’ to address global water pollution challenges. *Hydrological Processes*, 36, e14525.
- Hayes, D.J., McGuire, A.D., Kicklighter, D.W., Gurney, K.R., Burnside, T.J. & Melillo, J.M. (2011). Is the northern high-latitude land-based CO₂ sink weakening? *Global Biogeochemical Cycles*, 25, n/a–n/a.
- Heddam, S., Kim, S., Danandeh Mehr, A., Zounemat-Kermani, M., Malik, A., Elbeltagi, A., *et al.* (2022). Chapter 1 - Predicting dissolved oxygen concentration in river using new advanced machines learning: Long-short term memory (LSTM) deep learning. In: *Computers in Earth and Environmental Sciences* (ed. Pourghasemi, H.R.). Elsevier, pp. 1–20.
- Heiner, M., Heaton, M.J., Abbott, B.W., White, P., Minaudo, C. & Dupas, R. (2022). Model-Based Clustering of Trends and Cycles of Nitrate Concentrations in Rivers Across France. *JABES*.
- Heinze, C., Blenckner, T., Martins, H., Rusiecka, D., Döschner, R., Gehlen, M., *et al.* (2021). The quiet crossing of ocean tipping points. *PNAS*, 118.

- Hill, S.K., Hale, R.L., Grinath, J.B., Folk, B.T., Nielson, R. & Reinhardt, K. (2022). Looking beyond leaves: variation in nutrient leaching potential of seasonal litterfall among different species within an urban forest. *Urban Ecosyst*, 25, 1097–1109.
- Houlton, B.Z., Almaraz, M., Aneja, V., Austin, A.T., Bai, E., Cassman, K.G., *et al.* (2019). A World of Cobenefits: Solving the Global Nitrogen Challenge. *Earth's Future*, 0.
- Hurteau, M.D., Liang, S., Westerling, A.L. & Wiedinmyer, C. (2019). Vegetation-fire feedback reduces projected area burned under climate change. *Sci Rep*, 9, 2838.
- ICC. (2022). *Inuit Circumpolar Council – United Voice of the Arctic*. Available at: <https://www.inuitcircumpolar.com/>. Last accessed 20 February 2022.
- ICCI. (2022). State of the Cryosphere Report 2022 – ICCI – International Cryosphere Climate Initiative.
- IPCC. (2021). *IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group 1 to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Jasinski, J.P.P. & Payette, S. (2005). The Creation of Alternative Stable States in the Southern Boreal Forest, Québec, Canada. *Ecological Monographs*, 75, 561–583.
- Jenny, H. (1941). *Factors of soil formation: a system of quantitative pedology*. Dover, New York.
- Jiang, J., Tang, S., Han, D., Fu, G., Solomatine, D. & Zheng, Y. (2020). A comprehensive review on the design and optimization of surface water quality monitoring networks. *Environmental Modelling & Software*, 132, 104792.
- Jiang, S., Zheng, Y., Wang, C. & Babovic, V. (2022). Uncovering Flooding Mechanisms Across the Contiguous United States Through Interpretive Deep Learning on Representative Catchments. *Water Resources Research*, 58, e2021WR030185.
- Johnson, K.M., Snyder, N.P., Castle, S., Hopkins, A.J., Waltner, M., Merritts, D.J., *et al.* (2019). Legacy sediment storage in New England river valleys: Anthropogenic processes in a postglacial landscape. *Geomorphology*, 327, 417–437.
- Johnson, L. (2010). The Fearful Symmetry of Arctic Climate Change: Accumulation by Degradation. *Environ Plan D*, 28, 828–847.
- Kearney, W.S., Fernandes, A. & Fagherazzi, S. (2019). Sea-level rise and storm surges structure coastal forests into persistence and regeneration niches. *PLOS ONE*, 14, e0215977.
- Kemp, D.B., Eichenseer, K. & Kiessling, W. (2015). Maximum rates of climate change are systematically underestimated in the geological record. *Nat Commun*, 6, 8890.
- Keys, P.W., Galaz, V., Dyer, M., Matthews, N., Folke, C., Nyström, M., *et al.* (2019). Anthropocene risk. *Nat Sustain*, 2, 667–673.
- Kimmerer, R.W. (2002). Weaving Traditional Ecological Knowledge into Biological Education: A Call to Action. *BioScience*, 52, 432–438.
- Kincaid, D.W., Seybold, E.C., Adair, E.C., Bowden, W.B., Perdrial, J.N., Vaughan, M.C.H., *et al.* (2020). Land Use and Season Influence Event-Scale Nitrate and Soluble Reactive Phosphorus Exports and Export Stoichiometry from Headwater Catchments. *Water Resources Research*, 56, e2020WR027361.
- King, M.D., Howat, I.M., Candela, S.G., Noh, M.J., Jeong, S., Noël, B.P.Y., *et al.* (2020). Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat. *Communications Earth & Environment*, 1, 1–7.
- Kokelj, S.V., Lantz, T.C., Tunnicliffe, J., Segal, R. & Lacelle, D. (2017). Climate-driven thaw of permafrost preserved glacial landscapes, northwestern Canada. *Geology*, 45, 371–374.
- Kolbe, T., de Dreuz, J.-R., Abbott, B.W., Aquilina, L., Babey, T., Green, C.T., *et al.* (2019). Stratification of reactivity determines nitrate removal in groundwater. *Proceedings of the National Academy of Sciences*, 116, 2494–2499.
- Kopáček, J., Hejzlar, J., Kaňá, J., Norton, S.A. & Stuchlík, E. (2015). Effects of Acidic Deposition on in-Lake Phosphorus Availability: A Lesson from Lakes Recovering from Acidification. *Environ. Sci. Technol.*, 49, 2895–2903.
- Kratzert, F., Klotz, D., Herrnegger, M., Sampson, A.K., Hochreiter, S. & Nearing, G.S. (2019). Toward Improved Predictions in Ungauged Basins: Exploiting the Power of Machine Learning. *Water Resources Research*, 55, 11344–11354.

- Kristoffersen, B. & Langhelle, O. (2017). Sustainable Development as a Global-Arctic Matter: Imaginaries and Controversies. In: *Governing Arctic Change: Global Perspectives* (eds. Keil, K. & Knecht, S.). Palgrave Macmillan UK, London, pp. 21–41.
- Landrigan, P.J., Fuller, R., Acosta, N.J.R., Adeyi, O., Arnold, R., Basu, N. (Nil), *et al.* (2017). The Lancet Commission on pollution and health. *The Lancet*, 0.
- Larouche, J.R., Abbott, B.W., Bowden, W.B. & Jones, J.B. (2015). The role of watershed characteristics, permafrost thaw, and wildfire on dissolved organic carbon biodegradability and water chemistry in Arctic headwater streams. *Biogeosciences*, 12, 4221–4233.
- Larsson, M. & Abbott, B.W. (2018). Is the Capacity for Vocal Learning in Vertebrates Rooted in Fish Schooling Behavior? *Evol Biol*.
- Leavitt, W.M. & Kiefer, J.J. (2006). Infrastructure Interdependency and the Creation of a Normal Disaster: The Case of Hurricane Katrina and the City of New Orleans. *Public Works Management & Policy*, 10, 306–314.
- Lees, T., Reece, S., Kratzert, F., Klotz, D., Gauch, M., De Bruijn, J., *et al.* (2022). Hydrological concept formation inside long short-term memory (LSTM) networks. *Hydrology and Earth System Sciences*, 26, 3079–3101.
- Leon, M., Lubinski, D.J., Bode, C.A., Marini, L., Seul, M., Derry, L.A., *et al.* (2019). Increasing reusability of Critical Zone data with CUAHSI HydroShare and CZ Manager, 2019, IN23D-0903.
- Lewis, S.L. & Maslin, M.A. (2015). Defining the Anthropocene. *Nature*, 519, 171–180.
- Likens, G.E. (2013). The Hubbard Brook Ecosystem Study: Celebrating 50 Years. *The Bulletin of the Ecological Society of America*, 94, 336–337.
- Likens, G.E. & Bormann, F.H. (1974). Acid Rain: A Serious Regional Environmental Problem. *Science*, 184, 1176–1179.
- Lindeman, R.L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23, 399–417.
- Lindgren, A., Hugelius, G. & Kuhry, P. (2018). Extensive loss of past permafrost carbon but a net accumulation into present-day soils. *Nature*, 560, 219.
- Linton, J. (2014). Modern water and its discontents: a history of hydrosocial renewal. *WIREs Water*, 1, 111–120.
- Locke, H., Rockström, J., Bakker, P., Bapna, M., Gough, M., Hilty, J., *et al.* (2021). A Nature-Positive World: The Global Goal for Nature.
- Loehle, C. (2006). Control theory and the management of ecosystems. *Journal of Applied Ecology*, 43, 957–966.
- Loranty, M.M., Abbott, B.W., Blok, D., Douglas, T.A., Epstein, H.E., Forbes, B.C., *et al.* (2018). Reviews and syntheses: Changing ecosystem influences on soil thermal regimes in northern high-latitude permafrost regions. *Biogeosciences*, 15, 5287–5313.
- Lum, P.Y., Singh, G., Lehman, A., Ishkanov, T., Vejdemo-Johansson, M., Alagappan, M., *et al.* (2013). Extracting insights from the shape of complex data using topology. *Sci Rep*, 3, 1236.
- Mack, M.C., Bret-Harte, M.S., Hollingsworth, T.N., Jandt, R.R., Schuur, E.A.G., Shaver, G.R., *et al.* (2011). Carbon loss from an unprecedented Arctic tundra wildfire. *Nature*, 475, 489–492.
- Mack, M.C. & D’Antonio, C.M. (1998). Impacts of biological invasions on disturbance regimes. *Trends in Ecology & Evolution*, 13, 195–198.
- Manney, G.L., Santee, M.L., Rex, M., Livesey, N.J., Pitts, M.C., Veefkind, P., *et al.* (2011). Unprecedented Arctic ozone loss in 2011. *Nature*, 478, 469–475.
- Marçais, J. & de Dreuzy, J.-R. (2017). Prospective Interest of Deep Learning for Hydrological Inference. *Groundwater*, 55, 688–692.
- Marinos, R.E. & Bernhardt, E.S. (2018). Soil carbon losses due to higher pH offset vegetation gains due to calcium enrichment in an acid mitigation experiment. *Ecology*, 99, 2363–2373.
- Marlon, J.R. (2020). What the past can say about the present and future of fire. *Quaternary Research*, 96, 66–87.
- Meerow, S. & Newell, J.P. (2019). Urban resilience for whom, what, when, where, and why? *Urban Geography*, 40, 309–329.
- Meerow, S., Newell, J.P. & Stults, M. (2016). Defining urban resilience: A review. *Landscape and Urban Planning*, 147, 38–49.

- Mlynczak, M.G., Hunt, L.A., Garcia, R.R., Harvey, V.L., Marshall, B.T., Yue, J., *et al.* (2022). Cooling and Contraction of the Mesosphere and Lower Thermosphere From 2002 to 2021. *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036767.
- Mo, Y., Kearney, M.S. & Turner, R.E. (2020). The resilience of coastal marshes to hurricanes: The potential impact of excess nutrients. *Environment International*, 138, 105409.
- Moskovchenko, D.V., Arefev, S.P., Moskovchenko, M.D. & Yurtaev, A.A. (2020). Spatiotemporal Analysis of Wildfires in the Forest Tundra of Western Siberia. *Contemp. Probl. Ecol.*, 13, 193–203.
- Mu, C., Abbott, B.W., Norris, A.J., Mu, M., Fan, C., Chen, X., *et al.* (2020). The status and stability of permafrost carbon on the Tibetan Plateau. *Earth-Science Reviews*, 211, 103433.
- NASEM. (2020). *Review of the New York City Watershed Protection Program*. National Academies Press, Washington, D.C.
- National Research Council. (2001). *Basic Research Opportunities in Earth Science*. The National Academies Press, Washington, DC.
- Newman, T.P. (2017). Tracking the release of IPCC AR5 on Twitter: Users, comments, and sources following the release of the Working Group I Summary for Policymakers. *Public Underst Sci*, 26, 815–825.
- Nizzetto, L., Macleod, M., Borgå, K., Cabrerizo, A., Dachs, J., Guardo, A.D., *et al.* (2010). Past, Present, and Future Controls on Levels of Persistent Organic Pollutants in the Global Environment. *Environ. Sci. Technol.*, 44, 6526–6531.
- Noe, G.B., Cashman, M.J., Skalak, K., Gellis, A., Hopkins, K.G., Moyer, D., *et al.* (2020). Sediment dynamics and implications for management: State of the science from long-term research in the Chesapeake Bay watershed, USA. *WIREs Water*, 7, e1454.
- North, M.P. & Keeton, W.S. (2008). Emulating natural disturbance regimes: an emerging approach for sustainable forest management. In: *Patterns and processes in forest landscapes*. Springer, pp. 341–372.
- Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P., *et al.* (2016). Circumpolar distribution and carbon storage of thermokarst landscapes. *Nature Communications*, 7, 13043.
- Paldor, A., Stark, N., Florence, M., Raubenheimer, B., Elgar, S., Housego, R., *et al.* (2022). Coastal topography and hydrogeology control critical groundwater gradients and potential beach surface instability during storm surges. *Hydrology and Earth System Sciences*, 26, 5987–6002.
- Palmer, M.A. & Stewart, G.A. (2020). Ecosystem restoration is risky ... but we can change that. *One Earth*, 3, 661–664.
- Pinay, G., Bernal, S., Abbott, B.W., Lupon, A., Marti, E., Sabater, F., *et al.* (2018). Riparian Corridors: A New Conceptual Framework for Assessing Nitrogen Buffering Across Biomes. *Front. Environ. Sci.*, 6.
- Poff, N.L., Allan, J.D., Bain, M.B., Karr, J.R., Prestegard, K.L., Richter, B.D., *et al.* (1997). The Natural Flow Regime. *BioScience*, 47, 769–784.
- Raymond, P.A., Saiers, J.E. & Sobczak, W.V. (2016). Hydrological and biogeochemical controls on watershed dissolved organic matter transport: pulse-shunt concept. *Ecol*, 97, 5–16.
- Redman, C. (2014). Should sustainability and resilience be combined or remain distinct pursuits? *Ecology and Society*, 19.
- Reisner, M. (1993). *Cadillac Desert: The American West and Its Disappearing Water, Revised Edition*. 2nd edition. Penguin Books, New York, N.Y., U.S.A.
- Ritchie, P.D.L., Clarke, J.J., Cox, P.M. & Huntingford, C. (2021). Overshooting tipping point thresholds in a changing climate. *Nature*, 592, 517–523.
- Rockström, J., Beringer, T., Hole, D., Griscom, B., Mascia, M.B., Folke, C., *et al.* (2021). Opinion: We need biosphere stewardship that protects carbon sinks and builds resilience. *PNAS*, 118.
- Rode, M., Wade, A.J., Cohen, M.J., Hensley, R.T., Bowes, M.J., Kirchner, J.W., *et al.* (2016). Sensors in the Stream: The High-Frequency Wave of the Present. *Environ. Sci. Technol.*, 50, 10297–10307.
- Rodríguez-Cardona, B.M., Coble, A.A., Wymore, A.S., Kolosov, R., Podgorski, D.C., Zito, P., *et al.* (2020). Wildfires lead to decreased carbon and increased nitrogen concentrations in upland arctic streams. *Scientific Reports*, 10, 8722.
- Rogers, C.D.F., Bouch, C.J., Williams, S., Barber, A.R.G., Baker, C.J., Bryson, J.R., *et al.* (2012). Resistance and resilience – paradigms for critical local infrastructure. *Proceedings of the Institution of Civil Engineers - Municipal Engineer*, 165, 73–83.

882 Rosi-Marshall, E.J., Bernhardt, E.S., Buso, D.C., Driscoll, C.T. & Likens, G.E. (2016). Acid rain mitigation
883 experiment shifts a forested watershed from a net sink to a net source of nitrogen. *Proceedings of the*
884 *National Academy of Sciences*, 113, 7580–7583.

885 Sanders, B.F., Schubert, J.E., Kahl, D.T., Mach, K.J., Brady, D., AghaKouchak, A., *et al.* (2022). Large and
886 inequitable flood risks in Los Angeles, California. *Nat Sustain*, 1–11.

887 Sandifer, P.A., Sutton-Grier, A.E. & Ward, B.P. (2015). Exploring connections among nature, biodiversity,
888 ecosystem services, and human health and well-being: Opportunities to enhance health and
889 biodiversity conservation. *Ecosystem Services*, 12, 1–15.

890 Sayedi, S.S., Abbott, B.W., Thornton, B.F., Frederick, J.M., Vonk, J.E., Overduin, P., *et al.* (2020). Subsea
891 permafrost carbon stocks and climate change sensitivity estimated by expert assessment. *Environ. Res.*
892 *Lett.*, 15, 124075.

893 Scheffer, M., Bascompte, J., Brock, W.A., Brovkin, V., Carpenter, S.R., Dakos, V., *et al.* (2009). Early-warning
894 signals for critical transitions. *Nature*, 461, 53–59.

895 Schlesinger, W.H. & Bernhardt, E.S. (2020). *Biogeochemistry: An Analysis of Global Change*. 4th edition. Academic
896 Press, SanDiego.

897 Schoolmaster Jr., D.R., Zirbel, C.R. & Cronin, J.P. (2020). A graphical causal model for resolving species
898 identity effects and biodiversity–ecosystem function correlations. *Ecology*, 101, e03070.

899 Schuster, R., Germain, R.R., Bennett, J.R., Reo, N.J. & Arcese, P. (2019). Vertebrate biodiversity on
900 indigenous-managed lands in Australia, Brazil, and Canada equals that in protected areas.
901 *Environmental Science & Policy*, 101, 1–6.

902 Schuur, E.A.G., Abbott, B.W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., *et al.* (2022).
903 Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic. *Annual Review*
904 *of Environment and Resources*, 47, 343–371.

905 Shade, A., Peter, H., Allison, S., Baho, D., Berga, M., Buergermann, H., *et al.* (2012). Fundamentals of Microbial
906 Community Resistance and Resilience. *Frontiers in Microbiology*, 3.

907 Shen, C., Laloy, E., Elshorbagy, A., Albert, A., Bales, J., Chang, F.-J., *et al.* (2018). HESS Opinions: Incubating
908 deep-learning-powered hydrologic science advances as a community. *Hydrology and Earth System*
909 *Sciences*, 22, 5639–5656.

910 Shogren, A.J., Zarnetske, J.P., Abbott, B.W., Bratsman, S., Brown, B., Carey, M.P., *et al.* (2022). Multi-year,
911 spatially extensive, watershed-scale synoptic stream chemistry and water quality conditions for six
912 permafrost-underlain Arctic watersheds. *Earth System Science Data*, 14, 95–116.

913 Shur, Y.L. & Jorgenson, M.T. (2007). Patterns of permafrost formation and degradation in relation to climate
914 and ecosystems. *Permafrost and Periglacial Processes*, 18, 7–19.

915 Slater, T., Hogg, A.E. & Mottram, R. (2020). Ice-sheet losses track high-end sea-level rise projections. *Nat.*
916 *Clim. Chang.*, 10, 879–881.

917 South, A., Eriksson, K. & Levitt, R. (2018). How Infrastructure Public–Private Partnership Projects Change
918 Over Project Development Phases. *Project Management Journal*, 49, 62–80.

919 Sowby, R.B. & Hotchkiss, R.H. (2022). Minimizing Unintended Consequences of Water Resources Decisions.
920 *Journal of Water Resources Planning and Management*, 148, 02522007.

921 Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O. & Ludwig, C. (2015a). The trajectory of the
922 Anthropocene: the great acceleration. *The Anthropocene Review*, 2, 81–98.

923 Steffen, W., Persson, Å., Deutsch, L., Zalasiewicz, J., Williams, M., Richardson, K., *et al.* (2011). The
924 Anthropocene: From Global Change to Planetary Stewardship. *Ambio*, 40, 739–761.

925 Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., *et al.* (2015b). Planetary
926 boundaries: Guiding human development on a changing planet. *Science*, 347, 1259855.

927 Strauss, J., Biasi, C., Sanders, T., Abbott, B.W., von Deimling, T.S., Voigt, C., *et al.* (2022). A globally relevant
928 stock of soil nitrogen in the Yedoma permafrost domain. *Nat Commun*, 13, 6074.

929 Syvitski, J., Ángel, J.R., Saito, Y., Overeem, I., Vörösmarty, C.J., Wang, H., *et al.* (2022). Earth’s sediment cycle
930 during the Anthropocene. *Nat Rev Earth Environ*, 3, 179–196.

931 Tank, S.E., Vonk, J.E., Walvoord, M.A., McClelland, J.W., Laurion, I. & Abbott, B.W. (2020). Landscape
932 matters: Predicting the biogeochemical effects of permafrost thaw on aquatic networks with a state
933 factor approach. *Permafrost and Periglacial Processes*.

934 Tansley, A.G. (1935). The Use and Abuse of Vegetational Concepts and Terms. *Ecology*, 16, 284–307.

- Tate, E., Rahman, M.A., Emrich, C.T. & Sampson, C.C. (2021). Flood exposure and social vulnerability in the United States. *Nat Hazards*, 106, 435–457.
- Thomas, Z., Abbott, B.W., Troccaz, O., Baudry, J. & Pinay, G. (2016). Proximate and ultimate controls on carbon and nutrient dynamics of small agricultural catchments. *Biogeosciences*, 13, 1863–1875.
- Thomas, Z., Rousseau-Gueutin, P., Abbott, B.W., Kolbe, T., Le Lay, H., Marçais, J., *et al.* (2019). Long-term ecological observatories needed to understand ecohydrological systems in the Anthropocene: a catchment-scale case study in Brittany, France. *Reg Environ Change*, 19, 363–377.
- Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A.G., *et al.* (2020). Carbon release through abrupt permafrost thaw. *Nature Geoscience*, 13, 138–143.
- Turner, B.L., Matson, P.A., McCarthy, J.J., Corell, R.W., Christensen, L., Eckley, N., *et al.* (2003). Illustrating the coupled human–environment system for vulnerability analysis: three case studies. *Proceedings of the National Academy of Sciences*, 100, 8080–8085.
- Turner, M.G., Dale, V.H. & Gardner, R.H. (1989). Predicting across scales: theory development and testing. *Landscape ecology*, 3, 245–252.
- Underwood, K.L., Rizzo, D.M., Dewoolkar, M.M. & Kline, M. (2021). Analysis of reach-scale sediment process domains in glacially-conditioned catchments using self-organizing maps. *Geomorphology*, 382, 107684.
- Underwood, K.L., Rizzo, D.M., Schroth, A.W. & Dewoolkar, M.M. (2017). Evaluating Spatial Variability in Sediment and Phosphorus Concentration-Discharge Relationships Using Bayesian Inference and Self-Organizing Maps. *Water Resources Research*, 53, 10293–10316.
- Van Loon, A.F., Gleeson, T., Clark, J., Van Dijk, A.I.J.M., Stahl, K., Hannaford, J., *et al.* (2016). Drought in the Anthropocene. *Nature Geoscience*, 9, 89–91.
- Vohra, K., Vodonos, A., Schwartz, J., Marais, E.A., Sulprizio, M.P. & Mickley, L.J. (2021). Global mortality from outdoor fine particle pollution generated by fossil fuel combustion: Results from GEOS-Chem. *Environmental Research*, 195, 110754.
- Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P., *et al.* (2010). Global threats to human water security and river biodiversity. *Nature*, 467, 555–561.
- Wang, H. & He, G. (2022). Rivers: Linking nature, life, and civilization. *River*, 1, 25–36.
- Watson, J.E.M., Venter, O., Lee, J., Jones, K.R., Robinson, J.G., Possingham, H.P., *et al.* (2018). Protect the last of the wild. *Nature*, 563, 27.
- Webber, Z.R., Webber, K.G.I., Rock, T., St. Clair, I., Thompson, C., Groenwald, S., *et al.* (2021). Diné citizen science: Phytoremediation of uranium and arsenic in the Navajo Nation. *Science of The Total Environment*, 794, 148665.
- Wettstad, J. (2018). *Clearing the Air: European Advances in Tackling Acid Rain and Atmospheric Pollution*. Routledge.
- Wilkinson, B.H. (2005). Humans as geologic agents: A deep-time perspective. *Geology*, 33, 161–164.
- Wilson, M.P., Foulger, G.R., Gluyas, J.G., Davies, R.J. & Julian, B.R. (2017). HiQuake: The Human-Induced Earthquake Database. *Seismological Research Letters*, 88, 1560–1565.
- Wohl, E. (2019). Forgotten Legacies: Understanding and Mitigating Historical Human Alterations of River Corridors. *Water Resources Research*, 55, 5181–5201.
- Wohl, E., Lane, S.N. & Wilcox, A.C. (2015). The science and practice of river restoration. *Water Resources Research*, 51, 5974–5997.
- Wolock, D.M. (1995). Effects of Subbasin Size on Topographic Characteristics and Simulated Flow Paths in Sleepers River Watershed, Vermont. *Water Resour. Res.*, 31, 1989–1997.
- Wu, R., Hamshaw, S.D., Yang, L., Kincaid, D.W., Etheridge, R. & Ghasemkhani, A. (2022). Data Imputation for Multivariate Time Series Sensor Data With Large Gaps of Missing Data. *IEEE Sensors Journal*, 22, 10671–10683.
- Zarnetske, J.P., Bouda, M., Abbott, B.W., Saiers, J. & Raymond, P.A. (2018). Generality of Hydrologic Transport Limitation of Watershed Organic Carbon Flux Across Ecoregions of the United States. *Geophysical Research Letters*, 45, 11,702–11,711.
- Zhang, B., Hu, X. & Gu, M. (2022). Promote pro-environmental behaviour through social media: An empirical study based on Ant Forest. *Environmental Science & Policy*, 137, 216–227.