

The Usefulness of Streamflow Reconstructions: Understanding the Management Perspective

The Usefulness of Streamflow Reconstructions: Understanding the Management Perspective
 Connie Woodhouse
 School of Geography, Development, and Environment and the Laboratory of Tree-Ring Research, University of Arizona, Tucson, AZ

I. Introduction

This goal of this presentation is to shed some light on how water managers might perceive the usefulness of tree-ring based hydrologic reconstructions. It is my hope that these insights may help others in their work with water resource managers. I am not a water resource manager (or hydrologist), so these reflections are based on my experience meeting with water managers and the insights I have gained on the management perspective from these.

The usefulness of extended records of streamflow have long been noted by hydrologic reconstruction system scholars. Sagar recent provides a broader range of the utility of extensive and often costly and/or ongoing records to be evaluated in a long-term context.

A very clear example of this is the discussion of Colorado River water in the 1970s. The Colorado River Commission (1970) reported that the reconstruction of the Colorado River water in the 1970s provided a broader range of the utility of extensive and often costly and/or ongoing records to be evaluated in a long-term context.

II. Challenges in using streamflow reconstructions in water resource management

There are a number of challenges involved in using streamflow reconstructions in management decisions. In general, for scientific information to be applicable to decision-making it must be deemed to be robust, credible, and legitimate (Cohn et al. 2002), and this is not often the case if managers and managers have not collaborated on the development of the reconstructions. In the world, streamflow reconstructions must be in a range of formats and the information I present must be relevant to a current management concern. The credibility of the reconstructions, assessed in terms of accuracy, valid and high quality information, is evaluated through the peer-review process, substantially, but via other types of "peer" assessments as well. Credibility can also be related to the skill of the reconstruction. Finally, in order to be perceived as legitimate, the process by which the reconstruction was developed must be transparent and clearly evident upon the introduction.

III. Applications of streamflow reconstructions in water resource management and planning

When managers engage with water resource managers, the potential for the application of streamflow reconstructions to decision making and planning is increased. Engagement can take a variety of forms, from the manager serving as a consultant to a fully interdisciplinary process with resource managers as equal partners in a research project at every step of the way. While most people in the management world are not experts in hydrology, they are experts in water use of streamflow reconstructions, an important assessment, of rapidly changing water resources, water use indicators appropriate to collaboration with water stakeholders who have more direct access to higher quality data.

There is a wide range of ways in which reconstructions of streamflow based on tree rings can be applied to water resource management. From my experience with colleagues and a number of different water resource agencies in Colorado, I provide some examples. It is understood that, in certain key moments of past streamflow provide an assessment of the nature and range of hydrologic variability beyond that documented through records. In the Rio Grande headwaters region, the reconstruction of Rio Grande streamflow (1990) allowed water managers insights on the long-term variability of the system, with implications for the current level of groundwater pumping in the basin.

IV. Concluding Thoughts

1. In my experience, there have been a very limited number of cases in which it is possible to point to a decision or a change in management that was made based on new information from a streamflow reconstruction.
2. Most often, water managers intended to use reconstructions do not have a specific application in mind. The information from the reconstruction may be used primarily for decision-making or education. If the information is used in a more specific manner of application may not be immediate or tangible. Actual use of this information may depend on a variety of factors, including timing, funding, leadership, climate conditions, and needs.
3. The application of information from streamflow reconstructions is often a matter of decision, not a matter of fact. It is important to consider the nature of the decision, the nature of the information, and the nature of the decision-making process. The decision-making process includes both a matter of fact and a matter of decision. It is important to consider the nature of the decision, the nature of the information, and the nature of the decision-making process.
4. There is a growing trend toward the use of streamflow reconstructions in water resource management.

Abstract **References** **Contact Author** **Get Poster**

Connie A. Woodhouse

School of Geography, Development, and Environment and the Laboratory of Tree-Ring Research, University of Arizona, Tucson AZ



PRESENTED AT:

AGU FALL MEETING
 Chicago, IL & Online Everywhere
 12–16 December 2022

SCIENCE LEADS THE FUTURE

I. INTRODUCTION


This goal of this presentation is to shed some light on how water managers might perceive the usefulness of tree-ring based hydrologic reconstructions in the hope that these insights may help others in their work with water resource managers. I am not a water resource manager (or a hydrologist!), so these reflections are based on my experiences working with water managers and the insights I have gained on the management perspective from these experiences.

The usefulness of extended records of streamflow from tree-ring based hydrologic reconstructions seems obvious -- a longer record provides a broader range of the variability of extremes and allows recent and/or ongoing events to be evaluated in a long-term context.

A very classic example of this is the allocation of Colorado River water in the 1920s. The Colorado River Commissioners (*photo below*), representing each of the seven basin states, negotiated the allocation based on an overly optimistic estimate of average annual flow (Kuhn and Fleck 2019). The resulting Colorado River Compact allocated 7.5 million acre feet (MAF) to the upper basin and the same to the lower basin, as measured at the Lees Ferry gage, with an additional 1.5 MAF allocated to Mexico in 1944, for a total of 16.5 MAF.



In 1976, the first reconstructions of Colorado River flow provided a long-term context for this allocation. Stockton and Jacoby (1976) produced several different reconstruction models, but their “best” reconstruction (1520–1961 CE), had a long-term average of just 13.5 MAF (*figure below*). These findings had clear implications for the over-allocation of Colorado River resources.



Another more recent example is the reconstruction of drought based on a gridded network of soil moisture for the US Southwest from Williams et al. (2022) which found the recent 22-year period, 2000-2021, to be the driest period in the past 1200 years. Moreover, this dryness was strongly exacerbated due to anthropogenic warming. This result clearly has implications for water resource managers in this region. What is not clear is exactly how this information could be applied to management.

Rapid intensification of the emerging southwestern North American megadrought in 2020-2021

A. Park Williams^{1,2}, Benjamin I. Cook^{2,3} and Jason E. Smerdon²


A previous reconstruction back to 800 CE indicated that the 2000-2018 soil moisture deficit in southwestern North America was exceeded during one megadrought in the late-1500s. Here, we show that after exceptional drought severity in 2021, ~19% of which is attributable to anthropogenic climate trends, 2000-2021 was the driest 22-yr period since at least 800. This drought will very likely persist through 2022, matching the duration of the late-1500s megadrought.

Since the year 2000, southwestern North America (SWNA 30-45°N, 105-125°W) has been unusually dry due to low precipitation totals and heat, punctuated most recently by exceptional drought in 2021¹. From 2000 to 2021, mean water-year (October-September) SWNA precipitation was 8.3% below the 1950-1998 average and temperature was 0.91 °C above average (Extended Data Fig. 1). No other 22-yr period since at least 1901 was as dry or as hot. While there have been single-year breaks in these anomalous conditions, aridity has dominated the 2000s, as evidenced




II. CHALLENGES IN USING STREAMFLOW RECONSTRUCTIONS IN WATER RESOURCE MANAGEMENT


There are a number of challenges involved in using streamflow reconstructions in management decisions. In general, for scientific information to be applicable to decision making, it must be deemed to be salient, credible, and legitimate (Cash et al. 2002), and this is often not the case if researchers and managers have not collaborated on the development of the reconstructions. To be useful, a streamflow reconstruction must be for a gage of interest and the information it provides must be relevant to a current management concern. The credibility of the reconstruction, assessed in terms of accurate, valid, and high-quality information, is evaluated through the peer-review process, scientifically, but via other types of “peer” assessments as well. Credibility can also be related to the skill of the reconstruction. Finally, in order to be perceived as legitimate, the process by which the reconstruction was developed must be transparent and ideally involve input from the intended users.



In addition to these three factors, the capacity of the water management agency or organization can determine whether (and how) information and data from a streamflow reconstruction can be applied. In my experience, agencies that employ water supply system models may be better positioned to apply these data to management questions, but technical challenges can include the need for spatial and temporal disaggregation in order to match the inputs required by the management model (*example below, for the upper Colorado River*).




Uncertainty in the reconstruction may be difficult to incorporate into system models as well, so while the uncertainty information is important to convey, reconstruction ensembles may not be useable. Agencies that do not use water system models may find it more difficult to determine the value of a reconstruction, but it can be informative in a more qualitative way (Woodhouse and Lukas 2006). Related to capacity, timing can also be a critical factor. Water managers are often receptive to new types of information when events such as droughts present management challenges, but much less so when water supplies appear to be plentiful (e.g., Lake Mead in 1999 vs 2021, *below*).



III. APPLICATIONS OF STREAMFLOW RECONSTRUCTIONS IN WATER RESOURCE MANAGEMENT AND PLANNING

When researchers engage with water resource managers, the potential for the application of streamflow reconstructions to decision making and planning is increased. Engagement can take a variety of forms, from the researcher serving as a consultant to a fully transdisciplinary process with resource managers as equal partners in a research project at every step of the way. While water providers or agencies with resources to support a scientific staff have the best capacity to make use of streamflow reconstructions, an increased awareness of equity issues may prompt researchers to use more inclusive approaches to collaboration with water stakeholders who have more limited capacity to explore new types of data.

There is a wide range of ways in which reconstructions of streamflow based on tree rings are being use in water resource management. I use my experiences with colleagues and a number of different water resource agencies in Colorado to provide some examples. At a fundamental level, centuries-long records of past streamflow provide an awareness of the nature and range of hydrologic variability, beyond that documented in gage records. In the Rio Grande headwaters region, the reconstruction of Rio Grande streamflow (*below*) offered water managers insights on the low frequency variability of the system, with implications for the current level of groundwater pumping in the basin.



This extended range of variability also documents extremes, including pluvial periods and multi-year droughts. Before the current drought, water managers in Colorado had commonly used the 1950s drought as a “worst-case” scenario. Reconstructions record how often 1950s-scale droughts occur, as well as droughts that were even more severe. Several water providers in the Colorado Front Range used this information to either confirm the robustness of their drought planning or to adopt new worst case scenarios (Denver Water is an example, *below*)




What do we know about past droughts? Will climate change cause worse droughts?

By studying the growth pattern of old trees, scientists have estimated streamflow patterns in Denver's watersheds going back 400 years. Denver Water uses these estimates to analyze past droughts against water supply needs. Tree ring data indicate several droughts occurred in the past 400 years that were much worse than the modern droughts of the 1950s and the 2000s. Unfortunately, climate change could cause droughts to be worse than those of the last 400 years.

<https://www.denverwater.org/your-water/water-supply-and-planning/climate-change>

Water providers and agencies who use water supply system models have used reconstructions of streamflow as input into models to see how the system performs under the worst droughts of the past and for sensitivity testing. An example of this is the Bureau of Reclamation's use of Colorado River reconstructions in their Colorado River Simulation System model, first to test system response, including changes in reservoir levels, to a multidecadal drought in the 1100s, and then in supply scenarios in their Basin Study (Woodhouse and Lukas 2020, Reclamation 2012) (*figure below*).

FIGURE A-8
Illustration of Combined Water Supply and Water Demand Scenarios



From Reclamation's Colorado River Basin Study (2012)

In addition, we have found that the reconstructions of past streamflow have been used by water managers to educate their customers and to communicate risk and support recommendations to decision makers (Rice et al. 2009). The table

below includes this and other uses from a survey of 29 water managers in the Colorado Front Range.



TABLE 5. Survey Responses to the Question “How Have Tree-Ring Data Been Used by You, Your Organization, or Organizations That You Consult For?” (respondents could select more than one answer).

| | |
|--|-----|
| To broaden understanding of hydrologic variability | 75% |
| To educate users/public | 46% |
| To educate board and other decision makers | 50% |
| As input into a water system model or other model | 25% |
| For quantitative analysis, but not in a modeling environment | 14% |
| To inform planning and decision making | 54% |
| I have not used tree-rings in my organization | 18% |

From Rice et al. 2009


IV. CONCLUDING THOUGHTS

1. In my experience, there have been a very limited number of cases in which it is possible to point to a decision or a change in management that was made based on new information from a streamflow reconstruction.
2. Most often, water managers interested in a reconstruction do not have a concrete application in mind. The information from the reconstruction may be useful primarily for awareness-raising or education. If the information is used in some way, evidence of application may not be immediate or tangible. Actual use of this information may depend on a variety of factors, including timing, funding, leadership, and climate conditions and events.
3. The application of information from streamflow reconstructions is often in support of decisions that consider a host of other factors. It is important to recognize that scientific information is just one of a number of considerations in decision making. The decision-making context includes issues such as existing policies and planning practices, economic and financial constraints, risk/benefit trade-offs, politics, and overall feasibility of incorporating new data or information.



4. There is a growing track record of the use of streamflow reconstructions in water management in watersheds across the western US (and in other areas as well). These examples may help water managers in other watersheds consider how this information may be useful to them. My experience is that water managers greatly value learning from their peers and hearing about their experiences with tree-ring data.
5. Some have suggested that records from the past are no longer relevant given anthropogenic climate change. It is true that the past is not an analogue for the future. However, plausible future scenarios for water management include the range of conditions documented by paleohydrologic data with the added influence of warming (*illustrated below*). The blending of these two is being accomplished in a variety of ways, providing important insights for future planning.

Plausible Scenarios for the Future



Acknowledgements

Many colleagues have collaborated with me over the past two decades on the work that I've described here, including both academic and governmental researchers and water resource managers. I am extremely appreciative of these collaborations. Support for this work has come from a variety of grants and agencies including the NOAA Climate Program Office RISA/CAP Programs (CLIMAS and WWA), the DOI Southwest Climate Adaptation Science Center, California Department of Water Resources, Bureau of Reclamation, and the National Science Foundation.

ABSTRACT

The usefulness of extended records of streamflow from tree-ring based hydrologic reconstructions seems obvious; a longer record provides a broader range of the variability of extremes and allows recent and/or ongoing events to be evaluated in a long-term context. The information from these centuries-long records may have clear implications for water resource management, but it is often unclear exactly how this information can be applied to management. In this presentation, I will discuss some of the challenges I have observed that are involved in using streamflow reconstructions in management decisions. These range from issues related to an agency's capacity to use new types of data to mismatches between what is needed (e.g., daily resolution, a network of gage inputs) and what reconstruction data provide. The skillfulness of a streamflow reconstruction also has a bearing on its perceived credibility in terms of useable data. In spite of these challenges, there is a variety of ways that these data have been used by water resource managers in the western US. The uses are often not immediately evident, but can take the form of, for example, sensitivity assessment, awareness raising, and shifts in prior assumptions. Relationship building between researchers and resource managers can yield mutual respect and understanding that lead to both interesting research questions and relevant and valuable information, even if the application to management is not tangible or immediate.

REFERENCES

- Cash, D., W.C. Clark, William C., F. Alcock, N.M. Dickson, N. Eckley, and J. Jäger. 2002. Salience, Credibility, Legitimacy and Boundaries: Linking Research, Assessment and Decision Making. Cambridge, MA: John F. Kennedy School of Government, Harvard University. 24 pp. Available at SSRN: <https://ssrn.com/abstract=372280> or <http://dx.doi.org/10.2139/ssrn.372280>
- Kuhn, E. and J. Fleck. 2019 Science Be Dammed: How Ignoring Inconvenient Science Drained the Colorado River. University of Arizona Press, Tucson. 288 pp.
- Lukas, Jeff, and Elizabeth Payton, eds. 2020. Colorado River Basin Climate and Hydrology: State of the Science. Western Water Assessment, University of Colorado Boulder. <https://doi.org/10.25810/3hcv-w477>
- Reclamation. 2012. "Colorado River Basin Water Supply and Demand Study." Report to US Bureau of Reclamation. https://www.usbr.gov/lc/region/programs/crbstudy/finalreport/Study%20Report/CRBS_Study_Report_FINAL.pdf
- Rice, J.L., C.A. Woodhouse, and J.J. Lukas, 2009. Science and decision-making: water management and tree-ring data in the western United States. *Journal of the American Water Resources Association* 45, 1248-1259. <https://doi.org/10.1111/j.1752-1688.2009.00358.x>
- Stockton, C.W., and G.C. Jacoby. 1976. "Long-Term Surface Water Supply and Streamflow Levels in the Upper Colorado River Basin." Lake Powell Research Project, Bulletin No 18. Los Angeles: Inst. of Geophysics and Planetary Physics, University of California, 70 pp.
- Williams, A.P., B.I. Cook & J.E. Smerdon. 2022. Rapid intensification of the emerging southwestern North American megadrought in 2020–2021. *Nature Climate Change* 12, 232–234. <https://doi.org/10.1038/s41558-022-01290-z>
- Woodhouse, C.A. and J.J. Lukas, 2006. Drought, tree rings, and water resource management. *Canadian Water Resources Journal* 31, 297-310. <https://doi.org/10.4296/cwrj3104297>
- Woodhouse, C.A. and J. Lukas. 2020. "Paleohydrology." Chap. 10 in *Colorado River Basin Climate and Hydrology: State of the Science*, edited by J. Lukas and E. Payton, 361-383. Western Water Assessment, University of Colorado Boulder. <https://doi.org/10.25810/3hcv-w477>

