


Role of Vanadium in Carbonatite Magma at Amba Dongar Diatreme Gujarat, India

From Magmatic to Hydrothermal



Perumala V Sunder Raju* and Shrinivas Viladkar+

* CSIR-National Geophysical Research Institute, Uppal Road, Hyderabad, India + IISER, Bhopal, India



PRESENTED AT:



INTRODUCTION

Carbonatites are volumetrically insignificant (~4%) igneous rocks, intrusive as well as extrusive. They contain more than 50% by volume of carbonate minerals.

They are characterised by the high Sr and rare earth elements abundances

There are around 529 known carbonatites worldwide

They range in age from Archean to present day

Found in about 70 countries and are confined to stable Archean continental crust, intracontinental rift, fault system, Within alkaline rock provinces

In India Amba Dongar is the only carbonatite complex having such high vanadium activity.

Sövite exposures in ring dike of Amba Dongar exhibit magmatic banding that is conspicuous in some parts of the ring dike.

The bands are usually rich in magnetite, titanite, apatite and pyrochlore (Fig. 1). In addition to banding some exposures show randomly oriented extensive veins containing secondary hydrothermal minerals.


These veins are rich in barite, REE-minerals and vanadinite. Barite and magnetite are ubiquitous in Amba Dongar carbonatite .

In the samples studied barite occurs in close association with vanadinite and magnetite.

Magnetite is abundant in these sövite and it forms thick bands (~2 cm) as well as disseminated grains.

Vanadinite occurs in the vicinity of barite, titanite and magnetite (BSE figures 2 and 3).

GEOLOGY



The Ambadongar Carbonatite (Fig 1) is located in Chottanagpur district, Gujarat State,India.

Majority of the area is covered by tholeiitic lava flow of Deccan Traps.

The complex consist of a Sovite ring dyke that has an inner rim of carbonatite breccia.

The diatreme is sourrounded by a Fenite zone at the conatct with the Bagh sandstone.

The first pahse is of thin dykes intruding sovite and fenite.

The second phase form large plugs within sovite ring dykes

The last phase is enriched with REE and Nb and end with flourite mineralization (Fig 2 below)




Fig 2 Geological features of the Amba Dongar area

METHODOLOGY

Powder XRD Methodology

The PXRD analysis was conducted at CSIR-NGRI, India X-Ray facility, using a Bruker D-8 advance model.

It is equipped with a source of Cu K α radiation with high precision LYNXE detector and used a Ni K β filter.

The diffraction patterns collected using knife-edge to reduce air scatter by scanning from 5 to 70 reduce 0.02° 2 θ steps and counting for \approx 3177 s/step for a total scan time of reducing 0.02° 2 θ steps diffraction patterns were evaluated using the Bruker Diffrac. Suite EVA V 4.0, software released in 2016.

Each diffraction pattern was corrected for background. Instrumental shifts in 2 θ were corrected by reference to the 100% intensity (104) K α peak of corundum at 35.149 2 θ positions.

Mineralogical identification was facilitated by using the International Centre of Diffraction Data (ICDD) PDF4+ 2022 database.

The K α 2 stripping is also carried out to avoid doublet and triplet peaks and artifacts



or match fit.

Scanning electron Microscope

The system is a LEO440 device, from Oxford instruments, operated at 15-20Kv with variable voltage ranging between 2-5 nA. Filament was working at the first maximum, about 2.5 nA.

RESULTS

XRD examination we found peaks for two vanadinite minerals (Fig 4) namely Karelianite and Cooperite (PXRD patters are presented in figures 3 and 5.




Fig 3- PXRD pattern of Karelianite and Cooperite




Fig 4 BSE Image of wakefieldite- Ce 47, 49, Vanadinite 48, 50), below is the enlarge view of area containing these minerals.BSE image of titanite (25) and magnetite (26, 28)




Fig 5 PXRD pattern of Wakefeildite 0-70 degrees and also enlarged inset

DISCUSSION

The identified mineral is Wakefieldite with La 71 Nd0.15O14 Pr 0.11 Sm 0.01 V Y 0.02 or La X V Y0.02 (X=Nd, Pr, Sm) (Figures 6, 7 and 8)

The new mineral (IMA 89-035a) is the La analogue of Wakefieldite (Ce) and Wakefieldite(Y) and a member of xenotime group. Wakefieldite (La) has a zircon type crystal structure (tetragonal space group) with $a=7.406(4)$, $C=6.50498$ Å and $V=356.8$ (6) Å³ from XRD data.

The peak positions identified at 2 theta positions as shown in figures at 32.34, 47 and 48 degrees.

Karelianite (V2O3)

The chemical composition is V₂O₃ and belongs to hexagonal crystal system. The PXRD pattern shows similarity by analogy to hematite group and presents as prismatic grains. The cell data with space group is R3C, $a=4.99$, $c=13.98$ Å, $Z=6$. It is associated with Pyrrhotite, chalcopyrite, pyrite, tremolite, graphite, titanite, quartz (Outokumpu mine, Finland); corvusite, montroseite, uraninite, quartz (Mounana mine, Gabon); millerite, violarite, montroseite (Guangxi, China). Because the vanadium-oxygen bonds have some covalent character V⁺³ is chosen as a compromise between unionized V and fully ionized V⁺⁵ (Trotter and Barnes 1958).

Cooperite belongs to Tetragonal crystal system and, space group P4₂/mmc. with Unit cell data: $a=3.465$ Å, $c=6.104$ Å, $V=73.29$ Å³, $Z=2$. Class: 4/mmm; $c/a=1.7616$. The crystal structure is of PtS type with strong and directed covalent bonds. M (M = Pt, Pd) forms four co-planar bonds with X (X = O, S), which itself is coordinated by four M in a distorted tetrahedral environment. The standard X-ray powder diffraction data: 3.013 (100) (011), 2.450 (60) (110), 1.911 (80) (112), 1.753 (60) (013), 1.732 (50) (020), 1.507 (70) (022), 1.231 (50) (123).

CONCLUSIONS

High Ti minerals in Amba Dongar are ilmenite, perovskite, niobianzirconolite and betafite

Cooperite is rarely found in carbonatites

Significant concentration of Vanadium is present in Study area

Both magmatic and hydrothermal inputs in study area,

Hence, there is a chance to relook for Vanadium, Titanium, Pt bearing minerals and multimetal association in Carbonatites .


AUTHOR INFORMATION

P V SUNDER RAJU


***CSIR-National Geophysical Research Institute, Uppal Road,
Hyderabad, India***

ABSTRACT


In carbonatites high vanadium concentration is not common and it is usually present in trace amounts in spinel, titanite or magnetite and some ferromagnesian silicate minerals. V-minerals (vanadinite, wakefieldite) have not been reported from magmatic carbonatites barring some volcanic carbonatites of Italy. In Amba Dongar sövite, however both these minerals along with Karelitanite and Platinum group mineral-cooperate have been detected. It is noted that vanadium in low concentration is found in magmatic stage while its concentration reaches high enough in hydrothermal solutions to form vanadinite, wakefieldite and karelitanite. So far perhaps Amba Dongar is the only carbonatite complex having such high vanadium activity.



XRD Pattern of whole range of 2theta from 2-70 degree
(Wakefieldite at 47 and 48 degree 2theta) hosted in
Calcite



XRD pattern showing Wakefieldite (2theta angle)



BSE Image of wakefieldite- Ce 47, 49,
Vanadinite 48, 50), below is the enlarge view of
area containing these minerals.

(https://agu.confex.com/data/abstract/agu/fm22/1/9/Paper_1049091_abstract_952128_0.jpg)

REFERENCES

- Bordage, E. Balan, J.P.R. de Villiers, R. Cromarty, A. Juhin, C. Carvallo, G. Calas, P.V. Sunder Raju, P. Glatzel (2011) V oxidation state in Fe- Ti oxides by high-energy resolution fluorescence-detected x-ray absorption spectroscopy. *Physics and Chemistry of Minerals*, 38: pp 449-458
- Cheng Xu, Liang Qi, Zhilong Huang, Yanjing Chen, Xuehui Yu, Linjun Wang, and Endong Li (2008) Abundances and significance of platinum group elements in carbonatites from China. *Lithos*, 105, pp 201-207
- Cabri, L.J., Laflamme, J.H.E., Turner, K and Skinner B.J. (1978) On cooperite, braggite, and vysotskite. *American Mineralogist*, 63, pp. 832-839.
- Doroshkevich, A.G., Viladkar, S.G, Ripp, G.S. and Burtseva, M.V. (2009) Hydrothermal REE mineralization in the Amba Dongar Carbonatite complex, Gujarat, India. *Can. Mineral.* V. 47, pp. 1105-1116.
- Jones, A.P., Wall, F., Williams, C.T., 1996. Rare earth minerals: chemistry, origin and ore deposits. In: *The Mineralogical Society Series* Springer, Netherlands, pp. 372
- Kelley, K.D., Scott, C.T., Polyak D.E, and Kimball, B.E (2005) Vanadium Chapter U of Critical Mineral Resources of the United State-Economic and Environmental Geology and Prospects for Future Supply Vanadium Professional Paper 1802-USGS
- Long, J.V.P. (1963) Karelitanite, a new vanadium minerals. *Amer. Mineral.* V. 48, pp 33-41
- Magna, T., Viladkar, S Rappricha, V., Poura, O, Hopp, J., and Čejkova, B., (2020) Nb–V-enriched sovites of the northeastern and eastern part of the Amba Dongar carbonatite ring dike, India – A reflection of post-Emplacement hydrothermal overprint? *Geochemistry*, 80, pp 1-11
- Mariano, A.N. (1989) Nature of economic mineralization in carbonatites and related rocks. In Bell K. (Ed) *Carbonatites, Genesis and Evolution*, Unwyn Hyman, London, pp. 149-172
- Miles, N.M., Hogarth, D.D. and Russel, D.S. (1971) Wakefieldite, Yttrium Vanadate, A new mineral from Quebec. *Amer. Mineral.* V. 56, pp 395-410.
- Mitchell, R.H., Keays, R.R., 1981. Abundance and distribution of gold, Palladium and iridium in some spinel and garnet lherzolites-implications for the nature and origin of precious metal-rich intergranular components in the upper mantle. *Geochimica et Cosmochimica Acta* 45, pp 2425–2442.
- Nelson, D.R., Chivas, A.R., Chappell, B.W., McCulloch, M.T., 1988. Geochemical and isotopic systematics in carbonatites and implications for the evolution of ocean–island sources. *Geochimica et Cosmochimica Acta* 52, pp 1–17.
- Rudashevsky, N.S., Kertser, Y.L., Bulakh, A.G., Rudashevsky, V.N., 2001. Two type of PGE mineralization in carbonatite deposit. *Journal of African Earth Sciences* 32, pp 30–30.
- Rudashevsky, N.S., Krester, Yu.L., Rudashevsky, V.N., and Sukharzhevskaya, E.S. (2004) A review and comparison of PGE, noble metal and sulphide mineralization in phosphorites and carbonatites from Kovodor and Phalaborwa. In: *Phosphorites and carbonatites from mantle to mine: The key example of the Kola Alkaline Province* (Eds Zeitsev, A, and Wall, F.) Mineralogical Society London, Series 10, pp 363-393
- Stoppa, F., Pirajno, F., Schiazza, M., Vladykin, N.V., 2016. State of the art: Italian carbonatites and their potential for critical-metal deposits. *Gondwana Res.* 37, 152–171.
- Trotter, J. and Barnes W, H. (1958): The structure of vanadinite. *Can. Mineral.* 6, pp. 161-173
- Viladkar, S.G. and Wimmenauer, W. (1992) Geochemical and petrological studies on the Amba Dongar carbonatite (Gujarat, India). *Chemie Erde*, 52, pp 277-291
- Zhou, M.F., Sun, M., Keays, R.R., Kerrich, R.W., 1998. Control on platinum-Group elemental distributions of podiform chromitites: a case study of high-Cr and high-Al chromitites from Chinese orogenic belts. *Geochimica et Cosmochimica Acta* 62, pp 677–688.

