

On n-dimension Extended cubic B-splines collocation algorithms

K. R. Raslan, Khalid K. Ali*

Mathematics Department, Faculty of Science, Al-Azhar University, Nasr-City, Cairo, Egypt.

Email: kamal_raslan@yahoo.com

Email*: khalidkaram2012@azhar.edu.eg

Abstract

In this work, we present a solution to a major problem that most researchers meet, which is the solution of differential equations of different dimensions by presenting a new structure to n-dimensions for the Extended cubic B-spline collocation algorithm. The Extended cubic B-spline collocation forms are displayed in one, two and three dimensions. These constructs are of prime importance in solving mathematical models that have applications in various sciences. The efficiency and accuracy of these algorithms through a few test problems in two and three dimensions. Also, comparing our solutions and with the results obtained by using other numerical methods available in the literature as much as possible.

Keywords: Collocation method; n-dimensions; The Extended cubic B-splines; Error estimation.

1 Introduction

We all know that obtaining solutions to partial differential equations is of great importance in various fields. Some of these equations arise from different problems in fields such as physics, chemistry, engineering and others. These problems are converted into mathematical models. The solutions to these equations are divided into analytical solutions and numerical solutions [1–5]. Recently, researchers have tended to use different methods to find these solutions, whether analytical or numerical. With the existence of models for these equations that have a degree of difficulty in finding solutions to them, especially if they are in two dimensions or in three dimensions or more than that, serious work continued on how to find these solutions. Over time, some researchers in this field found it difficult to find analytical solutions with different dimensions for these models, so some of them went to find numerical solutions to them. Several researchers have used different numerical methods to find solutions to these equations with different dimensions [6–16]. Now, we are continuing to work on developing a different basis B-spline collocation method to find numerical solutions to partial differential equations in two and three dimensions and so on. This work is a continuation of the works in [17,18]. The collocation strategy began by Frazer et al. [19] in 1937. Afterward, the collocation strategy at the side the least-squares strategy and Galerkin strategy was utilized by Bickley [20] to ponder shaky heat condition issues. Afterward on, in 1975, the collocation strategy beside B-splines was connected to shaky warm conduction and boundary layer streams [21] and it was found that the comes about gotten were superior when compared with comes about gotten with limited contrast strategies. Since at that point, the

collocation strategy is being utilized over a wide extend of issues [22–29]. In combination with the collocation strategy, there has been serious utilization of polynomial B-splines for understanding halfway differential conditions. Cubic B-splines, quasi B-splines, quartic B-splines, quintic B-splines, and so on are utilized in combination with the collocation strategy in [22–27] for managing with different straight and nonlinear boundary esteem issues. Strategies like Haar wavelet collocation strategy [30], a slope replicating part collocation strategy [31] and Newton premise capacities collocation strategy [32] are moreover picking up ubiquity to illuminate differential conditions.

In this work, we present the Extended cubic B-spline collocation algorithm forms in n-dimensions. In addition, some numerical examples are proposed to study the effectiveness and accuracy of this method.

This article is organized as follows: The second section introduces n-dimensions Extended cubic B-spline formulations. The third section introduces numerical examples. The error estimates are present in section four. Finally, show the conclusion part.

2 Construct Extended cubic B-spline Formulas

The forms for n-dimensions Extended cubic B-splines were introduced in this part.

2.1 One dimension Extended cubic B-spline [21]

Let $x \in [a, b]$ and $\phi_i(x)$ are those Extended cubic B-spline with knots at the points x_i . Then the set of Extended cubic B-splines $\phi_{-1}(x), \phi_0(x), \dots, \phi_{N-1}(x), \phi_N(x), \phi_{N+1}$, forms a basis for functions defined over the interval. The approximation $U^N(x)$ to $U(x)$ which uses these splines as :

$$U^N(x) = \sum_{i=-1}^{N+1} \chi_i \phi_i(x), \quad (1)$$

where χ_i unknown term. The formulations of $U_i, \frac{dU_i}{dx}, \frac{d^2U_i}{dx^2}$ are given by:

$$\begin{aligned} U_i &= \frac{1}{24} (-(\lambda - 4)\chi_{i-1} + 2(\lambda + 8)\chi_i - (\lambda - 4)\chi_{i+1}), \\ \frac{dU_i}{dx} &= \frac{\chi_{i+1} - \chi_{i-1}}{2h}, \\ \frac{d^2U_i}{dx^2} &= \frac{(\lambda + 2)(\chi_{i-1} - 2\chi_i + \chi_{i+1})}{2h^2}. \end{aligned} \quad (2)$$

2.2 Two dimensions Extended cubic B-spline

In this subsection, we show the formula of Extended cubic B-spline in two dimensions on a rectangular grid divided into regular rectangular finite elements on both sides. $h = \Delta x, k = \Delta y$ by the knots (x_m, y_n) where $m = 0, 1, \dots, M, n = 0, 1, \dots, N$. The approximation $U^N(x, y)$ to $U(x, y)$ given by:

$$U^N(x, y) = \sum_{m=-1}^{M+1} \sum_{n=-1}^{N+1} \chi_{m,n} B_{m,n}(x, y), \quad (3)$$

where $\chi_{m,n}$ are the amplitudes of the Extended cubic B-splines $B_{m,n}(x,y)$ given by

$$B_{m,n}(x,y) = \phi_m(x) \phi_n(y).$$

Which peaks on the knot (x_m, y_n) and $\phi_m(x)$, $\phi_n(y)$ are identical in form to the one dimension Extended cubic B-splines. Then the formulations of $U_{m,n}$, $\frac{\partial U_{m,n}}{\partial x}$, $\frac{\partial U_{m,n}}{\partial y}$, $\frac{\partial^2 U_{m,n}}{\partial x^2}$, $\frac{\partial^2 U_{m,n}}{\partial y^2}$, $\frac{\partial^2 U_{m,n}}{\partial x \partial y}$, ... are given by:

$$\begin{aligned} U_{m,n} &= \frac{1}{576} \\ &\left(-2(\lambda^2 + 4\lambda - 32)\chi_{m-1,n} + \lambda^2\chi_{m-1,n+1} - 2\lambda^2\chi_{m,n-1} \right. \\ &\quad + 4\lambda^2\chi_{m,n} - 2\lambda^2\chi_{m,n+1} + \lambda^2\chi_{m+1,n-1} - 2\lambda^2\chi_{m+1,n} + \lambda^2\chi_{m+1,n+1} \\ &\quad + (\lambda - 4)^2\chi_{m-1,n-1} - 8\lambda\chi_{m-1,n+1} - 8\lambda\chi_{m,n-1} + 64\lambda\chi_{m,n} - 8\lambda\chi_{m,n+1} \\ &\quad - 8\lambda\chi_{m+1,n-1} - 8\lambda\chi_{m+1,n} - 8\lambda\chi_{m+1,n+1} + 16\chi_{m-1,n+1} + 64\chi_{m,n-1} \\ &\quad \left. + 256\chi_{m,n} + 64\chi_{m,n+1} + 16\chi_{m+1,n-1} + 64\chi_{m+1,n} + 16\chi_{m+1,n+1} \right). \end{aligned} \quad (4)$$

$$\begin{aligned} \frac{\partial U_{m,n}}{\partial x} &= \frac{1}{48h} \left((\lambda - 4)\chi_{m-1,n-1} - 2(\lambda + 8)\chi_{m-1,n} + \lambda\chi_{m-1,n+1} - \lambda\chi_{m+1,n-1} \right. \\ &\quad + 2\lambda\chi_{m+1,n} - \lambda\chi_{m+1,n+1} - 4\chi_{m-1,n+1} + 4\chi_{m+1,n-1} + 16\chi_{m+1,n} \\ &\quad \left. + 4\chi_{m+1,n+1} \right), \\ \frac{\partial U_{m,n}}{\partial y} &= \frac{1}{48k} \left((\lambda - 4)\chi_{m-1,n-1} - (\lambda - 4)\chi_{m-1,n+1} - 2\lambda\chi_{m,n-1} \right. \\ &\quad + 2\lambda\chi_{m,n+1} + \lambda\chi_{m+1,n-1} - \lambda\chi_{m+1,n+1} - 16\chi_{m,n-1} + 16\chi_{m,n+1} \\ &\quad \left. - 4\chi_{m+1,n-1} + 4\chi_{m+1,n+1} \right). \end{aligned} \quad (5)$$

$$\begin{aligned} \frac{\partial^2 U_{m,n}}{\partial x^2} &= -\frac{\lambda + 2}{48h^2} \\ &\left((\lambda - 4)\chi_{m-1,n-1} - 2(\lambda + 8)\chi_{m-1,n} + \lambda\chi_{m-1,n+1} - 2\lambda\chi_{m,n-1} + 4\lambda\chi_{m,n} \right. \\ &\quad - 2\lambda\chi_{m,n+1} + \lambda\chi_{m+1,n-1} - 2\lambda\chi_{m+1,n} + \lambda\chi_{m+1,n+1} - 4\chi_{m-1,n+1} \\ &\quad \left. + 8\chi_{m,n-1} + 32\chi_{m,n} + 8\chi_{m,n+1} - 4\chi_{m+1,n-1} - 16\chi_{m+1,n} - 4\chi_{m+1,n+1} \right), \\ \frac{\partial^2 U_{m,n}}{\partial y^2} &= -\frac{\lambda + 2}{48k^2} \left((\lambda - 4)\chi_{m-1,n-1} - 2(\lambda - 4)\chi_{m-1,n} \right. \\ &\quad + \lambda\chi_{m-1,n+1} - 2\lambda\chi_{m,n-1} + 4\lambda\chi_{m,n} - 2\lambda\chi_{m,n+1} + \lambda\chi_{m+1,n-1} \\ &\quad - 2\lambda\chi_{m+1,n} + \lambda\chi_{m+1,n+1} - 4\chi_{m-1,n+1} - 16\chi_{m,n-1} + 32\chi_{m,n} - 16\chi_{m,n+1} \\ &\quad \left. - 4\chi_{m+1,n-1} + 8\chi_{m+1,n} - 4\chi_{m+1,n+1} \right). \end{aligned} \quad (6)$$

$$\begin{aligned}
\frac{\partial^2 U_{m,n}}{\partial x \partial y} &= \frac{\chi_{m-1,n-1} - \chi_{m-1,n+1} - \chi_{m+1,n-1} + \chi_{m+1,n+1}}{4hk}, \\
\frac{\partial^3 U_{m,n}}{\partial x^2 \partial y} &= -\frac{(\lambda + 2)}{4h^2k} \\
&(\chi_{m-1,n-1} - \chi_{m-1,n+1} - 2\chi_{m,n-1} + 2\chi_{m,n+1} + \chi_{m+1,n-1} - \chi_{m+1,n+1}), \quad (7) \\
\frac{\partial^3 U_{m,n}}{\partial x \partial y^2} &= -\frac{(\lambda + 2)}{4hk^2} \\
&(\chi_{m-1,n-1} - 2\chi_{m-1,n} + \chi_{m-1,n+1} - \chi_{m+1,n-1} + 2\chi_{m+1,n} - \chi_{m+1,n+1}), \\
&\vdots
\end{aligned}$$

2.3 The three dimensions Extended cubic B-spline

Now, we obtain the Extended cubic B-spline in three measurements approximates on a framework subdivided into limited components of sides $h = \Delta x$, $k = \Delta y$, $q = \Delta z$ by the knots (x_m, y_n, z_r) where $m = 0, 1, \dots, M$, $n = 0, 1, \dots, N$, $r = 0, 1, \dots, R$ can be interpolated in terms of piecewise Extended cubic B-splines. If $U(x, y, z)$ is a function of x, y and z , it can be shown there exists a unique approximation $U^N(x, y, z)$ as

$$U^N(x, y, z) = \sum_{m=-1}^{M+1} \sum_{n=-1}^{N+1} \sum_{r=-1}^{R+1} \chi_{m,n,r} B_{m,n,r}(x, y, z), \quad (8)$$

where $\chi_{m,n,r}$ are the amplitudes of the Extended cubic B-splines $B_{m,n,r}(x, y, z)$ given by

$$B_{m,n,r}(x, y, z) = \phi_m(x) \phi_n(y) \phi_r(z).$$

Also, $\phi_m(x)$, $\phi_n(y)$ and $\phi_r(z)$ have the same form as the one dimension Extended cubic B-splines. The formulations of $U_{m,n,r}$, $\frac{\partial U_{m,n,r}}{\partial x}$, $\frac{\partial U_{m,n,r}}{\partial y}$, $\frac{\partial U_{m,n,r}}{\partial z}$, $\frac{\partial^2 U_{m,n,r}}{\partial x^2}$, $\frac{\partial^2 U_{m,n,r}}{\partial y^2}$, $\frac{\partial^2 U_{m,n,r}}{\partial z^2}$, $\frac{\partial^2 U_{m,n,r}}{\partial x \partial y}$, $\frac{\partial^2 U_{m,n,r}}{\partial x \partial z}$, \dots , are given in terms of the $\chi_{m,n,r}$ by:

$$\begin{aligned}
U_{m,n,r} &= \frac{1}{13824} \left(2(\lambda + 8)\chi_{m-1,n-1,r}(\lambda - 4)^2 - (\lambda - 4)^3\chi_{m-1,n-1,r-1} \right. \\
&\quad - \lambda^3\chi_{m-1,n-1,r+1} + 12\lambda^2\chi_{m-1,n-1,r+1} - 48\lambda\chi_{m-1,n-1,r+1} \\
&\quad + 64\chi_{m-1,n-1,r+1} + 2\lambda^3\chi_{m-1,n,r-1} - 96\lambda\chi_{m-1,n,r-1} \\
&\quad + 256\chi_{m-1,n,r-1} - 4\lambda^3\chi_{m-1,n,r} - 48\lambda^2\chi_{m-1,n,r} + 1024\chi_{m-1,n,r} \\
&\quad + 2\lambda^3\chi_{m-1,n,r+1} - 96\lambda\chi_{m-1,n,r+1} + 256\chi_{m-1,n,r+1} - \lambda^3\chi_{m-1,n+1,r-1} \\
&\quad + 12\lambda^2\chi_{m-1,n+1,r-1} - 48\lambda\chi_{m-1,n+1,r-1} + 64\chi_{m-1,n+1,r-1} \\
&\quad + 2\lambda^3\chi_{m-1,n+1,r} - 96\lambda\chi_{m-1,n+1,r} + 256\chi_{m-1,n+1,r} - \lambda^3\chi_{m-1,n+1,r+1} \\
&\quad + 12\lambda^2\chi_{m-1,n+1,r+1} - 48\lambda\chi_{m-1,n+1,r+1} + 64\chi_{m-1,n+1,r+1} \\
&\quad + 2\lambda^3\chi_{m,n-1,r-1} - 96\lambda\chi_{m,n-1,r-1} + 256\chi_{m,n-1,r-1} - 4\lambda^3\chi_{m,n-1,r} \\
&\quad - 48\lambda^2\chi_{m,n-1,r} + 1024\chi_{m,n-1,r} + 2\lambda^3\chi_{m,n-1,r+1} - 96\lambda\chi_{m,n-1,r+1} \\
&\quad + 256\chi_{m,n-1,r+1} - 4\lambda^3\chi_{m,n,r-1} - 48\lambda^2\chi_{m,n,r-1} + 1024\chi_{m,n,r-1} \\
&\quad \left. + 8\lambda^3\chi_{m,n,r} + 192\lambda^2\chi_{m,n,r} + 1536\lambda\chi_{m,n,r} + 4096\chi_{m,n,r} - 4\lambda^3\chi_{m,n,r+1} \right)
\end{aligned}$$

$$\begin{aligned}
& + 256\chi_{m,n+1,r-1} - 4\lambda^3\chi_{m,n+1,r} - 48\lambda^2\chi_{m,n+1,r} + 1024\chi_{m,n+1,r} \\
& + 2\lambda^3\chi_{m,n+1,r+1} - 96\lambda\chi_{m,n+1,r+1} + 256\chi_{m,n+1,r+1} - \lambda^3\chi_{m+1,n-1,r-1} \\
& + 12\lambda^2\chi_{m+1,n-1,r-1} - 48\lambda\chi_{m+1,n-1,r-1} + 64\chi_{m+1,n-1,r-1} \\
& + 2\lambda^3\chi_{m+1,n-1,r} - 96\lambda\chi_{m+1,n-1,r} + 256\chi_{m+1,n-1,r} - \lambda^3\chi_{m+1,n-1,r+1} \\
& + 12\lambda^2\chi_{m+1,n-1,r+1} - 48\lambda\chi_{m+1,n-1,r+1} + 64\chi_{m+1,n-1,r+1} \\
& + 2\lambda^3\chi_{m+1,n,r-1} - 96\lambda\chi_{m+1,n,r-1} + 256\chi_{m+1,n,r-1} - 4\lambda^3\chi_{m+1,n,r} \\
& - 48\lambda^2\chi_{m+1,n,r} + 1024\chi_{m+1,n,r} + 2\lambda^3\chi_{m+1,n,r+1} - 96\lambda\chi_{m+1,n,r+1} \\
& + 256\chi_{m+1,n,r+1} - \lambda^3\chi_{m+1,n+1,r-1} + 12\lambda^2\chi_{m+1,n+1,r-1} \\
& - 48\lambda\chi_{m+1,n+1,r-1} + 64\chi_{m+1,n+1,r-1} + 2\lambda^3\chi_{m+1,n+1,r} - 96\lambda\chi_{m+1,n+1,r} \\
& + 256\chi_{m+1,n+1,r} - \lambda^3\chi_{m+1,n+1,r+1} + 12\lambda^2\chi_{m+1,n+1,r+1} \\
& - 48\lambda\chi_{m+1,n+1,r+1} + 64\chi_{m+1,n+1,r+1} \Big). \tag{9}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial U_{m,n,r}}{\partial x} = \frac{1}{1152h} & \Big(2(\lambda^2 + 4\lambda - 32)\chi_{m-1,n-1,r} - \lambda^2\chi_{m-1,n-1,r+1} \\
& + 2\lambda^2\chi_{m-1,n,r-1} - 4\lambda^2\chi_{m-1,n,r} + 2\lambda^2\chi_{m-1,n,r+1} - \lambda^2\chi_{m-1,n+1,r-1} \\
& + 2\lambda^2\chi_{m-1,n+1,r} - \lambda^2\chi_{m-1,n+1,r+1} + \lambda^2\chi_{m+1,n-1,r-1} - 2\lambda^2\chi_{m+1,n-1,r} \\
& + \lambda^2\chi_{m+1,n-1,r+1} - 2\lambda^2\chi_{m+1,n,r-1} + 4\lambda^2\chi_{m+1,n,r} - 2\lambda^2\chi_{m+1,n,r+1} \\
& + \lambda^2\chi_{m+1,n+1,r-1} - 2\lambda^2\chi_{m+1,n+1,r} + \lambda^2\chi_{m+1,n+1,r+1} \\
& - (\lambda - 4)^2\chi_{m-1,n-1,r-1} + 8\lambda\chi_{m-1,n-1,r+1} + 8\lambda\chi_{m-1,n,r-1} \\
& - 64\lambda\chi_{m-1,n,r} + 8\lambda\chi_{m-1,n,r+1} + 8\lambda\chi_{m-1,n+1,r-1} + 8\lambda\chi_{m-1,n+1,r} \\
& + 8\lambda\chi_{m-1,n+1,r+1} - 8\lambda\chi_{m+1,n-1,r-1} - 8\lambda\chi_{m+1,n-1,r} - 8\lambda\chi_{m+1,n-1,r+1} \\
& - 8\lambda\chi_{m+1,n,r-1} + 64\lambda\chi_{m+1,n,r} - 8\lambda\chi_{m+1,n,r+1} - 8\lambda\chi_{m+1,n+1,r-1} \\
& - 8\lambda\chi_{m+1,n+1,r} - 8\lambda\chi_{m+1,n+1,r+1} - 16\chi_{m-1,n-1,r+1} - 64\chi_{m-1,n,r-1} \\
& - 256\chi_{m-1,n,r} - 64\chi_{m-1,n,r+1} - 16\chi_{m-1,n+1,r-1} - 64\chi_{m-1,n+1,r} \\
& - 16\chi_{m-1,n+1,r+1} + 16\chi_{m+1,n-1,r-1} + 64\chi_{m+1,n-1,r} + 16\chi_{m+1,n-1,r+1} \\
& + 64\chi_{m+1,n,r-1} + 256\chi_{m+1,n,r} + 64\chi_{m+1,n,r+1} + 16\chi_{m+1,n+1,r-1} \\
& + 64\chi_{m+1,n+1,r} + 16\chi_{m+1,n+1,r+1} \Big),
\end{aligned}$$

$$\begin{aligned}
\frac{\partial U_{m,n,r}}{\partial y} = \frac{1}{1152k} & \Big(2(\lambda^2 + 4\lambda - 32)\chi_{m-1,n-1,r} - \lambda^2\chi_{m-1,n-1,r+1} \\
& + \lambda^2\chi_{m-1,n+1,r-1} - 2\lambda^2\chi_{m-1,n+1,r} + \lambda^2\chi_{m-1,n+1,r+1} + 2\lambda^2\chi_{m,n-1,r-1} \\
& - 4\lambda^2\chi_{m,n-1,r} + 2\lambda^2\chi_{m,n-1,r+1} - 2\lambda^2\chi_{m,n+1,r-1} + 4\lambda^2\chi_{m,n+1,r} \\
& - 2\lambda^2\chi_{m,n+1,r+1} - \lambda^2\chi_{m+1,n-1,r-1} + 2\lambda^2\chi_{m+1,n-1,r} - \lambda^2\chi_{m+1,n-1,r+1} \\
& + \lambda^2\chi_{m+1,n+1,r-1} - 2\lambda^2\chi_{m+1,n+1,r} + \lambda^2\chi_{m+1,n+1,r+1} \\
& - (\lambda - 4)^2\chi_{m-1,n-1,r-1} + 8\lambda\chi_{m-1,n-1,r+1} - 8\lambda\chi_{m-1,n+1,r-1} \\
& - 8\lambda\chi_{m-1,n+1,r} - 8\lambda\chi_{m-1,n+1,r+1} + 8\lambda\chi_{m,n-1,r-1} - 64\lambda\chi_{m,n-1,r} \\
& + 8\lambda\chi_{m,n-1,r+1} - 8\lambda\chi_{m,n+1,r-1} + 64\lambda\chi_{m,n+1,r} - 8\lambda\chi_{m,n+1,r+1} \\
& + 8\lambda\chi_{m+1,n-1,r-1} + 8\lambda\chi_{m+1,n-1,r} + 8\lambda\chi_{m+1,n-1,r+1} - 8\lambda\chi_{m+1,n+1,r-1}
\end{aligned}$$

$$\begin{aligned}
& -8\lambda\chi_{m+1,n+1,r} - 8\lambda\chi_{m+1,n+1,r+1} - 16\chi_{m-1,n-1,r+1} + 16\chi_{m-1,n+1,r-1} \\
& + 64\chi_{m-1,n+1,r} + 16\chi_{m-1,n+1,r+1} - 64\chi_{m,n-1,r-1} - 256\chi_{m,n-1,r} \\
& - 64\chi_{m,n-1,r+1} + 64\chi_{m,n+1,r-1} + 256\chi_{m,n+1,r} + 64\chi_{m,n+1,r+1} \\
& - 16\chi_{m+1,n-1,r-1} - 64\chi_{m+1,n-1,r} - 16\chi_{m+1,n-1,r+1} + 16\chi_{m+1,n+1,r-1} \\
& + 64\chi_{m+1,n+1,r} + 16\chi_{m+1,n+1,r+1} \Big), \\
\frac{\partial U_{m,n,r}}{\partial z} = & \frac{1}{1152q} \Big(2\lambda^2\chi_{m-1,n,r-1} - 2\lambda^2\chi_{m-1,n,r+1} - \lambda^2\chi_{m-1,n+1,r-1} \\
& + \lambda^2\chi_{m-1,n+1,r+1} + 2\lambda^2\chi_{m,n-1,r-1} - 2\lambda^2\chi_{m,n-1,r+1} - 4\lambda^2\chi_{m,n,r-1} \\
& + 4\lambda^2\chi_{m,n,r+1} + 2\lambda^2\chi_{m,n+1,r-1} - 2\lambda^2\chi_{m,n+1,r+1} - \lambda^2\chi_{m+1,n-1,r-1} \\
& + \lambda^2\chi_{m+1,n-1,r+1} + 2\lambda^2\chi_{m+1,n,r-1} - 2\lambda^2\chi_{m+1,n,r+1} - \lambda^2\chi_{m+1,n+1,r-1} \\
& + \lambda^2\chi_{m+1,n+1,r+1} + (\lambda-4)^2\chi_{m-1,n-1,r+1} - (\lambda-4)^2\chi_{m-1,n-1,r-1} \\
& + 8\lambda\chi_{m-1,n,r-1} - 8\lambda\chi_{m-1,n,r+1} + 8\lambda\chi_{m-1,n+1,r-1} - 8\lambda\chi_{m-1,n+1,r+1} \\
& + 8\lambda\chi_{m,n-1,r-1} - 8\lambda\chi_{m,n-1,r+1} - 64\lambda\chi_{m,n,r-1} + 64\lambda\chi_{m,n,r+1} \\
& + 8\lambda\chi_{m,n+1,r-1} - 8\lambda\chi_{m,n+1,r+1} + 8\lambda\chi_{m+1,n-1,r-1} - 8\lambda\chi_{m+1,n-1,r+1} \\
& + 8\lambda\chi_{m+1,n,r-1} - 8\lambda\chi_{m+1,n,r+1} + 8\lambda\chi_{m+1,n+1,r-1} - 8\lambda\chi_{m+1,n+1,r+1} \\
& - 64\chi_{m-1,n,r-1} + 64\chi_{m-1,n,r+1} - 16\chi_{m-1,n+1,r-1} + 16\chi_{m-1,n+1,r+1} \\
& - 64\chi_{m,n-1,r-1} + 64\chi_{m,n-1,r+1} - 256\chi_{m,n,r-1} + 256\chi_{m,n,r+1} \\
& - 64\chi_{m,n+1,r-1} + 64\chi_{m,n+1,r+1} - 16\chi_{m+1,n-1,r-1} + 16\chi_{m+1,n-1,r+1} \\
& - 64\chi_{m+1,n,r-1} + 64\chi_{m+1,n,r+1} - 16\chi_{m+1,n+1,r-1} + 16\chi_{m+1,n+1,r+1} \Big). \\
\frac{\partial^2 U_{m,n,r}}{\partial x \partial y} = & -\frac{1}{96hk} \Big((\lambda-4)\chi_{m-1,n-1,r-1} - 2(\lambda+8)\chi_{m-1,n-1,r} \\
& + \lambda\chi_{m-1,n-1,r+1} - \lambda\chi_{m-1,n+1,r-1} + 2\lambda\chi_{m-1,n+1,r} - \lambda\chi_{m-1,n+1,r+1} \\
& - \lambda\chi_{m+1,n-1,r-1} + 2\lambda\chi_{m+1,n-1,r} - \lambda\chi_{m+1,n-1,r+1} + \lambda\chi_{m+1,n+1,r-1} \\
& - 2\lambda\chi_{m+1,n+1,r} + \lambda\chi_{m+1,n+1,r+1} - 4\chi_{m-1,n-1,r+1} + 4\chi_{m-1,n+1,r-1} \\
& + 16\chi_{m-1,n+1,r} + 4\chi_{m-1,n+1,r+1} + 4\chi_{m+1,n-1,r-1} + 16\chi_{m+1,n-1,r} \\
& + 4\chi_{m+1,n-1,r+1} - 4\chi_{m+1,n+1,r-1} - 16\chi_{m+1,n+1,r} - 4\chi_{m+1,n+1,r+1} \Big), \\
\frac{\partial^2 U_{m,n,r}}{\partial x \partial z} = & \frac{1}{96hs} \Big(-(\lambda-4)\chi_{m-1,n-1,r-1} + (\lambda-4)\chi_{m-1,n-1,r+1} \\
& + 2\lambda\chi_{m-1,n,r-1} - 2\lambda\chi_{m-1,n,r+1} - \lambda\chi_{m-1,n+1,r-1} + \lambda\chi_{m-1,n+1,r+1} \\
& + \lambda\chi_{m+1,n-1,r-1} - \lambda\chi_{m+1,n-1,r+1} - 2\lambda\chi_{m+1,n,r-1} + 2\lambda\chi_{m+1,n,r+1} \\
& + \lambda\chi_{m+1,n+1,r-1} - \lambda\chi_{m+1,n+1,r+1} + 16\chi_{m-1,n,r-1} - 16\chi_{m-1,n,r+1} \\
& + 4\chi_{m-1,n+1,r-1} - 4\chi_{m-1,n+1,r+1} - 4\chi_{m+1,n-1,r-1} + 4\chi_{m+1,n-1,r+1} \\
& - 16\chi_{m+1,n,r-1} + 16\chi_{m+1,n,r+1} - 4\chi_{m+1,n+1,r-1} + 4\chi_{m+1,n+1,r+1} \Big), \\
\frac{\partial^2 U_{m,n,r}}{\partial y \partial z} = & \frac{1}{96kq} \Big(-(\lambda-4)\chi_{m-1,n-1,r-1} + (\lambda-4)\chi_{m-1,n-1,r+1} \\
& + \lambda\chi_{m-1,n+1,r-1} - \lambda\chi_{m-1,n+1,r+1} + 2\lambda\chi_{m,n-1,r-1} - 2\lambda\chi_{m,n-1,r+1} \\
& - 2\lambda\chi_{m,n+1,r-1} + 2\lambda\chi_{m,n+1,r+1} - \lambda\chi_{m+1,n-1,r-1} + \lambda\chi_{m+1,n-1,r+1}
\end{aligned}
\tag{10}$$

$$\begin{aligned}
& + \lambda \chi_{m+1,n+1,r-1} - \lambda \chi_{m+1,n+1,r+1} - 4\chi_{m-1,n+1,r-1} + 4\chi_{m-1,n+1,r+1} \\
& + 16\chi_{m,n-1,r-1} - 16\chi_{m,n-1,r+1} - 16\chi_{m,n+1,r-1} + 16\chi_{m,n+1,r+1} \\
& + 4\chi_{m+1,n-1,r-1} - 4\chi_{m+1,n-1,r+1} - 4\chi_{m+1,n+1,r-1} + 4\chi_{m+1,n+1,r+1} \Big), \\
\frac{\partial^3 U_{m,n,r}}{\partial x \partial y \partial z} &= \frac{1}{8hkq} \Big(-\chi_{m-1,n-1,r-1} + \chi_{m-1,n-1,r+1} + \chi_{m-1,n+1,r-1} \\
& - \chi_{m-1,n+1,r+1} + \chi_{m+1,n-1,r-1} - \chi_{m+1,n-1,r+1} \\
& - \chi_{m+1,n+1,r-1} + \chi_{m+1,n+1,r+1} \Big), \\
& \vdots \\
\frac{\partial^2 U_{m,n,r}}{\partial x^2} &= \frac{\lambda + 2}{1152h^2} \Big(\chi_{m-1,n-1,r-1}(\lambda - 4)^2 - 2(\lambda^2 + 4\lambda - 32) \\
& \chi_{m-1,n-1,r} + \lambda^2 \chi_{m-1,n-1,r+1} - 8\lambda \chi_{m-1,n-1,r+1} + 16\chi_{m-1,n-1,r+1} \\
& - 2\lambda^2 \chi_{m-1,n,r-1} - 8\lambda \chi_{m-1,n,r-1} + 64\chi_{m-1,n,r-1} + 4\lambda^2 \chi_{m-1,n,r} \\
& + 64\lambda \chi_{m-1,n,r} + 256\chi_{m-1,n,r} - 2\lambda^2 \chi_{m-1,n,r+1} - 8\lambda \chi_{m-1,n,r+1} \\
& + 64\chi_{m-1,n,r+1} + \lambda^2 \chi_{m-1,n+1,r-1} - 8\lambda \chi_{m-1,n+1,r-1} + 16\chi_{m-1,n+1,r-1} \\
& - 2\lambda^2 \chi_{m-1,n+1,r} - 8\lambda \chi_{m-1,n+1,r} + 64\chi_{m-1,n+1,r} + \lambda^2 \chi_{m-1,n+1,r+1} \\
& - 8\lambda \chi_{m-1,n+1,r+1} + 16\chi_{m-1,n+1,r+1} - 2\lambda^2 \chi_{m,n-1,r-1} + 16\lambda \chi_{m,n-1,r-1} \\
& - 32\chi_{m,n-1,r-1} + 4\lambda^2 \chi_{m,n-1,r} + 16\lambda \chi_{m,n-1,r} - 128\chi_{m,n-1,r} \\
& - 2\lambda^2 \chi_{m,n-1,r+1} + 16\lambda \chi_{m,n-1,r+1} - 32\chi_{m,n-1,r+1} + 4\lambda^2 \chi_{m,n,r-1} \\
& + 16\lambda \chi_{m,n,r-1} - 128\chi_{m,n,r-1} - 8\lambda^2 \chi_{m,n,r} - 128\lambda \chi_{m,n,r} - 512\chi_{m,n,r} \\
& + 4\lambda^2 \chi_{m,n,r+1} + 16\lambda \chi_{m,n,r+1} - 128\chi_{m,n,r+1} - 2\lambda^2 \chi_{m,n+1,r-1} \\
& + 16\lambda \chi_{m,n+1,r-1} - 32\chi_{m,n+1,r-1} + 4\lambda^2 \chi_{m,n+1,r} + 16\lambda \chi_{m,n+1,r} \\
& - 128\chi_{m,n+1,r} - 2\lambda^2 \chi_{m,n+1,r+1} + 16\lambda \chi_{m,n+1,r+1} - 32\chi_{m,n+1,r+1} \\
& + \lambda^2 \chi_{m+1,n-1,r-1} - 8\lambda \chi_{m+1,n-1,r-1} + 16\chi_{m+1,n-1,r-1} - 2\lambda^2 \chi_{m+1,n-1,r} \\
& - 8\lambda \chi_{m+1,n-1,r} + 64\chi_{m+1,n-1,r} + \lambda^2 \chi_{m+1,n-1,r+1} - 8\lambda \chi_{m+1,n-1,r+1} \\
& + 16\chi_{m+1,n-1,r+1} - 2\lambda^2 \chi_{m+1,n,r-1} - 8\lambda \chi_{m+1,n,r-1} + 64\chi_{m+1,n,r-1} \\
& + 4\lambda^2 \chi_{m+1,n,r} + 64\lambda \chi_{m+1,n,r} + 256\chi_{m+1,n,r} - 2\lambda^2 \chi_{m+1,n,r+1} \\
& - 8\lambda \chi_{m+1,n,r+1} + 64\chi_{m+1,n,r+1} + \lambda^2 \chi_{m+1,n+1,r-1} - 8\lambda \chi_{m+1,n+1,r-1} \\
& + 16\chi_{m+1,n+1,r-1} - 2\lambda^2 \chi_{m+1,n+1,r} - 8\lambda \chi_{m+1,n+1,r} + 64\chi_{m+1,n+1,r} \\
& + \lambda^2 \chi_{m+1,n+1,r+1} - 8\lambda \chi_{m+1,n+1,r+1} + 16\chi_{m+1,n+1,r+1} \Big), \\
\frac{\partial^2 U_{m,n,r}}{\partial y^2} &= \frac{\lambda + 2}{1152k^2} \Big(\chi_{m-1,n-1,r-1}(\lambda - 4)^2 - 2(\lambda^2 + 4\lambda - 32) \\
& \chi_{m-1,n-1,r} + \lambda^2 \chi_{m-1,n-1,r+1} - 8\lambda \chi_{m-1,n-1,r+1} + 16\chi_{m-1,n-1,r+1} \\
& - 2\lambda^2 \chi_{m-1,n,r-1} + 16\lambda \chi_{m-1,n,r-1} - 32\chi_{m-1,n,r-1} + 4\lambda^2 \chi_{m-1,n,r} \\
& + 16\lambda \chi_{m-1,n,r} - 128\chi_{m-1,n,r} - 2\lambda^2 \chi_{m-1,n,r+1} + 16\lambda \chi_{m-1,n,r+1} \\
& - 32\chi_{m-1,n,r+1} + \lambda^2 \chi_{m-1,n+1,r-1} - 8\lambda \chi_{m-1,n+1,r-1} + 16\chi_{m-1,n+1,r-1}
\end{aligned}
\tag{11}$$

$$\begin{aligned}
& -2\lambda^2\chi_{m-1,n+1,r} - 8\lambda\chi_{m-1,n+1,r} + 64\chi_{m-1,n+1,r} + \lambda^2\chi_{m-1,n+1,r+1} \\
& - 8\lambda\chi_{m-1,n+1,r+1} + 16\chi_{m-1,n+1,r+1} - 2\lambda^2\chi_{m,n-1,r-1} - 8\lambda\chi_{m,n-1,r-1} \\
& + 64\chi_{m,n-1,r-1} + 4\lambda^2\chi_{m,n-1,r} + 64\lambda\chi_{m,n-1,r} + 256\chi_{m,n-1,r} \\
& - 2\lambda^2\chi_{m,n-1,r+1} - 8\lambda\chi_{m,n-1,r+1} + 64\chi_{m,n-1,r+1} + 4\lambda^2\chi_{m,n,r-1} \\
& + 16\lambda\chi_{m,n,r-1} - 128\chi_{m,n,r-1} - 8\lambda^2\chi_{m,n,r} - 128\lambda\chi_{m,n,r} - 512\chi_{m,n,r} \\
& + 4\lambda^2\chi_{m,n,r+1} + 16\lambda\chi_{m,n,r+1} - 128\chi_{m,n,r+1} - 2\lambda^2\chi_{m,n+1,r-1} \\
& - 8\lambda\chi_{m,n+1,r-1} + 64\chi_{m,n+1,r-1} + 4\lambda^2\chi_{m,n+1,r} + 64\lambda\chi_{m,n+1,r} \\
& + 256\chi_{m,n+1,r} - 2\lambda^2\chi_{m,n+1,r+1} - 8\lambda\chi_{m,n+1,r+1} + 64\chi_{m,n+1,r+1} \\
& + \lambda^2\chi_{m+1,n-1,r-1} - 8\lambda\chi_{m+1,n-1,r-1} + 16\chi_{m+1,n-1,r-1} \\
& - 2\lambda^2\chi_{m+1,n-1,r} - 8\lambda\chi_{m+1,n-1,r} + 64\chi_{m+1,n-1,r} + \lambda^2\chi_{m+1,n-1,r+1} \\
& - 8\lambda\chi_{m+1,n-1,r+1} + 16\chi_{m+1,n-1,r+1} - 2\lambda^2\chi_{m+1,n,r-1} + 16\lambda\chi_{m+1,n,r-1} \\
& - 32\chi_{m+1,n,r-1} + 4\lambda^2\chi_{m+1,n,r} + 16\lambda\chi_{m+1,n,r} - 128\chi_{m+1,n,r} \\
& - 2\lambda^2\chi_{m+1,n,r+1} + 16\lambda\chi_{m+1,n,r+1} - 32\chi_{m+1,n,r+1} + \lambda^2\chi_{m+1,n+1,r-1} \\
& - 8\lambda\chi_{m+1,n+1,r-1} + 16\chi_{m+1,n+1,r-1} - 2\lambda^2\chi_{m+1,n+1,r} - 8\lambda\chi_{m+1,n+1,r} \\
& + 64\chi_{m+1,n+1,r} + \lambda^2\chi_{m+1,n+1,r+1} - 8\lambda\chi_{m+1,n+1,r+1} + 16\chi_{m+1,n+1,r+1} \Big), \\
& \frac{\partial^2 U_{m,n,r}}{\partial z^2} = \frac{\lambda+2}{1152q^2} \Big(\chi_{m-1,n-1,r-1}(\lambda-4)^2 - 2\chi_{m-1,n-1,r}(\lambda-4)^2 \\
& + \lambda^2\chi_{m-1,n-1,r+1} - 8\lambda\chi_{m-1,n-1,r+1} + 16\chi_{m-1,n-1,r+1} \\
& - 2\lambda^2\chi_{m-1,n,r-1} - 8\lambda\chi_{m-1,n,r-1} + 64\chi_{m-1,n,r-1} + 4\lambda^2\chi_{m-1,n,r} \\
& + 16\lambda\chi_{m-1,n,r} - 128\chi_{m-1,n,r} - 2\lambda^2\chi_{m-1,n,r+1} - 8\lambda\chi_{m-1,n,r+1} \\
& + 64\chi_{m-1,n,r+1} + \lambda^2\chi_{m-1,n+1,r-1} - 8\lambda\chi_{m-1,n+1,r-1} + 16\chi_{m-1,n+1,r-1} \\
& - 2\lambda^2\chi_{m-1,n+1,r} + 16\lambda\chi_{m-1,n+1,r} - 32\chi_{m-1,n+1,r} + \lambda^2\chi_{m-1,n+1,r+1} \\
& - 8\lambda\chi_{m-1,n+1,r+1} + 16\chi_{m-1,n+1,r+1} - 2\lambda^2\chi_{m,n-1,r-1} - 8\lambda\chi_{m,n-1,r-1} \\
& + 64\chi_{m,n-1,r-1} + 4\lambda^2\chi_{m,n-1,r} + 16\lambda\chi_{m,n-1,r} - 128\chi_{m,n-1,r} \\
& - 2\lambda^2\chi_{m,n-1,r+1} - 8\lambda\chi_{m,n-1,r+1} + 64\chi_{m,n-1,r+1} + 4\lambda^2\chi_{m,n,r-1} \\
& + 64\lambda\chi_{m,n,r-1} + 256\chi_{m,n,r-1} - 8\lambda^2\chi_{m,n,r} - 128\lambda\chi_{m,n,r} - 512\chi_{m,n,r} \\
& + 4\lambda^2\chi_{m,n,r+1} + 64\lambda\chi_{m,n,r+1} + 256\chi_{m,n,r+1} - 2\lambda^2\chi_{m,n+1,r-1} \\
& - 8\lambda\chi_{m,n+1,r-1} + 64\chi_{m,n+1,r-1} + 4\lambda^2\chi_{m,n+1,r} + 16\lambda\chi_{m,n+1,r} \\
& - 128\chi_{m,n+1,r} - 2\lambda^2\chi_{m,n+1,r+1} - 8\lambda\chi_{m,n+1,r+1} + 64\chi_{m,n+1,r+1} \\
& + \lambda^2\chi_{m+1,n-1,r-1} - 8\lambda\chi_{m+1,n-1,r-1} + 16\chi_{m+1,n-1,r-1} \\
& - 2\lambda^2\chi_{m+1,n-1,r} + 16\lambda\chi_{m+1,n-1,r} - 32\chi_{m+1,n-1,r} + \lambda^2\chi_{m+1,n-1,r+1} \\
& - 8\lambda\chi_{m+1,n-1,r+1} + 16\chi_{m+1,n-1,r+1} - 2\lambda^2\chi_{m+1,n,r-1} - 8\lambda\chi_{m+1,n,r-1} \\
& + 64\chi_{m+1,n,r-1} + 4\lambda^2\chi_{m+1,n,r} + 16\lambda\chi_{m+1,n,r} - 128\chi_{m+1,n,r} \\
& - 2\lambda^2\chi_{m+1,n,r+1} - 8\lambda\chi_{m+1,n,r+1} + 64\chi_{m+1,n,r+1} + \lambda^2\chi_{m+1,n+1,r-1} \\
& - 8\lambda\chi_{m+1,n+1,r-1} + 16\chi_{m+1,n+1,r-1} - 2\lambda^2\chi_{m+1,n+1,r} + 16\lambda\chi_{m+1,n+1,r} \\
& - 32\chi_{m+1,n+1,r} + \lambda^2\chi_{m+1,n+1,r+1} - 8\lambda\chi_{m+1,n+1,r+1} + 16\chi_{m+1,n+1,r+1} \Big). \tag{12}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial^3 U_{m,n,r}}{\partial x^2 \partial y} &= \frac{\lambda+2}{96h^2k} \left((\lambda-4)\chi_{m-1,n-1,r-1} - 2(\lambda+8)\chi_{m-1,n-1,r} \right. \\
&+ \lambda\chi_{m-1,n-1,r+1} - \lambda\chi_{m-1,n+1,r-1} + 2\lambda\chi_{m-1,n+1,r} - \lambda\chi_{m-1,n+1,r+1} \\
&- 2\lambda\chi_{m,n-1,r-1} + 4\lambda\chi_{m,n-1,r} - 2\lambda\chi_{m,n-1,r+1} + 2\lambda\chi_{m,n+1,r-1} \\
&- 4\lambda\chi_{m,n+1,r} + 2\lambda\chi_{m,n+1,r+1} + \lambda\chi_{m+1,n-1,r-1} - 2\lambda\chi_{m+1,n-1,r} \\
&+ \lambda\chi_{m+1,n-1,r+1} - \lambda\chi_{m+1,n+1,r-1} + 2\lambda\chi_{m+1,n+1,r} - \lambda\chi_{m+1,n+1,r+1} \\
&- 4\chi_{m-1,n-1,r+1} + 4\chi_{m-1,n+1,r-1} + 16\chi_{m-1,n+1,r} + 4\chi_{m-1,n+1,r+1} \\
&+ 8\chi_{m,n-1,r-1} + 32\chi_{m,n-1,r} + 8\chi_{m,n-1,r+1} - 8\chi_{m,n+1,r-1} \\
&- 32\chi_{m,n+1,r} - 8\chi_{m,n+1,r+1} - 4\chi_{m+1,n-1,r-1} - 16\chi_{m+1,n-1,r} \\
&- 4\chi_{m+1,n-1,r+1} + 4\chi_{m+1,n+1,r-1} + 16\chi_{m+1,n+1,r} + 4\chi_{m+1,n+1,r+1} \left. \right), \\
\frac{\partial^3 U_{m,n,r}}{\partial x^2 \partial z} &= \frac{\lambda+2}{96h^2q} \left((\lambda-4)\chi_{m-1,n-1,r-1} - (\lambda-4)\chi_{m-1,n-1,r+1} \right. \\
&- 2\lambda\chi_{m-1,n,r-1} + 2\lambda\chi_{m-1,n,r+1} + \lambda\chi_{m-1,n+1,r-1} - \lambda\chi_{m-1,n+1,r+1} \\
&- 2\lambda\chi_{m,n-1,r-1} + 2\lambda\chi_{m,n-1,r+1} + 4\lambda\chi_{m,n,r-1} - 4\lambda\chi_{m,n,r+1} \\
&- 2\lambda\chi_{m,n+1,r-1} + 2\lambda\chi_{m,n+1,r+1} + \lambda\chi_{m+1,n-1,r-1} - \lambda\chi_{m+1,n-1,r+1} \\
&- 2\lambda\chi_{m+1,n,r-1} + 2\lambda\chi_{m+1,n,r+1} + \lambda\chi_{m+1,n+1,r-1} - \lambda\chi_{m+1,n+1,r+1} \\
&- 16\chi_{m-1,n,r-1} + 16\chi_{m-1,n,r+1} - 4\chi_{m-1,n+1,r-1} + 4\chi_{m-1,n+1,r+1} \\
&+ 8\chi_{m,n-1,r-1} - 8\chi_{m,n-1,r+1} + 32\chi_{m,n,r-1} - 32\chi_{m,n,r+1} \\
&+ 8\chi_{m,n+1,r-1} - 8\chi_{m,n+1,r+1} - 4\chi_{m+1,n-1,r-1} + 4\chi_{m+1,n-1,r+1} \\
&- 16\chi_{m+1,n,r-1} + 16\chi_{m+1,n,r+1} - 4\chi_{m+1,n+1,r-1} + 4\chi_{m+1,n+1,r+1} \left. \right), \\
\frac{\partial^3 U_{m,n,r}}{\partial y^2 \partial z} &= \frac{\lambda+2}{96k^2q} \left((\lambda-4)\chi_{m-1,n-1,r-1} - (\lambda-4)\chi_{m-1,n-1,r+1} \right. \\
&- 2\lambda\chi_{m-1,n,r-1} + 2\lambda\chi_{m-1,n,r+1} + \lambda\chi_{m-1,n+1,r-1} - \lambda\chi_{m-1,n+1,r+1} \\
&- 2\lambda\chi_{m,n-1,r-1} + 2\lambda\chi_{m,n-1,r+1} + 4\lambda\chi_{m,n,r-1} - 4\lambda\chi_{m,n,r+1} \\
&- 2\lambda\chi_{m,n+1,r-1} + 2\lambda\chi_{m,n+1,r+1} + \lambda\chi_{m+1,n-1,r-1} - \lambda\chi_{m+1,n-1,r+1} \\
&- 2\lambda\chi_{m+1,n,r-1} + 2\lambda\chi_{m+1,n,r+1} + \lambda\chi_{m+1,n+1,r-1} - \lambda\chi_{m+1,n+1,r+1} \\
&+ 8\chi_{m-1,n,r-1} - 8\chi_{m-1,n,r+1} - 4\chi_{m-1,n+1,r-1} + 4\chi_{m-1,n+1,r+1} \\
&- 16\chi_{m,n-1,r-1} + 16\chi_{m,n-1,r+1} + 32\chi_{m,n,r-1} - 32\chi_{m,n,r+1} \\
&- 16\chi_{m,n+1,r-1} + 16\chi_{m,n+1,r+1} - 4\chi_{m+1,n-1,r-1} + 4\chi_{m+1,n-1,r+1} \\
&+ 8\chi_{m+1,n,r-1} - 8\chi_{m+1,n,r+1} - 4\chi_{m+1,n+1,r-1} + 4\chi_{m+1,n+1,r+1} \left. \right), \\
\frac{\partial^3 U_{m,n,r}}{\partial y \partial z^2} &= \frac{\lambda+2}{96kq^2} \left((\lambda-4)\chi_{m-1,n-1,r-1} - 2(\lambda-4)\chi_{m-1,n-1,r} \right. \\
&+ \lambda\chi_{m-1,n-1,r+1} - \lambda\chi_{m-1,n+1,r-1} + 2\lambda\chi_{m-1,n+1,r} - \lambda\chi_{m-1,n+1,r+1} \\
&- 2\lambda\chi_{m,n-1,r-1} + 4\lambda\chi_{m,n-1,r} - 2\lambda\chi_{m,n-1,r+1} + 2\lambda\chi_{m,n+1,r-1} \\
&- 4\lambda\chi_{m,n+1,r} + 2\lambda\chi_{m,n+1,r+1} + \lambda\chi_{m+1,n-1,r-1} - 2\lambda\chi_{m+1,n-1,r} \\
&+ \lambda\chi_{m+1,n-1,r+1} - \lambda\chi_{m+1,n+1,r-1} + 2\lambda\chi_{m+1,n+1,r} - \lambda\chi_{m+1,n+1,r+1} \\
&- 4\chi_{m-1,n-1,r+1} + 4\chi_{m-1,n+1,r-1} - 8\chi_{m-1,n+1,r} + 4\chi_{m-1,n+1,r+1} \\
&- 16\chi_{m,n-1,r-1} + 32\chi_{m,n-1,r} - 16\chi_{m,n-1,r+1} + 16\chi_{m,n+1,r-1} \\
&- 32\chi_{m,n+1,r} + 16\chi_{m,n+1,r+1} - 4\chi_{m+1,n-1,r-1} + 8\chi_{m+1,n-1,r} \left. \right)
\end{aligned}$$

$$\begin{aligned}
& -4\chi_{m+1,n-1,r+1} + 4\chi_{m+1,n+1,r-1} - 8\chi_{m+1,n+1,r} + 4\chi_{m+1,n+1,r+1} \Big), \\
& \frac{\partial^3 U_{m,n,r}}{\partial x \partial z^2} = \frac{\lambda+2}{96hq^2} \Big((\lambda-4)\chi_{m-1,n-1,r-1} - 2(\lambda-4)\chi_{m-1,n-1,r} \\
& + \lambda\chi_{m-1,n-1,r+1} - 2\lambda\chi_{m-1,n,r-1} + 4\lambda\chi_{m-1,n,r} - 2\lambda\chi_{m-1,n,r+1} \\
& + \lambda\chi_{m-1,n+1,r-1} - 2\lambda\chi_{m-1,n+1,r} + \lambda\chi_{m-1,n+1,r+1} - \lambda\chi_{m+1,n-1,r-1} \\
& + 2\lambda\chi_{m+1,n-1,r} - \lambda\chi_{m+1,n-1,r+1} + 2\lambda\chi_{m+1,n,r-1} - 4\lambda\chi_{m+1,n,r} \\
& + 2\lambda\chi_{m+1,n,r+1} - \lambda\chi_{m+1,n+1,r-1} + 2\lambda\chi_{m+1,n+1,r} - \lambda\chi_{m+1,n+1,r+1} \\
& - 4\chi_{m-1,n-1,r+1} - 16\chi_{m-1,n,r-1} + 32\chi_{m-1,n,r} - 16\chi_{m-1,n,r+1} \\
& - 4\chi_{m-1,n+1,r-1} + 8\chi_{m-1,n+1,r} - 4\chi_{m-1,n+1,r+1} + 4\chi_{m+1,n-1,r-1} \\
& - 8\chi_{m+1,n-1,r} + 4\chi_{m+1,n-1,r+1} + 16\chi_{m+1,n,r-1} - 32\chi_{m+1,n,r} \\
& + 16\chi_{m+1,n,r+1} + 4\chi_{m+1,n+1,r-1} - 8\chi_{m+1,n+1,r} + 4\chi_{m+1,n+1,r+1} \Big), \quad (13) \\
& \frac{\partial^3 U_{m,n,r}}{\partial x \partial y^2} = \frac{\lambda+2}{96hk^2} \Big((\lambda-4)\chi_{m-1,n-1,r-1} - 2(\lambda+8)\chi_{m-1,n-1,r} \\
& + \lambda\chi_{m-1,n-1,r+1} - 2\lambda\chi_{m-1,n,r-1} + 4\lambda\chi_{m-1,n,r} - 2\lambda\chi_{m-1,n,r+1} \\
& + \lambda\chi_{m-1,n+1,r-1} - 2\lambda\chi_{m-1,n+1,r} + \lambda\chi_{m-1,n+1,r+1} - \lambda\chi_{m+1,n-1,r-1} \\
& + 2\lambda\chi_{m+1,n-1,r} - \lambda\chi_{m+1,n-1,r+1} + 2\lambda\chi_{m+1,n,r-1} - 4\lambda\chi_{m+1,n,r} \\
& + 2\lambda\chi_{m+1,n,r+1} - \lambda\chi_{m+1,n+1,r-1} + 2\lambda\chi_{m+1,n+1,r} - \lambda\chi_{m+1,n+1,r+1} \\
& - 4\chi_{m-1,n-1,r+1} + 8\chi_{m-1,n,r-1} + 32\chi_{m-1,n,r} + 8\chi_{m-1,n,r+1} \\
& - 4\chi_{m-1,n+1,r-1} - 16\chi_{m-1,n+1,r} - 4\chi_{m-1,n+1,r+1} + 4\chi_{m+1,n-1,r-1} \\
& + 16\chi_{m+1,n-1,r} + 4\chi_{m+1,n-1,r+1} - 8\chi_{m+1,n,r-1} - 32\chi_{m+1,n,r} \\
& - 8\chi_{m+1,n,r+1} + 4\chi_{m+1,n+1,r-1} + 16\chi_{m+1,n+1,r} + 4\chi_{m+1,n+1,r+1} \Big), \\
& \vdots
\end{aligned}$$

In all n-dimensions PDE's with collocation method we get a system of algebraic equations in this form

$$A\chi = b, \quad (14)$$

We solve the above system using newton's method to find the unknown values of χ .

3 The error estimates

Lemma 1 Suppose that \hat{U} is an estimation of smoothness class C^2 . At that point error gauges of the insertion on a square work of side h are

$$\begin{aligned}
\|U - \hat{U}\| &\leq \beta_0 h^4, \quad \left\| \frac{\partial U}{\partial x} - \frac{\partial \hat{U}}{\partial x} \right\| \leq \beta_1 h^3, \quad \left\| \frac{\partial U}{\partial z} - \frac{\partial \hat{U}}{\partial z} \right\| \leq \beta_2 h^3, \quad \left\| \frac{\partial U}{\partial y} - \frac{\partial \hat{U}}{\partial y} \right\| \leq \beta_3 h^3, \\
\left\| \frac{\partial^2 U}{\partial x^2} - \frac{\partial^2 \hat{U}}{\partial x^2} \right\| &\leq \beta_4 h^2, \quad \left\| \frac{\partial^2 U}{\partial y^2} - \frac{\partial^2 \hat{U}}{\partial y^2} \right\| \leq \beta_5 h^2, \quad \left\| \frac{\partial^2 U}{\partial z^2} - \frac{\partial^2 \hat{U}}{\partial z^2} \right\| \leq \beta_6 h^2,
\end{aligned}$$

$$\left\| \frac{\partial^2 U}{\partial x \partial y} - \frac{\partial^2 \hat{U}}{\partial x \partial y} \right\| \leq \beta_7 h^2, \quad \left\| \frac{\partial^2 U}{\partial x \partial z} - \frac{\partial^2 \hat{U}}{\partial x \partial z} \right\| \leq \beta_8 h^2, \quad \left\| \frac{\partial^2 U}{\partial y \partial z} - \frac{\partial^2 \hat{U}}{\partial y \partial z} \right\| \leq \beta_9 h^2,$$

where the β_i are constants.

The proof of above lemma see [9].

4 The numerical results

Presently, we must know whether this method, which was developed by presenting its constructions in different dimensions, is accurate and effective or not. To prove that this method is of high accuracy, we present in this section various numerical examples in different dimensions. We also show some figures of the results obtained. In addition to providing comparisons of our results with pre-existing results.

The first test problem: [18]

We take the first test problem in the 2-dimensional in this form:

$$u_{xx}(x, y) + u_{yy}(x, y) + u_x(x, y) + u_y(x, y) - 3e^{2x+3y}(x^2(18y^2 - 4y - 5) + x(5 - 8y^2 - 6y)) - 3y^2 + 3y = 0, \quad x, y \in [a, b] \quad (15)$$

The exact solution to that problem given as follows:

$$u(x, y) = 3e^{2x+3y}(x - x^2)(y - y^2). \quad (16)$$

We take the boundary conditions to the first problem in this form:

$$u(a, y) = u(x, a) = \alpha, \quad u(b, y) = u(x, b) = \beta. \quad (17)$$

By substitution from (4)-(6) into (15) with (17) we obtain the numerical results as in the next table:

Table 1: The computational results to the first problem at $y = 0.5, x, y \in [0, 1]$

x	Numerical results	Exact results	Absolute error	Quadratic B-Spline [18]
0.1	0.36856	0.36949	9.22844 E-4	1.06992 E-3
0.2	0.80015	0.80230	2.15026 E-3	2.32385 E-3
0.3	1.28295	1.28617	3.21794 E-3	3.40917 E-3
0.4	1.79122	1.79535	4.12938 E-3	4.31609 E-3
0.5	2.27931	2.28422	4.90872 E-3	5.04294 E-3
0.6	2.67274	2.67835	5.61237 E-3	5.60652 E-3
0.7	2.85609	2.86243	6.34146 E-3	6.05466 E-3
0.8	2.65652	2.66375	7.23762 E-3	6.46809 E-3
0.9	1.82167	1.83010	8.43355 E-3	6.93102 E-3

In Table 1, we compared the results about the 2-dimensions Extended cubic B-spline method employing a work of 50×50 and the exact results together. We show that our results are accepted with regard to the exact results. In Fig. 1, we introduce the numerical arrangements with the exact solution at $y = 0.5$. In Fig. 2, we show the numerical results and the exact solution at $x = 0.5$.




Figure 1: The numerical results with the exact results at $y = 0.5$.




Figure 2: The numerical results with the exact results at $x = 0.5$.

The second test problem: MHD duct flow [7–9, 16]

The cross-section of an infinitely long rectangular duct is oriented with its sides parallel to the x - and y -axes and the origin of coordinates at its center. The duct width is $2a$ and height $2b$ so that the sides of the duct have equations $x = \pm a$ and $y = \pm b$. A conducting fluid flows in the z direction along the duct and is subjected to a constant applied magnetic field M acting in a direction lying in the xy -plane and making an angle ϕ with the y -axis. The equations governing the flow may be expressed in the normalized form [7, 15].

$$\frac{\partial P}{\partial z} = \mu\nu\nabla^2\nabla_z + \frac{A_{0x}}{\mu_0}\frac{\partial P_z}{\partial x'} + \frac{A_{0y}}{\mu_0}\frac{\partial P_z}{\partial y'}, \quad (18)$$

and the z -component of the curl of Ohm's law,

$$\nabla^2 A_z + \xi\mu_0(A_{0x}\frac{\partial U_z}{\partial x'} + A_{0y}\frac{\partial U_z}{\partial y'}) = 0, \quad (19)$$

with the boundary conditions: $U = A = 0$ at $x' = \pm a, y' = \pm b$, where ν , μ and ξ are, respectively, the kinematic viscosity, density and electric conductivity of the fluid; μ_0 is the magnetic permeability in vacuum; dP/dz is the constant axial pressure gradient; B_{0x} and B_{0y} are the x' and y' components of the applied magnetic field; and U_z and A_z are the z components of velocity

and induced magnetic field, respectively. Following the notation of P. C. Lu [15], who solved this problem using the Kantorovich method, Eqs. (18) and (19) become in non-dimensionized form,

$$(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})U + M_x \frac{\partial A}{\partial x} + M_y \frac{\partial A}{\partial y} = -1, \quad (20)$$

and

$$(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})A + M_x \frac{\partial U}{\partial x} + M_y \frac{\partial U}{\partial y} = -1, \quad (21)$$

with boundary conditions $U = A = 0, x = \pm\alpha, y = \pm 1$. Distance has been scaled to the duct semi-height b so that $x = x'/b, y = y'/b$, and $\alpha = a/b$. The following normalisations have also been used.

$$\begin{aligned} U &= \frac{U_z}{\frac{-b^2}{\nu\mu} \frac{dP}{dz}}, \\ A &= \frac{A_z}{\frac{-b^2}{\nu\mu} \frac{dP}{dz} \mu_0 (\nu\mu\xi)^{\frac{1}{2}}}, \\ M_x &= A_{0x'} b \left(\frac{\xi}{\nu\mu} \right)^{\frac{1}{2}} = M \sin(\phi), \\ M_y &= A_{0y'} b \left(\frac{\xi}{\nu\mu} \right)^{\frac{1}{2}} = M \cos(\phi), \\ M &= \text{Hartmann no.} = (M_x^2 + M_y^2)^{\frac{1}{2}} = A_0 b \left(\frac{\xi}{\nu\mu} \right)^{\frac{1}{2}}. \end{aligned} \quad (22)$$

The Hartmann number is the ratio of magnetic to fluid viscosity. If $M = 0$, the flow field is the classical laminar pipe flow. If $M \geq 1$, the flow field is determined primarily by the $E \times A$ drift. To uncouple (20) and (21), the functions

$$H_1 = U + A, \quad (23)$$

and

$$H_2 = U - A, \quad (24)$$

$$(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})H_1 + M_x \frac{\partial H_1}{\partial x} + M_y \frac{\partial H_1}{\partial y} = -1, \quad (25)$$

and

$$(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})H_2 - M_x \frac{\partial H_2}{\partial x} - M_y \frac{\partial H_2}{\partial y} = -1, \quad (26)$$

with boundary conditions $H_1 = H_2 = 0, x = \pm\alpha, y = \pm 1$. Thus, if H_1 is solved as $H_1(M_x, M_y)$ from (26), then

$$H_2(M_x, M_y) = H_1(-M_x, -M_y). \quad (27)$$

So that the solution is completely determined when either H_1 or H_2 , are known. Having determined H_1 the function H_2 is found from (27) and hence the velocity field U from

$$U = \frac{1}{2}(H_1 + H_2). \quad (28)$$

Now, we will introduce some numerical results for the flow in a square duct with an applied magnetic field parallel to the x -axis so that $M_y = 0$. To compare with earlier results [7–9, 14], we give to M , the following values $M_x = 0, 2, 5$ and 8.

By substituting from (4)-(6) in (25) and (26) we get the numerical solutions as follows:

Table 2: U at the centre of the duct

M_x	Alexander [7]	ones and Xenophonos [8]	Bi-CBSG approach [9]	Finite difference approach [16]	2-dimensions Extended cubic B-spline approach	Analytic [14]
0	0.2982	0.2982	0.29468	0.29410	0.29359	0.29468
2	0.2632	0.2631	0.25890	0.25862	0.25828	0.25890
5	0.1743	0.1742	0.17160	0.17147	0.17148	0.17160
8	0.1201	0.1201	0.11878	0.11865	0.11875	0.11878

In Table 2, the results of the 2-dimensions Extended cubic B-spline method using a mesh of 20×20 were compared with those the numerical [7–9, 16] and also with the analytic solution of Shercliff [14].

In Fig. 3, we show The profile of velocity with Hartmann numbers 0 (top curve), to 8 (bottom curve) at $[-1, 1]$ using a mesh of 20×20 .




Figure 3: The profile of velocity with various values of Hartmann numbers

In Table 3, some other results are presented where the period with from $[-1, 1]$ to $[-0.5, 0.5]$ is changed and we also compare these results with Finite difference method [16] and the analytical solution found in the research [14].

Table 3: U at the centre of the duct. Finite difference and analytic simulations compared

M_x	Finite differenc method using a mesh of 50×50 [16]	2-dimensions Ex- tended cubic B-spline method 50×50	Analytic [14]
0	0.073648	0.0736279	0.073671
2	0.071109	0.0710908	0.071128
5	0.060838	0.0608273	0.060846
8	0.049359	0.0493563	0.049363

In Fig. 4, we show the profile of the velocity with various values of Hartmann numbers at $[-0.5, 0.5]$ using a mesh of 50×50 . For diverse values of the Hartmann number,




Figure 4: The profile of velocity with various values of Hartmann numbers.

the course of action for the speed profile along the x -axis has shown up in figs 4 and 5. As would be expected, growing the appealing field (growing the Hartmann number) has an affect on the speed of the fluid where it is the speed reduces near to the center of the channel, this clear affect of the alluring field concentrated is as of presently known. In this way, we see that the comes about are totally congruous with the physical meaning of the affect of the alluring field.

The third test problem: [18, 33–36]

We take the third test problem in the 2-dimensional in this form:

$$u_{xx}(x, y, z) + u_{yy}(x, y, z) - \sin(\pi x) \sin(\pi y) = 0, \quad x, y \in [a, b] \quad (29)$$

the exact solution to that problem given as follows:

$$u(x, y, z) = -\frac{\sin(\pi x) \sin(\pi y)}{2\pi^2}. \quad (30)$$

We take the boundary conditions to the third problem in this form:

$$u(a, y) = u(x, a) = \alpha, \quad u(b, y) = u(x, b) = \beta. \quad (31)$$

By substitution from (4)-(6) into (29) with (31) we obtain the numerical results as in the next table:

Table 4: The numerical results for third problem at $y = 0.4$, $x, y \in [0, 1]$

x	Numerical results	Exact results	Absolute error
0.2	-0.0282154	-0.0283201	1.04701 E-4
0.4	-0.0456535	-0.0458229	1.69408 E-4
0.6	-0.0456535	-0.0458229	1.69408 E-4
0.8	-0.0282154	-0.0283201	1.04701 E-4

In Table 4, we compared the results of the 2-dimensions Extended cubic B-spline strategy employing at 15×15 and the exact results together. From our results we can say that results are accepted with regard to the exact results. In Figs. 5, 6 we show the numerical results with the exact results at $y = 0.5$.




Figure 5: The numerical results with the exact results at $y = 0.4$.




Figure 6: The numerical results with the exact results at $x = 0.4$.

Let 15×15 grid points, we compar between the results of the proposed method and the results of using different methods that shown in Table 6 [18, 33–36].

Table 5: Maximum absolute error according to the method used for third problem.

The pro- posed method	Quadratic B-spline approach [18]	MCBDQM approach [33]	Spline- based DQM approach [34]	Haar wavelet ap- proach [35]	Cubic B-spline ap- proach [36]
1.69 E-4	3.72 E-5	2.11 E-5	1.62 E-4	3.08 E-4	1.67 E-4

The fourth test problem: [18]

We take the fourth test problem in the 3-dimensional in this form:

$$u_{xx}(x, y, z) + u_{yy}(x, y, z) + u_{zz}(x, y, z) - xyz(e^{x+y+z})(3yxz + yx + zx - 5x + zy - 5y - 5z + 9) = 0, \quad x, y, z \in [a, b] \quad (32)$$

The exact solution to that problem given as follows:

$$u(x, y, z) = (x - x^2)(y - y^2)(z - z^2)e^{x+y+z}. \quad (33)$$

We take the boundary conditions to the fourth problem in this form:

$$u(a, y, z) = u(x, a, z) = u(x, y, a) = \alpha, \quad u(b, y, z) = u(x, b, z) = u(x, y, b) = \beta. \quad (34)$$

By substitution from (12) into (32) with (34) we obtain the numerical results as in the next table:

Table 6: The numerical results for test problem at $z = y = 0.5$, $x, y, z \in [0, 1]$

x	Numerical solution	Exact solution	Absolute error	Quadratic B-spline method [18]
0.1	0.0168635	0.0168984	3.48852 E-5	3.24947 E-5
0.2	0.0331304	0.0332012	7.07378 E-5	6.49943 E-5
0.3	0.0480531	0.0481595	1.06445 E-4	9.65554 E-5
0.4	0.0606859	0.0608280	1.42149 E-4	1.27075 E-4
0.5	0.0698464	0.0700264	1.79967 E-4	1.57835 E-4
0.6	0.0740704	0.0742955	2.25088 E-4	1.92337 E-4
0.7	0.0715583	0.0718456	2.87275 E-4	2.37433 E-4
0.8	0.0601139	0.0604965	3.82576 E-4	3.04639 E-4
0.9	0.0370736	0.0376082	5.34586 E-4	4.11161 E-4

In Table 6, we compared the results of the 3-dimensions Extended cubic B-spline strategy employing at 20×20 and the exact results together. From our results we can say that results are accepted with regard to the exact results. In Fig. 7, we show the numerical results with the exact results at $y = z = 0.5$.




Figure 7: The numerical results with the exact results at $y = z = 0.5$.

5 Conclusion

Perhaps by the end of this work, we will have made a clear contribution to solving some of the problems facing most researchers in various fields through how to deal with mathematical models of different dimensions. The topic studied is very important and we believe that most researchers are waiting for its results. Thinking about this work came after we followed what was presented by some researchers in solutions of partial differential equations in one, two and three dimensions, and we noticed how difficult it is for them to deal with these models as the dimension increases. So we thought to develop the Extended cubic B-spline method that was used previously in solving one-dimensional mathematical problems and we were able to present a shape for this method in two and three dimensions. We tested the accuracy and effectiveness of the derived shapes by providing some numerical examples with different dimensions. The numerical results were compared with the real solution, and the inferred formulas were found effective and accurate. From this perspective, we can say that a clear contribution has been made to overcome the problems of partial differential equations of different dimensions. Amid long-term work, we are going moreover generalize a few other B-Splines shapes to serve as a solution to differential equations in n-dimensions.

References

- [1] R. I. Nuruddeen, K. suliman and K. K. Ali, Analytical Investigation of Soliton Solutions to Three Quantum Zakharov-Kuznetsov Equations, *Commun. Theor. Phy.*, 70(4), 405-412 (2018).
- [2] K. K. Ali, R. I. Nuruddeen and K. R. Raslan, New structures for the space-time fractional simplified MCH and SRLW equations, *Chaos, Solitons & Fractals* 106, 304-309 (2018).
- [3] K. R. Raslan, K. K. Ali and M. A. Shallal, The modified extended tanh method with the Riccati equation for solving the space-time fractional EW and MEW equations, *Chaos, Solitons & Fractals*, 103, 404-409 (2017).
- [4] K. R. Raslan, T. S. El-Danaf, K. K. Ali, Exact solution of the space-time fractional coupled EW and coupled MEW equations, *The European Physical Journal Plus*, 132,1-11 (2017).
- [5] K. K. Ali, R. I. Nuruddeen, K. R. Raslan, New hyperbolic structures for the conformable time-fractional variant bussinesq equations , *Optical and Quantum Electronics*, 50, 61 (2018).
- [6] G.W. Sutton and A. Sherman, *Engineering Magnetohydrodynamics* (McGraw-Hill, New York, 1965).
- [7] H. Alexander, An application of the extended Kantorovich method to MHD duct flow, *Acta Mech.* 11, 171(1971).
- [8] R. Jones and J. Xenophontos, An iterative finite element method using the Kantorovich technique, *Acta Mech.* 26, 91-101 (1977).
- [9] L.R.T. Gardner, G.A. Gardner, A two dimensions cubic B-spline finite element: used in a study of MHD-duct flow, *Comput. Methods Appl. Mech. Engrg.* 124, 365-375 (1995).
- [10] Xiujie Zhang, Zengyu Xu and Chuanjie Pan, Numerical analysis of MHD duct flow with a flow channel insert, *Fusion Engineering and Design*, 85, (2010), 2090-2094.
- [11] C. N. Kim, Numerical analysis of a magnetohydrodynamic duct flow with flow channel insert under a non-uniform magnetic field, *Journal of Hydrodynamics*, 30(6), , 1134-1142 (2018).

- [12] C. R. Brewitt-Taylor and J. T. Weave, On the finite difference solution of two-dimensional induction problems, *Geophys. J. R. astr. SOC.*, 47, 375-396 (1976).
- [13] Chia-Ming Fana, Po-Wei Lia, Generalized finite difference method for solving two-dimensional Burgers' equations, *Procedia Engineering*, 79, 55-60 (2014).
- [14] J. A. Shercliff, Steady motion of conducting fluids in pipes under transverse magnetic fields, *Proc. Camb. Philos. Soc.* 49, 139 (1953).
- [15] P.C. Lu, A study of Kantorovich's variational method in MHD duct flow, *AIAA J.*, 5, 1519 (1967).
- [16] K. R. Raslan and Khalid K. Ali, Numerical study of MHD-duct flow using the two-dimensional finite difference method, *Appl. Math. Inf. Sci.* 14, No. 4, 1-5 (2020).
- [17] R. Arora, Swarn Singh and Suruchi Singh, Numerical solution of second-order two-dimensional hyperbolic equation by bi-cubic B-spline collocation method, *Mathematical Sciences*, <https://doi.org/10.1007/s40096-020-00331-y> (2020).
- [18] K. R. Raslan and Khalid K. Ali, On n-dimensional quadratic B-splines, *Numer. Methods Partial Differential Eq.*, 1-15, (2020). <https://doi.org/10.1002/num.22566>
- [19] Frazer, R.A., Jones, W.P., Skan, S.W.: Springer, New York (1937, ARC R and M 1799).
- [20] Bickley, W.G.: Experiments in approximating to solutions of a partial differential equation. *Philos. Mag.* 32(7), 50-66 (1941).
- [21] K. R. Raslan, Talaat S. EL-Danaf, Khalid K. Ali, Application of linear combination between cubic B-spline collocation methods with different basis for solving the KdV equation, *Computational methods for differential equations*, 4 (3), 191-204, (2016).
- [22] K. R. Raslan, Talaat S. EL-Danaf and Khalid K. Ali, Collocation Method with Quintic B-Spline Method for Solving the Hirota equation, *Journal of Abstract and Computational Mathematics*, 1, 1-12, (2016).
- [23] T. S. EL-Danaf, K. R. Raslan and Khalid K. Ali, " collocation method with cubic B- Splines for solving the GRLW equation", *Int. J. of Num. Meth. and Appl.*, 15(1), 39-59, (2016).
- [24] K. R. Raslan, Talaat S. EL-Danaf and Khalid K. Ali, Collocation method with Quintic b-spline method for solving hirota-satsuma coupled KDV equation, *International Journal of Applied Mathematical Research*, 5 (2), 123-131, (2016).
- [25] K. R. Raslan, Talaat S. EL-Danaf and Khalid K. Ali, Application of Septic B-Spline Collocation Method for Solving the Coupled-BBM System, *Appl. & Comput. Math.*, 5 (5), 2-7, (2016).
- [26] K. R. Raslan, Talaat S. EL-Danaf and Khalid K. Ali, collocation method with cubic trigonometric B- Splines algorithm for solving Coupled Burgers' Equations, *Far East Journal of Applied Mathematics*, 95 (2), 109-123, (2016).
- [27] Muhannad Shallal, Khalid K. Ali, K. R. Raslan, Abbas Taqi, Septic B-spline collocation method for numerical solution of the coupled Burgers' equations, *Arab Journal of Basic and Applied Sciences University*, 26(1), 331-341, (2019).
- [28] R.C. Mittal and Amit Tripathi, Numerical solutions of two-dimensional unsteady convection-diffusion problems using modified bicubic B-spline finite elements, *International Journal of Computer Mathematics*, 94(1), 1-21, (2017).
- [29] R.C. Mittal and Amit Tripathi, Numerical solutions of generalized Burgers-Fisher and generalized Burgers-Huxley equations using collocation of cubic B-splines, *International Journal of Computer Mathematics*, 92(5), 1053-1077, (2015).

- [30] Mohammadi, A., Aghazadeh, N., Rezapour, S.: Haar wavelet collocation method for solving singular and nonlinear fractional time-dependent Emden-Fowler equations with initial and boundary conditions. *Math. Sci.* 255-265 (2019).
- [31] AzarNavid, B., Emamjome, M., Nabati, M., Abbasbandy, S.: A reproducing kernel Hilbert space approach in meshless collocation method. *Comput. Appl. Math.* 38-72 (2019).
- [32] Nojavana H., Abbasbandya S., Mohammadi M., Local variably scaled Newton basis functions collocation method for solving Burgers' equation. *Appl. Math. Comput.* 330, 23-41 (2018).
- [33] A. M. Elsherbeny, R. M.I. El-hassani, H. El-badry and M. I. Abdallah, *Solving 2D-Poisson equation using modified cubic B-spline differentialquadrature method*, *Ain Shams Engineering Journal*, 9(4), 2879-2885, (2018).
- [34] G. Mohammad, *Spline-based DQM for multi-dimensions PDEs:Application to bi-harmonic and Poisson Equations in 2D and 3D*, *ComputMath Appl*, 73(7), 1576-1592, (2017).
- [35] S. Zhi, C. Yong-yan and j. Qing, *Solving 2D and 3D Poisson equations andbi-harmonic equations by the Haar wavelet method*, *Appl. Math. Model.*, 36(11), 5134-5161, (2012).
- [36] K. R. Raslan and Khalid K. Ali, A new structure formulations for cubic B-spline collocation method in three and four-dimensions, *Nonlinear Engineering*, 9, 432?448, (2020).