

Canada Under Fire – Drivers and Impacts of the Record-Breaking 2023 Wildfire Season

Piyush Jain¹
Quinn E. Barber¹
Steve Taylor²
Ellen Whitman¹
Dante Castellanos Acuna³
Yan Boulanger⁴
Raphaël D. Chavardès^{4,5,6}
Jack Chen⁷
Peter Englefield¹
Mike Flannigan⁸
Martin P. Girardin⁴
Chelene C. Hanes⁹
John Little¹
Kimberly Morrison²
Rob S. Skakun¹
Dan K. Thompson⁹
Xianli Wang¹
Marc-André Parisien¹

¹ *Northern Forestry Centre, Canadian Forest Service, Natural Resources Canada, 5320-122 St. NW, Edmonton, AB, T6H 3S5, Canada.*

² *Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, 506 West Burnside Rd., Victoria, BC, V8Z 1M5, Canada.*

³ *Department of Renewable Resources, University of Alberta, General Services Building, 9007-116 St. NW, Edmonton, AB, T6G 2H1, Canada.*

⁴ *Laurentian Forestry Centre, Canadian Forest Service, Natural Resources Canada, 1055 du P.E.P.S., P.O. Box 10380, Stn. Sainte-Foy, QC, G1V 4C7, Canada.*

⁵ *Atlantic Forestry Centre, Canadian Forest Service, Natural Resources Canada, 1350 Regent Street, P.O. Box 4000, Fredericton, NB, E3B 5P7, Canada.*

⁶ *Institut de recherche sur les forêts, Université du Québec en Abitibi-Témiscamingue, 445 boulevard de l'Université, Rouyn-Noranda, QC, J9X 5E4, Canada.*

⁷ *Air Quality Research Division, Atmospheric Science and Technology Directorate, Environment and Climate Change Canada, Ottawa, ON, K1V 1C7, Canada.*

⁸ *Natural Resource Science, Thompson Rivers University, 805 TRU Way, Kamloops, BC, V2C 0C8, Canada.*

⁹ *Great Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, 1219 Queen Street E, Sault Ste. Marie, ON, P6A 2E5, Canada.*

Abstract

The 2023 wildfire season in Canada was unprecedented in its scale and intensity. Spanning from late April to early November and extending across much of the forested regions of Canada, the season resulted in a record-breaking total area burned of approximately 15 million hectares, over seven times the historic national annual average. The impacts were profound with more than 200 communities evacuated (approximately 232,000 people), periods of dense smoke that caused significant public health concerns, and unprecedented demands on fire-fighting resources. The exceptional area burned can be attributed to several environmental factors that converged early in the season to enable extreme fire danger over much of the country. These factors included early snowmelt, interannual drought conditions in western Canada, and the rapid transition to drought in eastern Canada. Furthermore, the mean May–October temperature over Canada in 2023 was a staggering 2.2°C warmer than normal (1991–2020), enabling sustained extreme fire weather conditions throughout the fire season. These conditions led to a larger than normal proportion of very large fires (> 50,000 hectares), many having burned for months from the spring into the fall. Fires that started in May or June accounted for over two-thirds of the total area burned. Overall, the 2023 wildfire season in Canada was characterized by its exceptional scale and major societal impacts, setting new records and highlighting the increasing challenges posed by wildfires in the country.

Significance Statement:

The 2023 wildfire season in Canada was unprecedented, extending from coast to coast, and burning approximately 15 million hectares between late April and the end of October – over seven times the historic national average – challenging firefighting efforts. More than 200 communities were evacuated, affecting hundreds of thousands of people, while dense smoke posed significant public health risks. Environmental factors, including early snowmelt, prolonged drought in the west and rapid transition to drought in the east, with unusually high temperatures, created extreme fire conditions. The season saw a larger than average proportion of very large fires, many burning for months, including one fire complex growing to over one million hectares. This record-setting season underscored the escalating challenges of wildfires in Canada.

Introduction

The 2023 wildfire season in Canada was one of superlatives and broken records. At approximately 15 Mha, the area burned was by far the highest since the start of comprehensive national reporting (c. 1975), shattering the previous record of 6.7 Mha in 1989 (Skakun et al. 2022). Four provinces and territories registered their highest recorded annual area burned, with a massive fire complex reaching over 1 Mha, larger than any fire since 1950. Over 200 communities had to be evacuated—some twice—under stressful

or logistically complex circumstances. There were several fatalities, and hundreds of homes and important infrastructure were destroyed. The smoke produced by Canadian wildfires had widespread consequences, affecting not only nearby communities, but also major population centres >1,000 km from the wildfires, notably in southern Canada and on the east coast of the USA (Wang et al. 2023). An unprecedented contingent of wildland firefighters was deployed to respond to the wildfires and international help was obtained from 12 countries and the European Union. The aftermath of the 2023 fire season will likely affect communities and ecosystems for years, if not decades to come.

Canada's climate has been warming at twice the global rate: the mean annual temperature has increased 1.7°C since 1948 nationally, with larger increases at high latitudes and during winter and spring (Bush et al. 2019). Nationally, the annual area burned in the country has been increasing since the 1950s (Hanes et al. 2019). While 2023 was unique in both magnitude and character, scientists and managers have long anticipated the growing potential for increased fire activity (Flannigan et al. 2005; Flannigan and Van Wagner, 1991). The length of the wildfire season—the window of opportunity for wildfire to ignite and grow—has been steadily increasing due to warming over the last 50 years (Wotton and Flannigan 1993; Jain et al. 2017; Hanes et al. 2019). Projections indicate more frequent and intense fire-conducive weather, especially in the central and western part of the country (Wotton et al. 2017; Wang et al. 2020). Moreover, recent wildfire seasons have been associated with extreme weather phenomena affecting an uncommonly large area. For example, the 2021 “heat dome” that covered much of northwestern North America brought record temperatures and entrained one of the worst wildfire seasons of the last century in the province of British Columbia, contributing to the abrupt increase in fire activity in the province since ~2000 (Parisien et al. 2023). Undoubtedly, the increasing potential for wildfires in Canada is symptomatic of the changing climates; in fact, it was estimated that the heat dome would have been 150 times less likely to occur without the ongoing anthropogenic climate disruption (Philip et al. 2021).

Despite this growing body of knowledge about changing fire patterns, the 2023 wildfire season has challenged our understanding of wildland fire in Canada. Canada is about the width of a Rossby wave, the planetary wave pattern (Madden 1979) associated with meridional flow of the Jet Stream (Hoskins and Ambrizzi 1993). Severe fire weather is more likely to develop on one side of a wave, leading typical active fire years to occur regionally, rather than throughout the country (Macias Fauria et al. 2011). Nationwide fire activity, affecting nearly all forested areas in a single season as experienced in 2023, was unprecedented and was not anticipated until later in the century based on climate change projections (Flannigan et al. 2005). Equally surprising, the fire activity in 2023 was relentless: except for a handful of days, many large wildfires burned without respite from late April to early October, and several regions underwent two or more significant surges in fire activity. Uncommon fire behavior, including a pyro-tornado in British Columbia (<https://www.uwo.ca/ntp/>), and a record number of pyrocumulonimbus events were observed due to the frequency and intensity of extreme weather and fire intensity. Consequently, large areas of less-flammable vegetation, such as broadleaf-dominated forests and recently burned stands, were affected by wildfires.

The goal of this article is to provide an overview of this exceptional year, and to examine the drivers and impacts of the record-breaking 2023 wildfire season in Canada. Specific objectives consist of: (i) describing the spatiotemporal patterns of fire activity, (ii) investigating the drivers of fire activity (i.e., weather conditions, ignition source, and fire duration), and (iii) examining the wildfires' observed and

potential impacts on people and ecosystems. To achieve this, we use several datasets quantifying wildfire activity, weather and climate, ignitions, fire management response, and societal impacts. We interpret and discuss our observations relative to past wildfire seasons, but also through the lens of a rapidly changing physical and human environment.

Results and Discussion

Overview of the wildfire season

The 2023 wildfire season extended across seven months (Fig. 1), following early snowmelt that coincided with record-breaking heat. Many of these wildfires caused evacuations, loss of homes, and infrastructure; several continued to burn for the rest of the fire season, reaching very large sizes (e.g., $\geq 10^5$ ha). The fire season abruptly began in mid April with an evacuation in southern British Columbia. Shortly afterward, hundreds of mainly human-caused wildfires began in Alberta in early May. In late May, fast-spreading wildfires on the eastern coast—an area where large wildfires are relatively rare—led to the evacuation of thousands of people from several communities in Nova Scotia and the destruction of hundreds of buildings outside of Halifax and in southern Nova Scotia. In early June, multiple convective cells with associated lightning ignited a string of fires across south-central Quebec, followed by another series of wildfires three weeks later that ignited to the north of the initial ones. The Quebec wildfires burned for weeks and produced a colossal plume of smoke that caused severe smoke pollution in several major cities in Canada and the USA, eventually spreading throughout much of the northern hemisphere (Wang et al. 2023). In midsummer, large wildfires burning in the Northwest Territories caused extensive structural losses and the evacuation of about 70% of the population, including the territorial capital city of Yellowknife. In mid-August, wildfires across British Columbia also burned close to communities, destroying hundreds of structures in south-central British Columbia. Embers transported over 3 km across Okanagan Lake started spot fires that threatened the city of Kelowna. Across Canada, there were many rapidly spreading crown fires over the fire season with extreme fire intensity and towering convection columns. Extreme fire behavior and instability generated 140 pyro-cumulonimbus events in the country, 83% of the total observed globally in 2023 (D. A. Peterson, pers. comm.). As a testament to the widespread and sustained nature of the 2023 wildfire activity, September 22nd—usually a quiet time at the end of the season—was the largest single-day area burned in Canada (~440,000 ha) since satellite records began.

Exceptionally high area burned

The 2023 Canadian fire season was remarkable due to coast-to-coast fire activity lasting from late April to late October, resulting in a record-breaking total burned area of approximately 15 Mha, corresponding to around 4% of Canada's forest area and more than seven times the historic national average. At the level of individual provinces and territories (Table S1), Quebec, the Northwest Territories, British Columbia, and Alberta all registered a record year for total area burned. In comparison with the 1986–2022 period, the area burned in 2023 eclipsed the previous record of 6.7 Mha burned in 1989 (Fig. 2). The widely reported preliminary figure of 18.5 Mha burned is based on active fire data from the fire management agencies for the National Fire Situation Report (CIFFC, <https://ciffc.net/>). The agencies provide invaluable real-time information during the fire season, but the data is considered to be preliminary estimates for daily

operations. Final quality control to account for water bodies and unburned vegetation islands is performed postseason. Our revised estimate is based on a hybrid NBAC-M3 dataset, combining ~12 Mha mapped at high resolution with Landsat and Sentinel-2 satellite imagery and ~3 Mha of perimeters estimated from thermal anomalies.

There were about ~6,700 reported wildfire ignitions in 2023, which represents—surprisingly— a lower number than the average of about 8,000 wildfires per year (Hanes et al. 2019). Lightning ignited 59% of the wildfires and lightning-caused wildfires accounted for 93% of the total area burned (long-term average = 91%) (Hanes et al. 2019). Lightning storms that caused multiple simultaneous ignitions that are beyond initial attack capacity led to several large fires, some of which eventually coalesced. Lightning fires that started on four days, May 13th, May 27th, June 1st and July 5th, were responsible for 30% of the total annual area burned. Interestingly, 2023 had the third lowest overall lightning detections since 1998. The lightning-ignition efficiency was high due to extensive areas of dry, receptive ground fuels coinciding with the specific timing and location of lightning events (Rao et al. 2023), a phenomenon that is expected to increase as a function of the ongoing warming and drying (Hessilt et al. 2022). Although they are responsible for a comparatively low proportion of the total area burned (7%), human-caused ignitions were numerous early in the season, before the greening of vegetation and the prevalence of lightning storms (Parisien et al. 2023). Many of these, notably in Alberta, burned for months and caused evacuations of communities. Yet, it is difficult to confidently assign a human cause to some fires due to ongoing investigations and the eventual intermingling of individual fires with multiple ignition sources in large wildfire complexes.

Drivers of fire activity

Early season drought conditions

In 2023, much of Canada faced a significant moisture deficit at the onset of the fire season, primarily attributed to prolonged drought and premature snowmelt. The country was snow free considerably earlier than the 2004–2022 average, particularly in northern and west-central Canada (British Columbia, Alberta, and Saskatchewan), and along the east coast (Fig. 3 a, b). The maximum root zone soil moisture drying rate increased in western Canada in early May, followed by rapid, intense drying in eastern Ontario and southern Quebec in late May and early June (Figs. 3 c, d). Standardized anomalies of the Canadian Fire Weather Index (FWI) System Drought Code (DC), show similar trends, with severe drought conditions in some northern and western areas in May, and new-onset drought conditions in Quebec and Nova Scotia in June (Fig. 3d, e). Early-season drought is a common occurrence in western Canada (Tymstra et al. 2021), due to persistent drought carried over from the previous year and exacerbated by a low winter snowpack (Hanes et al. 2020) (Fig. S1). In contrast, the 2023 fire season started with near-normal levels of soil moisture following snowmelt in the eastern provinces, but above-normal temperatures and rapid drying caused what could be described as a ‘flash drought’, an emerging phenomenon that we are only beginning to understand (Christian et al. 2024).

Canada experienced extreme fire weather conditions in 2023, recognized as the hottest year on record globally (NOAA 2024). Mean fire season daily temperature, precipitation, and vapor pressure deficit (VPD) anomalies confirm that it was significantly hotter and drier in Canada than normal (Fig. 4). The mean May to October fire season temperature was 2.2°C warmer than the baseline period (1991–2020), with higher values in northern Canada and average-to-below average values in southern Ontario, Quebec, and New Brunswick. This anomaly is already comparable to what is expected in summertime temperature increases over Canada for ~2050, based on projections from CMIP6 climate models under the SSP5-8.5 scenario (<https://climate-scenarios.canada.ca/?page=cmip6-scenarios>). VPD is a good predictor of fuel moisture globally (Rodrigues et al. 2024) and area burned regionally (Parks and Abatzoglou, 2020). Many daily mean temperature and VPD extremes were observed across the country in 2023; for example, on July 7 the temperature at Norman Wells, NWT (65° N), just south of the Arctic Circle, reached 38 °C. Precipitation deficits were widespread, particularly in regions of western and northern Canada and western Quebec; all areas that experienced unusually large wildfires during 2023.

Fire growth is largely determined by the occurrence of extreme weather conditions (Wang et al. 2014). Here, we define extreme fire weather when the Fire Weather Index (FWI) exceeds the climatological 95th percentile of FWI calculated at each location (FWI₉₅). The FWI is a meteorological-based measure of overall fire danger used extensively in fire management (Van Wagner 1987). In 2023, there was a strong correlation between the forested area where FWI exceeded FWI₉₅ and the daily burned area in each province and territory (Fig. 5). Because conditions were extreme for many regions of the country for an extended period of time (e.g., 2-5 months), this enabled the synchronous burning observed in both western and eastern Canada. Synchronous extreme fire weather conditions across large areas has previously been identified as a proxy for constraints on fire suppression resources (Magnussen and Taylor 2012, Abatzoglou et al. 2021). The highest number of extreme fire weather days occurred in northeastern British Columbia, northern Alberta, and southern Northwest Territories (Fig. 6a). Nationwide, fire weather was most extreme in May and June (Fig. 6b), as a result of significant spring drought conditions over much of the country (Figs. 3 & 4). When examining the mean proportion of daily forested area exceeding FWI₉₅, 2023 was also the most extreme fire weather year since at least 1940 (Fig. 6d). Similar relationships were also found for two other FWI System components (Van Wagner 1987), the Initial Spread Index (ISI) and the Buildup Index (BUI) (Fig. S2). Although drought conditions dominated much of the 2023 fire season, days of extreme fire growth were also related to the presence of high near-surface winds, as evidenced by the largest growth days typically coinciding with short-lived (1–3 days) surges in the FWI (Fig. S6).

Large-scale atmospheric circulation patterns have a strong influence on weather extremes. Atmospheric ridges and the associated positive 500-hPa geopotential height anomalies are associated with fire conducive weather, such as ‘blocking’ highs (Skinner et al. 2002, Jain and Flannigan 2021). Blocking can be characterized by persistent positive anomalies (PPAs) in 500-hPa geopotential heights (Tibaldi and Molteni, 1990). Sharma et al. (2022) found PPAs in North America were associated with positive fire-weather anomalies and a seven-fold increase in the likelihood of wildfire ignitions. In 2023, the persistent and widespread extreme fire weather conditions were likely related to the presence of frequent blocking events. Whereas for the baseline period (1991–2020) the mean number of annual

(April–October) blocking days in Canada is ~15, Figure 6c shows the number of blocking days during 2023 far exceeded this value in most parts of the country, both in western and eastern Canada.

Fire size and duration

Most wildfires in Canada’s managed forest are contained or self-extinguished before they reach 200 ha in size; however, fires that survive and exceed 200 ha (“large fires”) account for approximately 97% of the area burned (Stocks et al. 2002; Hanes et al. 2019). In 2023, there were many more large fires in Canada (834) than in previous years (average of 320 for 1986–2022, Fig. S4), as well as a substantially higher frequency of very large fires > 50,000 ha (Fig. 7a). The contribution of very large fires in 2023 is noteworthy: 60 of them were responsible for 73% of the total area burned, compared to an average of 7 very large fires with 41% of the total area burned in the historical period (Fig. 7a). An uncommon event during most years, wildfire complexes where two or more wildfires eventually burn together resulted in 6 of the 10 largest, long-burning wildfires on record (Figs. S5, S6).

How did so many fires become so large in 2023? The likelihood of fire growth in any time period depends on the probability of extreme weather favoring growth, and the absence of fire-ending events such as significant rain or the onset of winter, assuming no fuel limitations (Wiitala and Carlton 1994). Because the number of possible spread days for a given fire is limited by the fire duration, fire size also covaries with the time from ignition to control or extinguishment (Xi et al. 2020). This is illustrated by the growth of the 10 largest wildfires that burned in Canada in 2023 (Figs S5, S6). Of the 43 very large fires (> 50,000 ha) that occurred in western Canada (west of 85°W), the median duration was 82 days (90% CI 32–156) in contrast to only 40 days (22–89) for the 17 very large fires that occurred in eastern Canada. The relationship between the number of potential spread days (defined here as days exceeding the 95th percentile of fire season FWI during each fire and at its location) and fire size is shown in Fig. 7b. In fact, a robust power-law relationship exists between extreme fire weather days (i.e., potential spread days) and fire size throughout Canada (Wang et al. 2020). Interestingly, large fires that burned in western Canada burned over a longer period than those in eastern Canada, but under similar numbers of extreme fire weather days, relative to the local FWI climatology. Although conditions were extreme in each regional context, typical fire weather conditions in western Canada are more severe overall due to climatologically higher aridity. The fires in eastern Canada that grew very large over a relatively short duration, compared to those in western Canada, may also reflect bottom-up factors such as fuel continuity and topography although further work is required to understand these regional differences (Wang et al. 2014).

The seasonal timing of ignitions also influences fire size, and this was particularly evident in 2023, as environmental conditions favoring fire growth over extinguishment persisted for extended periods in many regions of Canada (Figs. S5 & S6), particularly in western Canada where some fires burned for five months. Wildfires that began in the spring and were not extinguished quickly grew larger (Fig. 7c) and accounted for an outsized proportion of the total burn area. The proportions of fires > 1,000 ha ignited in May, June, July, and August corresponded to 0.12, 0.24, 0.37, 0.08, respectively, but were responsible for substantially different proportions of the total area burned (0.35, 0.34, 0.24, 0.05, respectively; see Fig. 7d). A smaller number of large fires that began in April and September had a negligible contribution to the area burned nationally.

Fire management considerations

In 2023, approximately 79% of wildfires detected in Canada received full suppression response, with the remaining 21% receiving modified or monitored (i.e., no direct intervention) response. However, modified and monitored fires contributed 61% percent of the total burned area. The number of active wildfires placed an overwhelming demand on resources throughout most of the 2023 fire season (Fig S3). A national preparedness level of 5, signifying an extreme fire load, insufficient national resources, and limited exchange capabilities, was declared on May 11 and persisted for a record 120 consecutive days. Unprecedented personnel imports and exchanges occurred within Canada (over 1,700 individuals), and over 5,500 individuals from 12 countries and the European Union provided assistance. Additionally, the collective fire-fighting effort included contract firefighting crews, structural firefighters from municipal and local governments, individuals who volunteered, and the Canadian Armed Forces that were deployed to aid in firefighting efforts in multiple provinces. A significant number of firebreaks were created to halt fire spread and protect communities and critical infrastructure. As a result of this record fire-suppression effort, approximately 85% of full-response fires were contained to under 200 hectares, in contrast to 62% of modified- and monitored-response fires that did not exceed this size.

Societal and ecosystem impacts

Aside from fire management, the human dimensions of wildfire also include the vulnerability of communities or individuals to wildfire smoke, evacuations, and the potential loss of structures or even lives. Wildfires in Canada in 2023 resulted in profound societal impacts. Hundreds of thousands of people in communities were affected by evacuations, structure losses, power outages, and business interruptions, and millions of people in North America were exposed to wildfire smoke. Tragically, eight people working on wildfires in Canada were killed during the 2023 fire season. The approximately 15 Mha burned represents a vast area of fire-affected forest, with important implications for post-fire ecological recovery, habitat conservation, human use and enjoyment of forest resources, and the Canadian forestry industry.

Air quality

Fine particulate matter with an aerodynamic diameter less than 2.5 μm ($\text{PM}_{2.5}$) is a known air pollutant; exposure to $\text{PM}_{2.5}$ from wildfires has been shown to significantly impact human health (Matz et al. 2020). For communities near sources of wildfires, smoke causes direct visibility degradation and hazardous exposures to air pollutants. Smoke can be transported hundreds to thousands of kilometers downwind, thus causing population-level exposure to elevated levels of $\text{PM}_{2.5}$ across large regions. Figure 8c shows the air quality warnings (bulletins) issued by forecasters at Environment and Climate Change Canada (ECCC) for possible onset of poor air quality conditions (Air Quality Health Index (AQHI), Stieb et al. 2008) for years between 2017 and 2023. Within the current decade, most AQHI alerts are the result of wildfire events across the country, with peaks during June to September. Compared to the annual average of about ~1,300 alerts from 2017–2022, 2023 stands out with ~5,000 alerts issued (Jan–Nov), due to the extreme poor air quality conditions from summer wildfires across Canada. Fires in the southern Northwest Territories, and northwestern Quebec were particularly influential sources of smoke in 2023 (Fig. 8a). In some regions, almost half of the summer (> 60 days) was spent in unsafe air quality

conditions, sometimes in air containing more than 18 times the level of PM_{2.5} required to trigger an air quality warning (Fig. 8b). Smoke transport from fires as point sources (Fig. 8b) also affected many large communities along several eastern states of the USA. Furthermore, satellite imagery from early June 2023 also showed smoke from Quebec fires transported across the Atlantic impacting countries in western Europe (Wang et al. 2023). In 2023, Canadians experienced approximately eight days of poor air quality on average during the fire season; however, severely affected regions with lower populations, such as the Northwest Territories, endured up to 44 poor air quality days (Tables S4 & S5).

Evacuations

Most of Canada's population of ~41 million is concentrated in the southern part of the country, where fire activity is less prevalent. However approximately 12.3% of Canadians live within or adjacent to forested areas, in the wildland urban interface (WUI; Johnston and Flannigan 2017), and these communities are more likely to experience an evacuation when threatened by fire. Indigenous communities, of which 32.1% of the on-reserve population is located within the WUI, are especially vulnerable to evacuation (Erni et al. 2021; Tepley et al. 2022). Evacuations are extremely disruptive for community cohesion and economics, and have negative physical and mental health impacts (Asfaw et al. 2019; Cherry and Haynes, 2017).

In 2023, wildfires triggered evacuations of communities in 12 of the country's 13 provinces and territories (Natural Resources Canada, 2023a). Preliminary estimates indicate that approximately 232,000 people were evacuated in 282 events; the most evacuees of any fire season since records began in 1980 (Beverly and Bothwell, 2011), and nearly three times as many people as were evacuated due to the Horse River Wildfire that burned through Fort McMurray in 2016. Five of the 10 largest evacuations ever recorded in Canada occurred this year (Table S3), with some communities such as Edson in Alberta and Lebel-sur-Quévillon in Quebec, experiencing repeated evacuations. The 2023 fire season will undoubtedly have lasting social and health impacts, throughout the country.

Ecological impacts

The 2023 wildfire season will significantly alter forest landscapes due to extensive stand-replacing wildfires. While most Canadian forests are adapted to large, intense fires and the recent burn rates have remained within the historical range of variability (Chavardès et al. 2022), the scale of this wildfire season (~4% of the total forest area burned) in Canada is well outside what was observed in the recent decades. The widespread burning of young forests (i.e., <30 years since fire or harvest) in 2023 has the potential to cause extensive post-fire tree regeneration failures, because immature trees cannot provide enough seeds following a fire (Reid et al. 2023; Splawinski et al. 2019). These failures, compounded by logging legacies, drought, and insect outbreaks, could reduce forest productivity (Cyr et al. 2022) and carbon stocks (Walker et al. 2019), and accelerate the transition from boreal forests to open taiga, prairies or parklands (Stralberg et al. 2018; Whitman et al. 2019). Forest landscape changes from the 2023 wildfire season will have profound effects on forest ecosystem processes and biodiversity, with species adapted to early-stage or open-canopy forests benefitting, whereas those reliant on mature or old-growth forests, such as woodland caribou (*Rangifer tarandus*), being most negatively affected in the near term (Rudolph et al. 2017; Stewart et al. 2020). The cumulative impacts of the area burned in 2023 coupled with the

extensive anthropogenic disturbance legacies on the landscape will challenge the resilience of forest ecosystems, especially if fire activity continues to increase, as projected (Gauthier et al. 2015).

Conclusions

The 2023 Canadian fire season shattered national and multiple regional records, causing profound societal and health impacts. The combination of extreme weather and an extended fire season led to longer-lasting, hence larger, wildfires, resulting in a burned area approximately seven times the national average. Unprecedented synchronous fire activity strained fire management and response capacities across the country. Climate change has increased wildfire activity globally (Sullivan et al. 2022), including in Canada, where recent extreme fire weather in British Columbia and Alberta has been linked to climate change (Kirchmeier-Young et al. 2019; Whitman et al. 2022). Although research is needed to quantify the importance of climate change to the 2023 fire season at a national scale, anthropogenic climate change strongly contributed to Quebec's worst-recorded fire season (Barnes et al. 2023). Disentangling the relative influence of human landscape legacies and worsening fire weather (Jain et al. 2022; 2017; Jones et al. 2022) on the current Canadian fire environment is ongoing, but it is inescapable that extreme heat and moisture deficits drove the record-breaking 2023 fire season. The disproportionate effect a few days of extreme weather can have on the total area burned is also evident in this fire season (Wang et al. 2023; Coop et al. 2022), leading to worrisome prospects given projected future conditions (Wang et al. 2020). Fire-fuel feedbacks may offset some potential for future fire activity (Boulanger et al. 2017; Gaboriau et al. 2023); however, increasingly severe weather overwhelms this resistance to burning. Although this trajectory cannot easily be changed, we are not without options. Increased focus on innovative and integrated fire management strategies that prioritize prevention and mitigation are gaining momentum. This will require all stakeholders to engage in fostering more fire resilient environments, as the 2023 fire season may foreshadow the emergence of severe climate change impacts on fire activity decades earlier than previously anticipated.

Materials and Methods

Wildfires in Canada

Fire-prone regions of Canada have predominantly boreal and temperate continental climates, with the majority of wildfires occurring between April 1 and September 30. Even though about half are ignited by humans, lightning-caused fires are responsible for a greater proportion (~90%) of the area burned (Hanes et al. 2019), occurring in remote areas (Erni et al. 2020). In some parts of the country, Indigenous cultural burning during spring or fall also drove local fire regimes, though these practices were suppressed beginning in the early 20th century (Lewis et al. 1988; Hoffman et al. 2022). From a meteorological standpoint, the Canadian wildfire season is typically characterized by a series of high-pressure atmospheric systems resulting in drying periods of several days to more than one week, interspersed with rain events (Taylor, 2020). Lightning storms occurring during or after these dry spells can cause several hundred fire starts within a few days in a given region. When followed by persistent warm and dry weather and strong winds, extensive forested areas can burn as crown fires (Frelich et al. 2021).

Fire data

In Canada, high-quality spatial wildfire perimeters are available from the National Burned Area Composite (NBAC; Skakun et al. 2022). At the time of writing, the NBAC product had not yet been completed and consolidated with agency data, so we considered a variety of provisional fire data sources to form a complete picture of fire activity for 2023.

Agency fire data

During the fire season in Canada, fire information reported by fire management agencies are collated by the Canadian Interagency Forest Fire Centre (CIFFC). This data includes reported date, fire ignition cause, and fire management response type. We downloaded reported data for 2023 (<https://ciffc.net/>, accessed December 29, 2023) to determine the number of incidents, management responses, and fire cause.

Subdaily fire detections

Near-real-time fire perimeter polygons were obtained from the M3 fire perimeters (Natural Resources Canada. 2023b.) derived from thermal anomaly (hotspot) point detections using a two-step buffering process as follows. First, VIIRS (375m) hotspots (VIIRS and MODIS (1km) in Quebec) are buffered out r_{out} meters to form circular polygons. Boundaries between intersecting circular polygons are dissolved to form larger polygons. Second, perimeters of the resulting polygons are buffered in (contracted) r_{in} meters. The choice of r_{out} and r_{in} for each ecozone were determined by comparing buffered polygons from 2012–2021 with higher-resolution fire perimeters from the NBAC. Selected radii maximize the intersecting area, and minimize commission and omission errors.

National Burned Area Composite

The provisional 2023 NBAC fire perimeters were created using cloud and shadow-free satellite imagery from the Google Earth Engine Collection-2 catalogue (Landsat-8 and -9 data combined at 30-metres resolution, and Sentinel-2 data at 20-metres). We produced post-fire mosaics using median compositing of images collected over a 30-day period starting after the last hotspot acquisition date of the corresponding M3 perimeter. A pre-fire mosaic was derived likewise using the dates of the post-fire mosaic one year earlier. Image compositing was performed independently for Landsat and Sentinel, and an analyst selected the best quality pre- and post-fire mosaics for image mapping. In this procedure, an adaptive threshold was applied based on the differenced Normalized Burn Ratio to map pixels corresponding to burned areas (Skakun et al. 2022). The largest fires contributing the greatest area burned were targeted for NBAC mapping.

NBAC-M3 2023 hybrid dataset

We produced a hybrid fire perimeter dataset ("NBAC-M3") using the combination of all preliminary NBAC fire perimeters and, for the 18% of the 2023 area burned which was not yet mapped, M3 perimeters (Natural Resources Canada, 2023b). We removed water bodies using the CanVec hydrographic

features polygons (Natural Resources Canada 2019). It was necessary to correct the apparent burned area from M3 buffered hotspots, since buffered points typically overestimate area burned due to the presence of unburned islands and water bodies that are not accounted for in the buffering process. We did this by dividing the NBAC area burned by the total area burned reported in historical M3 buffered hotspots for 2012–2022, stratified by ecozone and year, thereby producing ecozone-specific calibration factors. We multiplied the calculated area of the M3 polygons for 2023 by the correction factors, within ecozones. Finally, polygons less than 1ha in size were removed from the final dataset to minimize any introduced artifacts from the above process. We then summed the calibrated M3 area burned, with the mapped area burned from NBAC polygons to estimate the total area burned in 2023.

Daily fire growth

To estimate daily fire growth, we interpolated the fire detection date from hotspots for all wildfires in the NBAC-interim dataset with a final area over 500 ha. Detection date (NRT VIIRS 2022) was corrected to local standard time and interpolated using ordinary kriging to a 180-m grid, following Parks (2014). A filter was applied to detections prior to April 1st to remove false detections outside of the main fire season. In total, 641 fires were interpolated, accounting for ~99.5% of the total area burned in 2023.

Snowmelt timing

Daily snow cover data was obtained from the Interactive Multisensor Snow and Ice Mapping System (IMS) produced by the United States National Ice Center (USNIC, 2008) at 4-km resolution. We defined snowmelt timing at each location as the first day of the longest snow free period at that location for any given year. In addition, snow melt timing departures for 2023 were determined by the anomaly (days) of the 2023 snowmelt timing from the mean timing of the period 2004–2022.

Drought data

The root zone soil moisture (RZSM) is an indicator of the moisture content of the top 1 m of soil, and as such, is a useful measure of drought. We obtained the RZSM from the Global Land Data Assimilation System (GLDAS-2.2, Li et al. 2020), a land surface model constrained by Data Assimilation of the Gravity Recovery and Climate Experiment (GRACE-DA). The GLDAS-2.2. data is daily at 0.25 degrees resolution, and was subset to North America. To determine rapid onset of drought, we calculated the maximum amount of drying (i.e., reduction in RZSM values, kg m⁻²) that occurred using a 14-day sliding window from May 1st to June 30th. The timing of rapid onset was defined as the midpoint of the window corresponding to the maximum 14-day drying.

Weather and fire weather

Weather data was obtained from the ERA5 reanalysis via the Copernicus Climate Change Service (Hersbach et al. 2023). ERA5 includes an extensive set of surface and upper air atmospheric variables, available hourly and globally at 0.25° resolution. We downloaded 2 m temperature, 2 m dewpoint temperature, precipitation, and the 10 m U and V components of wind. Relative humidity and Vapor

Pressure Deficit (VPD) were estimated from temperature and dewpoint temperature, using the equation from Alduchov and Eskridge for VPD (1996).

We examined mean daily anomalies of temperature, precipitation and vapor pressure deficit (VPD) averaged between May 1st and October 31st (fire season) of 2023 compared with the baseline period (1991–2020); we also plotted the rank of the fire season mean of each variable for 2023 compared with the period 1940–2023.

The Canadian Fire Weather Index System (CFWIS) — part of the Canadian Forest Fire Danger Rating System (CFFDRS; Van Wagner, 1987) — is an empirical model that combines screen level temperature, relative humidity, 10-m open wind speed, and 24-hour accumulated precipitation collected at solar noon (Van Wagner, 1987). The System outputs represent proxies to fuel moisture in the forest floor, potential fire behavior and an overall fire danger rating (the Fire Weather Index, FWI). We calculated CFWIS outputs from ERA5 weather following a procedure to account for interseasonal drought conditions (McElhinny et al. 2020). Fire season was determined using the startup condition that daily maximum temperature should exceed 12° C for three consecutive days, and annual shutdown conditions when temperatures fall below 5° C for three consecutive days, serving as a proxy for snow free periods (Wotton and Flannigan 1993).

We calculated a climatology of extreme FWI values at each ERA5 grid cell, defined as the 95th percentile of FWI values (FWI_{95}) between 1991–2020 (omitting values outside of the fire season). An extreme fire weather day is then any day for which $FWI > FWI_{95}$. Using percentiles to define extreme fire weather accounts for the fact that the fire-weather relationship (e.g., fire ignition and spread) are heterogeneous across the landscape. The extent of extreme fire weather conditions was further defined as the forested area exceeding the local (i.e., grid-level) 95th percentile of the FWI (FWI_{95} , defined by the fire season period during 1991–2020, Wang et al. 2015). We defined forested area by applying a threshold of > 20% canopy cover to the NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Vegetation Continuous Fields (VCF) Version 1 data product (VCF5KYR, Hansen et al. 2018)

To examine the influence of large-scale atmospheric patterns over Canada in 2023 — and in particular, blocking events — we applied the identification algorithm for persistent positive anomalies (PPAs) in 500-hPa geopotential heights, previously developed by Sharma et al. (2022). A PPA was identified for contiguous regions where the 500-hPa geopotential heights exceeded one standard deviation above the local seasonal climatological mean (for the baseline period 1991–2020) and persisted for at least 5 days, with the additional criterion that each event reached at least 100,000 km² during its evolution. We applied the algorithm using a domain that covers the entirety of the forested area of Canada.

Air quality

We compiled PM_{2.5} concentration maps from the first 24-hr model outputs of the ECCC operational Regional Air Quality Deterministic Prediction System-FireWork (ECCC RAQDPSFW TechNote, 2021). FireWork is a state-of-science numerical air quality forecast system with near-real-time wildfire emissions (Chen et al. 2019; O'Neill et al. 2023). The core of the FireWork system is the GEM-MACH

meteorology-chemical-coupled model that simulates the dynamics of chemical composition, accounting for detailed gas-, aqueous- and particle-phase chemical reactions and atmospheric physics. The model domain covers North America with a 10-km grid resolution. Hourly wildfire emissions were estimated using the Canadian Forest Fire Emissions Prediction System (CFFEPS, Chen et al. 2019) for fire grid locations representing hotspot areas (past-24 hr satellite detection by VIIRS sensors). We estimated emissions with forecast meteorology, fuel type, and burn areas through the NRCan Canadian Wildland Fire Information System (Natural Resources Canada, 2023c). Output of the FireWork system consists of hourly concentrations of gas-phase and aerosol species. The total fine PM components represent simulated PM_{2.5} concentrations. We adopted thresholds for ambient levels of PM_{2.5} as regulated under the Canadian Ambient Air Quality Standards (CAAQS, <https://ccme.ca/en/air-quality-report>), with daily and annual standards determined as a 24-hour average concentration of 27 $\mu\text{g m}^{-3}$ and an annual average of 8.8 $\mu\text{g m}^{-3}$.

Air Quality (AQ) Alert Bulletins issued by MSC-ECCC, 2017-2023. Data representing the number of all AQ Alert bulletins issued throughout the period. Each AQ alert will have a minimum of 2 bulletins issued (issuance and termination), plus a number of continued bulletins based on the persistence of the event. Each AQ Alert's area of coverage varies in size based on the geographical extent of the AQ event.

Evacuations

Evacuation data for Canada for 1980–2023 were obtained by request from the Canadian Wildland Fire Evacuation Database (Natural Resources Canada, 2023a). This data is compiled using an exhaustive search of media reports and quality control measures as described by Tepley et al. (2022). Evacuation data from 2023 are provisional and are subject to updates as evacuation numbers are confirmed.

Acknowledgments

The authors would like to thank Dawn McVittie (CFS) for compiling the 2023 evacuation data, Melissa MacDonald at Environment and Climate Change Canada for providing the air quality alert data, and Claudia Castillo for graphic design. We would also like to thank David Peterson, Michael Fromm and Rene Servranckx for sharing pyro-cumulonimbus event statistics for 2023.

Data Availability

1. Snow cover data was obtained from the United States National Ice Center, <https://nsidc.org/data/g02156/versions/1>
2. Data from the North American Drought Monitor was downloaded from: <https://droughtmonitor.unl.edu/NADM/Statistics.aspx>
3. GLDAS-2.2 data was obtained from: https://disc.gsfc.nasa.gov/datasets/GLDAS_CLSM025_DA1_D_EP_2.2/summary
4. ERA5 reanalysis data was obtained from the Copernicus Climate Change Service (2023): ERA5 hourly data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.adbb2d47 (Accessed on 07-Jan-2024)

5. We acknowledge the use of data and/or imagery from NASA's Fire Information for Resource Management System (FIRMS) (<https://earthdata.nasa.gov/firms>), part of NASA's Earth Observing System Data and Information System (EOSDIS).

References

1. Abatzoglou, J.T., Juang, C.S., Williams, A.P., Kolden, C.A. and Westerling, A.L., 2021. Increasing synchronous fire danger in forests of the western United States. *Geophysical Research Letters*, 48(2), p.e2020GL091377.
2. Alduchov, O. A. and Eskridge, R. E. Improved Magnus form approximation of saturation vapor pressure. *J. Appl. Meteorol.* 35, 601–609 (1996).
3. Asfaw, H.W., McGee, T. and Christianson, A.C., 2019. The role of social support and place attachment during hazard evacuation: the case of Sandy Lake First Nation, Canada. *Environmental Hazards*, 18(4), pp.361-381.
4. Barnes C., Boulanger Y., Keeping T., Gachon P., Gillett N., Boucher J., Roberge F., Kew S., Haas O., Heinrich D., Vahlberg M., Singh R., Elbe M., Sivanu S., Arrighi J., Van Aalst M., Otto F., Zacharian M., Krikken F., Wang X., Erni S., Pietropalo E., Avis A., Bisailon A., Kimutai, J. 2023. Climate change more than doubled the likelihood of extreme fire weather conditions in eastern Canada. *Scientific Reports*, <https://doi.org/10.25561/105981>.
5. Beverly, J.L. and Bothwell, P., 2011. Wildfire evacuations in Canada 1980–2007. *Natural Hazards*, 59, pp.571-596.
6. Boulanger, Y., Girardin, M., Bernier, P.Y., Gauthier, S., Beaudoin, A. and Guindon, L., 2017. Changes in mean forest age in Canada's forests could limit future increases in area burned but compromise potential harvestable conifer volumes. *Canadian Journal of Forest Research*, 47(6), pp.755-764.
7. Bush, E. and Lemmen, D.S., editors (2019): Canada's Changing Climate Report; Government of Canada, Ottawa, ON. 444 p.
8. Chavardès RD, Danneyrolles V, Portier J, Girardin MP, Gaboriau DM, Gauthier S, ..., Bergeron Y (2022) Converging and diverging burn rates in North American boreal forests from the Little Ice Age to the present. *International Journal of Wildland Fire*, 31(12), 1184-1193. <https://doi.org/10.1071/WF22090>
9. Chen, J., Anderson, K., Pavlovic, R., Moran, M. D., Englefield, P., Thompson, D. K., Munoz-Alpizar, R., and Landry, H.: The FireWork v2.0 air quality forecast system with biomass burning emissions from the Canadian Forest Fire Emissions Prediction System v2.03, *Geosci. Model Dev.*, <https://doi.org/10.5194/gmd-2019-63>, 2019.
10. Cherry, N. and Haynes, W., 2017. Effects of the Fort McMurray wildfires on the health of evacuated workers: follow-up of 2 cohorts. *Canadian Medical Association Open Access Journal*, 5(3), pp.E638-E645.

11. Christian, J.I., Hobbins, M., Hoell, A., Otkin, J.A., Ford, T.W., Cravens, A.E., Powlen, K.A., Wang, H. and Mishra, V., 2024. Flash drought: A state of the science review. *Wiley Interdisciplinary Reviews: Water*, p.e1714.
12. Coop, J. D., Parks S. A., Stevens-Rumann C. S., Ritter S. M., and Hoffman C. M. (2022). Extreme fire spread events and area burned under recent and future climate in the western USA. *Global Ecology and Biogeography*, 31, 1949–1959. <https://doi.org/10.1111/geb.13496>
13. Cyr, D., Splawinski, T. B., Pascual Puigdevall, J., Valeria, O., Leduc, A., Thiffault, N., ... and Gauthier, S. (2022). Mitigating post-fire regeneration failure in boreal landscapes with reforestation and variable retention harvesting: At what cost?. *Canadian Journal of Forest Research*, 52(4), 568-581.
14. ECCC, 2021. Regional Air Quality Deterministic Prediction System with Near-Real-Time Wildfire Emissions (RAQDPSFW): Upgrade to version 023. Technical note, December 2021, Canadian Centre for Meteorological and Environmental Prediction, Montreal, 31pp. [available https://eccc-msc.github.io/open-data/msc-data/nwp_raqdps-fw/readme_raqdps-fw_en/]
15. Erni S, Wang X, Taylor S, Boulanger Y, Swystun T, Flannigan M, Parisien MA (2020) Developing a two-level fire regime zonation system for Canada. *Canadian Journal of Forest Research*. 50(3): 259-273. <https://doi.org/10.1139/cjfr-2019-0191>
16. Erni, S., Johnston, L., Boulanger, Y., Manka, F., Bernier, P., Eddy, B., Christianson, A., Swystun, T., Gauthier, S. 2021. Exposure of the Canadian wildland–human interface and population to wildland fire, under current and future climate conditions. *Canadian Journal of Forest Research*. 51(9): 1357-1367. <https://doi.org/10.1139/cjfr-2020-0422>
17. Flannigan, M.D. and Van Wagner, C.E., 1991. Climate change and wildfire in Canada. *Canadian Journal of Forest Research*. 21(1): 66-72. <https://doi.org/10.1139/x91-010>
18. Flannigan, M.D., Logan, K.A., Amiro, B.D., Skinner, W.R. and Stocks, B.J., 2005. Future area burned in Canada. *Climatic change*, 72(1-2), pp.1-16.
19. Frelich LE, Lorimer CG, Stambaugh MC (2021) History and future of fire in hardwood and conifer forests of the Great Lakes-Northeastern forest region, USA. *Fire ecology and management: past, present, and future of US forested ecosystems*, 243-285.
20. Gaboriau, D. M., Chaste, É., Girardin, M. P., Asselin, H., Ali, A. A., Bergeron, Y., and Hély, C. (2023). Interactions within the climate-vegetation-fire nexus may transform 21st century boreal forests in northwestern Canada. *Iscience*, 26(6).
21. Gauthier, S., Bernier, P., Kuuluvainen, T., Shvidenko, A.Z. and Schepaschenko, D.G., 2015. Boreal forest health and global change. *Science*, 349(6250), pp.819-822.
22. Hanes CC, Wang XL, Jain P, Parisien MA, Little JM, Flannigan MD (2019) Fire-regime changes in Canada over the last half century. *Canadian Journal of Forest Research*. 49(3): 256-269. <https://doi.org/10.1139/cjfr-2018-0293>

23. Hanes C, Wotton M, Woolford DG, Martell DL, Flannigan M (2020) Preceding Fall Drought Conditions and Overwinter Precipitation Effects on Spring Wildland Fire Activity in Canada. *Fire* 3, 24
24. Hansen, M., X. Song. Vegetation Continuous Fields (VCF) Yearly Global 0.05 Deg. 2018, distributed by NASA EOSDIS Land Processes Distributed Active Archive Center, <https://doi.org/10.5067/MEaSUREs/VCF/VCF5KYR.001>. Accessed 2023-12-12.
25. Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N. 2023: ERA5 hourly data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.adbb2d47 (Accessed on 07-Dec-2023)
26. Hessilt, T.D., Abatzoglou, J.T., Chen, Y., Randerson, J.T., Scholten, R.C., Van Der Werf, G. and Veraverbeke, S., 2022. Future increases in lightning ignition efficiency and wildfire occurrence expected from drier fuels in boreal forest ecosystems of western North America. *Environmental Research Letters*, 17(5), p.054008.
27. Hoffman KM, Cardinal Christianson A, Dickson-Hoyle S, Copes-Gerbitz K, Nikolakis W, Diabo DA, McLeod R, et al. (2022) “The Right to Burn: Barriers and Opportunities for Indigenous-Led Fire Stewardship in Canada.” *FACETS* 7: 464–81.
28. Hoskins BJ, Ambrizzi T (1993) Rossby wave propagation on a realistic longitudinally varying flow. *J Atmos Sci* 50:1661–1671
29. Jain, P. and Flannigan, M., 2021. The relationship between the polar jet stream and extreme wildfire events in North America. *Journal of Climate*, 34(15), pp.6247-6265.
30. Jain, P., Castellanos-Acuna, D., Coogan, S.C., Abatzoglou, J.T. and Flannigan, M.D., 2022. Observed increases in extreme fire weather driven by atmospheric humidity and temperature. *Nature Climate Change*, 12(1), pp.63-70.
31. Jain, P., Wang, X. and Flannigan, M.D., 2017. Trend analysis of fire season length and extreme fire weather in North America between 1979 and 2015. *International journal of wildland fire*, 26(12), pp.1009-1020.
32. Jones, M.W., Abatzoglou, J.T., Veraverbeke, S., Andela, N., Lasslop, G., Forkel, M., Smith, A.J., Burton, C., Betts, R.A., van der Werf, G.R. and Sitch, S., 2022. Global and regional trends and drivers of fire under climate change. *Reviews of Geophysics*, 60(3), p.e2020RG000726.
33. Johnston, L.M. and Flannigan, M.D., 2017. Mapping Canadian wildland fire interface areas. *International journal of wildland fire*, 27(1), pp.1-14.
34. Kirchmeier-Young, M.C., Gillett, N.P., Zwiers, F.W., Cannon, A.J. and Anslow, F.S., 2019. Attribution of the influence of human-induced climate change on an extreme fire season. *Earth's Future*, 7(1), pp.2-10.

35. Lewis, H. T., and Ferguson, T. A. (1988). Yards, corridors, and mosaics: how to burn a boreal forest. *Human ecology*, 16, 57-77. <https://doi.org/10.1007/BF012620>
36. Li B, Beaudoin H, Rodell M, NASA/GSFC/HSL (2020) GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree GRACE-DA1 V2.2, Greenbelt, Maryland, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: [Data Access 16-Oct-2023], 10.5067/TXBMLX370XX8
37. Macias Fauria M, Michaletz ST, Johnson EA (2011) Predicting climate change effects on wildfires requires linking processes across scales. *Wiley Interdisciplinary Reviews: Climate Change*, 2(1), 99-112.
38. Madden, R. A. (1979), Observations of large-scale traveling Rossby waves, *Rev. Geophys.*, 17(8), 1935–1949, doi:10.1029/RG017i008p01935.
39. Magnussen, S. and Taylor, S.W., 2012. Inter-and intra-annual profiles of fire regimes in the managed forests of Canada and implications for resource sharing. *International Journal of Wildland Fire*, 22: 328-341.
40. Matz, C.J., Egyed, M., Xi, G., Racine, J., Pavlovic, R., Rittmaster, R., Henderson, S.B. and Stieb, D.M., 2020. Health impact analysis of PM_{2.5} from wildfire smoke in Canada (2013–2015, 2017–2018). *Science of The Total Environment*, 725, p.138506.
41. McElhinny, M., Beckers, J.F., Hanes, C., Flannigan, M. and Jain, P., 2020. A high-resolution reanalysis of global fire weather from 1979 to 2018—overwintering the Drought Code. *Earth System Science Data*, 12(3), pp.1823-1833.
42. Natural Resources Canada. 2019. Lakes, Rivers and Glaciers in Canada - CanVec Series - Hydrographic Features. Available from open.canada.ca/data/en/dataset/2dac78ba-8543-48a6-8f07-faeef56f9895 [accessed May 20, 2021].
43. Natural Resources Canada. 2023a. Canadian Wildland Fire Evacuation Database. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta. 2023. Unpublished.
44. Natural Resources Canada. 2023b. Fire M3 Season-to-date Buffered Hotspots. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta. Available from: <https://cwfis.cfs.nrcan.gc.ca/> [Accessed 15 November 2023].
45. Natural Resources Canada. 2023c. Canadian Wildland Fire Information System [online]. Available from <http://cwfis.cfs.nrcan.gc.ca/datamart> [Accessed 10 November 2023].
46. NOAA National Centers for Environmental Information, Monthly Global Climate Report for Annual 2023, published online January 2024, retrieved on February 10, 2024 from <https://www.ncei.noaa.gov/access/monitoring/monthly-report/global/202313>.

47. NRT VIIRS 375 m Active Fire product VJ114IMGTDL_NRT distributed from NASA FIRMS. Available on-line <https://earthdata.nasa.gov/firms>. doi:10.5067/FIRMS/VIIRS/VJ114IMGT_NRT.002
48. O'Neill, S.M., Xian, P., Flemming, J., Cope, M., Baklanov, A., Larkin, N.K., Vaughan, J.K., Tong, D., Howard, R., Stull, R. and Davignon, D., 2023. Profiles of Operational and Research Forecasting of Smoke and Air Quality Around the World. Landscape Fire, Smoke, and Health: Linking Biomass Burning Emissions to Human Well-Being, pp.149-191.
49. Parisien, M.A., Barber, Q.E., Bourbonnais, M.L., Daniels, L.D., Flannigan, M.D., Gray, R.W., Hoffman, K.M., Jain, P., Stephens, S.L., Taylor, S.W. and Whitman, E., 2023. Abrupt, climate-induced increase in wildfires in British Columbia since the mid-2000s. Communications Earth & Environment, 4(1), p.309.
50. Parks, S.A. and Abatzoglou, J.T., 2020. Warmer and drier fire seasons contribute to increases in area burned at high severity in western US forests from 1985 to 2017. Geophysical Research Letters, 47(22), p.e2020GL089858.
51. Parks, S.A., 2014. Mapping day-of-burning with coarse-resolution satellite fire-detection data. *International Journal of Wildland Fire*, 23(2), pp.215-223.
52. Philip, S.Y., Kew, S.F., Van Oldenborgh, G.J., Anslow, F.S., Seneviratne, S.I., Vautard, R., Coumou, D., Ebi, K.L., Arrighi, J., Singh, R. and Van Aalst, M., 2021. Rapid attribution analysis of the extraordinary heatwave on the Pacific Coast of the US and Canada June 2021. *Earth System Dynamics Discussions*, 2021, pp.1-34.
53. Rao, K., Williams, A.P., Diffenbaugh, N.S., Yebra, M., Bryant, C. and Konings, A.G., 2023. Dry Live Fuels Increase the Likelihood of Lightning-Caused Fires. *Geophysical Research Letters*, 50(15), p.e2022GL100975.
54. Reid, K. A., Day, N. J., Alfaro-Sánchez, R., Johnstone, J. F., Cumming, S. G., Mack, M. C., ... and Baltzer, J. L. (2023). Black spruce (*Picea mariana*) seed availability and viability in boreal forests after large wildfires. *Annals of Forest Science*, 80(1), 1-16.
55. Rodrigues, M., de Dios, V.R., Sil, Â., Camprubí, À.C. and Fernandes, P.M., 2024. VPD-based models of dead fine fuel moisture provide best estimates in a global dataset. *Agricultural and Forest Meteorology*, 346, p.109868.
56. Rudolph, T. D., Drapeau, P., Imbeau, L., Brodeur, V., Légaré, S., and St-Laurent, M. H. (2017). Demographic responses of boreal caribou to cumulative disturbances highlight elasticity of range-specific tolerance thresholds. *Biodiversity and Conservation*, 26, 1179-1198.
57. Skakun, R.; Castilla, G.; Metsaranta, J.; Whitman, E.; Rodrigue, S.; Little, J.; Groenewegen, K.; Coyle, M. 2022. Extending the National Burned Area Composite Time Series of Wildfires in Canada. *Remote Sensing*, 14, 3050. DOI:[10.3390/rs14133050](https://doi.org/10.3390/rs14133050)

58. Sharma, A.R., Jain, P., Abatzoglou, J.T. and Flannigan, M., 2022. Persistent Positive Anomalies in Geopotential Heights Promote Wildfires in Western North America. *Journal of Climate*, 35(19), pp.6469-6486.
59. Skinner, W., Flannigan, M., Stocks, B. et al. A 500 hPa synoptic wildland fire climatology for large Canadian forest fires, 1959–1996. *Theor Appl Climatol* 71, 157–169 (2002).
<https://doi.org/10.1007/s007040200002>
60. Splawinski, T. B., Greene, D. F., Michaletz, S. T., Gauthier, S., Houle, D., and Bergeron, Y. 2019. Position of cones within cone clusters determines seed survival in black spruce during wildfire. *Canadian Journal of Forest Research*, 49(2), 121-127.
61. Stewart, F. E., Nowak, J. J., Micheletti, T., McIntire, E. J., Schmiegelow, F. K., and Cumming, S. G. 2020. Boreal caribou can coexist with natural but not industrial disturbances. *The Journal of Wildlife Management*, 84(8), 1435-1444.
62. Stieb, D. M., Burnett, R. T., Smith-Doiron, M., Brion, O., Shin, H. H., and Economou, V. 2008. A new multipollutant, no-threshold air quality health index based on short-term associations observed in daily time-series analyses. *Journal of the Air & Waste Management Association*, 58(3), 435-450.
63. Stocks, B. J., J. A. Mason, J. B. Todd, E. M. Bosch, B. M. Wotton, B. D. Amiro, M. D. Flannigan et al. 2002. Large forest fires in Canada, 1959–1997." *Journal of Geophysical Research: Atmospheres* 107, no. D1 : FFR-5.
64. Stralberg, D., Wang, X., Parisien, M. A., Robinne, F. N., Sólymos, P., Mahon, C. L., ... and Bayne, E. M. 2018. Wildfire-mediated vegetation change in boreal forests of Alberta, Canada. *Ecosphere*, 9(3), e02156.
65. Sullivan, A., Baker, E., Kurvits, T., Popescu, A., Paulson, A.K., Cardinal Christianson, A., Tulloch, A., Bilbao, B., Mathison, C., Robinson, C. and Burton, C., 2022. Spreading like wildfire: The rising threat of extraordinary landscape fires.
66. Taylor, S. W. 2020. Atmospheric cascades shape wildfire activity and fire management decision spaces across scales– A conceptual framework for fire prediction. *Frontiers in Environmental Science*, 8, 527278.
67. Tepley, A.J., Parisien, M.A., Wang, X., Oliver, J.A. and Flannigan, M.D., 2022. Wildfire evacuation patterns and syndromes across Canada's forested regions. *Ecosphere*, 13(10), p.e4255.
68. Tibaldi, S., and F. Molteni, 1990: On the operational predictability of blocking. *Tellus*, 42A, 343–365, <https://doi.org/10.1034/j.1600-0870.1990.t01-2-00003.x>.
69. Tymstra, C., Jain, P. and Flannigan, M.D., 2021. Characterisation of initial fire weather conditions for large spring wildfires in Alberta, Canada. *International journal of wildland fire*, 30(11), pp.823-835.

70. USNIC, U.S. National Ice Center. 2008. IMS Daily Northern Hemisphere Snow and Ice Analysis at 1 km, 4 km, and 24 km Resolutions, Version 1.2 and 1.3 [G02156]. Boulder, Colorado USA. National Snow and Ice Data Center. <https://doi.org/10.7265/N52R3PMC>. Date Accessed 11-07-2023.
71. Van Wagner, C. E. 1987. Development and Structure of the Canadian Forest Fire Weather Index System. Technical Report 35. Canadian Forest Service, Ottawa, Ontario, Canada.
72. Wang, X., Parisien, M.A., Flannigan, M.D., Parks, S.A., Anderson, K.R., Little, J.M. and Taylor, S.W. 2014. The potential and realized spread of wildfires across Canada. *Global change biology*, 20(8), pp.2518-2530.
73. Wang, X., Oliver, J., Swystun, T., Hanes, C.C., Erni, S. and Flannigan, M.D. 2023. Critical fire weather conditions during active fire spread days in Canada. *Science of the total environment*, 869, p.161831.
74. Wang, X., Studens, K., Parisien, M.A., Taylor, S.W., Candau, J.N., Boulanger, Y. and Flannigan, M.D. 2020. Projected changes in fire size from daily spread potential in Canada over the 21st century. *Environmental Research Letters*, 15(10), p.104048.
75. Wang, X., Thompson, D., Marshall, G.A., Tymstra, C., Carr, R., Flannigan, M.D. 2015. Increasing frequency of extreme fire weather in Canada with climate change. *Clim. Chang.* 130(4), 573–586. <https://doi.org/10.1007/s10584-015-1375-5>.
76. Wang, Z., Wang, Z., Zou, Z. et al. 2023. Severe Global Environmental Issues Caused by Canada's Record-Breaking Wildfires in 2023. *Adv. Atmos. Sci.* <https://doi.org/10.1007/s00376-023-3241-0>
77. Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., ... and Mack, M. C. 2019. Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature*, 572(7770), 520-523.
78. Whitman, E., Parisien, M. A., Thompson, D. K., and Flannigan, M. D. 2019. Short-interval wildfire and drought overwhelm boreal forest resilience. *Scientific Reports*, 9(1), 18796.
79. Whitman, E., Parks, S.A., Holsinger, L.M. and Parisien, M.A. 2022. Climate-induced fire regime amplification in Alberta, Canada. *Environmental Research Letters*, 17(5), p.055003.
80. Wiitala, M. R. And Carlton, D. W. 1994. Assessing long term fire movement risk in wilderness fire management. In 12th Conf. on Fire and Forest Meteorology 187–194. Jekyll Island, GA.
81. Wotton, B.M. and Flannigan, M.D., 1993. Length of the fire season in a changing climate. *The Forestry Chronicle*, 69(2), pp.187-192. <https://doi.org/10.5558/TFC69187-2>,
82. Wotton, B.M., Flannigan, M.D. and Marshall, G.A., 2017. Potential climate change impacts on fire intensity and key wildfire suppression thresholds in Canada. *Environmental Research Letters*, 12(9), p.095003.

83. Xi, D.D., Dean, C.B. and Taylor, S.W., 2020. Modeling the duration and size of extended attack wildfires as dependent outcomes. *Environmetrics*, 31(5), p.e2619.

Figure 1: Infographic showing NBAC-M3 mapped fires, regional area burned and timeline and location of key events that occurred during the Canadian 2023 wildfire season.








Figure 2: Annual area burned for Canada from the National Burned Area Composite (NBAC; 1986–2022) and NBAC-M3 (Natural Resources Canada, 2023b) datasets. During 2023, 15 Mha burned, compared to the annual mean of 2.1 Mha (1986–2022, dashed line). The next largest annual area burned occurred in 1989 with 6.7 Mha.


a) snowmelt timing




b) snowmelt timing departure




c) maximum RZSM drying timing



d) maximum RZSM drying amount



e) May DC



f) June DC




Figure 3: Early 2023 fire season conditions for Canada. Panels: a) timing of snow melt; b) snow melt timing departure (days) from historical norm (2004–2022); c) mid-point timing of maximum Root Zone Soil Moisture (RZSM) drying in any two-week period between May and July; d) the corresponding maximum RZSM drying amount; e) standardized anomalies of the May Drought Code (DC); f) standardized anomalies of the June DC. Data sources described in methods.





Figure 4: 2023 fire season (May–October) anomalies relative to baseline period (1991–2020) for 2 m temperature a), total precipitation c) and Vapor Pressure Deficit (VPD; e); Fire season (May–October) mean value ranking during period 1940–2023 for 2 m temperature b), total precipitation d) and VPD f). Data derived from the ERA5 reanalysis.

a) Daily area burned



b) Area extreme fire weather




Figure 5: a) Daily area burned in 2023, by province/territory estimated from interpolated day of burn for fires ≥ 500 ha; b) the daily forested area in each province/territory with FWI values exceeding the 95th percentile of FWI (during the fire season). Nunavut is excluded from the plots due to its low total area burned and low forest cover.




Figure 6 : a) Total number of days exceeding local 95th percentile of FWI for 2023; b) Proportion forested area exceeding local 95th percentile of FWI (black) compared with mean (red) values for 1991–2020 (confidence intervals shaded for 5th and 95th percentiles); c) The number of blocking days between April 1st and October 31st 2023; d) Mean proportion forested area exceeding local 95th percentile of FWI during fire season (May-Sep) for each year (1940–2023).




Figure 7 a) Left: number of fires > 50k ha for each fire size class for historical period (1986–2022, annual mean) and for 2023; Right: proportion of total area burned for each fire size class for 1986–2022 and 2023; b) Fire size as function of potential spread days (extreme fire weather days; FWI₉₅) for Western Canada (west of -85 deg longitude) and eastern Canada (east of -85 degrees longitude). c) Distribution of fire sizes by start month for all fires > 1,000 ha. (colored outliers correspond to fires > 100,000 ha). d) Cumulative daily area burned for fires starting in each month between May–October.




Figure 8: Panel a) The number of days in 2023 with daily mean $\text{PM}_{2.5}$ greater than $27 \mu\text{g}/\text{m}^3$; b) the maximum daily value of $\text{PM}_{2.5}$ that occurred during 2023; panel c) the number of air quality bulletins each month issued by Environment and Climate Change Canada between 2017 and 2023 (Health and Air Quality Forecast Services Program, MSC-ECCC, December 2023); panel d) The number of annual evacuation orders and evacuees from 1980–2023 using data from the Canadian wildfire evacuation database.

Supplementary Material for:
Canada Under Fire – Drivers and Impacts of the Record-breaking 2023 Wildfire Season
Jain et al. 2024

Table S1: Regional summaries for total area burned in 2023

Region	2023 Area burned (ha)	2023 rank (1986-2023)
Quebec	4,258,416	1
Northwest Territories	3,574,813	1
Alberta	2,735,063	1
British Columbia	2,342,294	1
Saskatchewan	1,137,683	3
Yukon Territory	379,668	2
Ontario	341,599	6
Manitoba	142,334	18
Maritimes	24,064	2
Newfoundland and Labrador	19,257	18
Nunavut	2,323	10

Table S2 : Data corresponding to Fig 10c in main text. Air Quality (AQ) Alert Bulletins issued by MSC-ECCC, 2017-2023. Data representing the number of all AQ Alert bulletins issued throughout the period. Each AQ alert will have a minimum of 2 bulletins issued (issuance and termination), plus a number of continued bulletins based on the persistence of the event. Each AQ Alert's area of coverage varies in size based on the geographical extent of the AQ event.

	2017	2018	2019	2020	2021	2022	2023
Jan	122	97	53	32	45	79	34
Feb	77	61	67	87	52	43	59
Mar	50	129	366	49	102	90	142
Apr	20	66	68	0	20	10	28
May	31	53	108	33	104	19	520
Jun	9	31	185	48	160	64	1412
Jul	223	217	273	22	728	245	932
Aug	413	589	66	70	407	191	979
Sep	385	54	15	96	101	247	762
Oct	32	22	6	9	77	153	72
Nov	85	67	17	12	27	138	55
Dec	127	37	69	27	10	43	21
Total	1574	1423	1293	485	1833	1322	5016

Table S3. Ten largest wildfire evacuations on record, by number of evacuees (Canadian Forest Service, 2023). Five of the ten largest evacuation events (by number of evacuees) since 1980 occurred in 2023.

Evacuation Date	Province/Territory	Location	Evacuees
2016-05-03	AB	Fort McMurray	88000
2003-08-18	BC	Kelowna	33050
2023-08-16	NT	Yellowknife	21720
2023-08-17	BC	West Kelowna	19809
2023-05-28	NS	Halifax	16400
2009-07-18	BC	West Kelowna	11000
2017-07-15	BC	Williams Lake	10753
2023-08-17	BC	Kelowna	9757
2023-05-05	AB	Edson	8414
1998-08-10	BC	Salmon Arm	8000

North American Drought Monitor

To evaluate general drought conditions in Canada, we used summary statistics from the North American Drought Monitor (NADM, National Drought Mitigation Center). The statistics consists of a monthly time series of drought classification for each province/territory. Classifications include D0 (Abnormally Dry), D1 (Moderate Drought), D2 (Severe Drought), D3 (Extreme Drought), D4 (Exceptional Drought). We retrieved data for the percentage area with conditions D0-D4, i.e., Abnormally dry or in drought.




Fig S1: Monthly time series of percentage of each province or territory that is abnormally dry or in drought conditions for 2022 and 2023. The climatological mean (2002-2023) is shown by the blue line with 95% confidence intervals shaded. Data from the North American Drought Monitor (see methods). See Lawrimore et al 2002 for drought definitions.

Extreme Fire Weather

The 95th percentile of the Initial Spread Index (ISI) and Build-up Index (BUI), indices which are components (output indices) of the FWI System (Van Wagner 1987) showed similar regional and temporal trends to FWI₉₅ in 2023 (Fig. 6 main text). Because of its dependence on wind speed, the ISI is related to wind-driven fire regimes, whereas the BUI - which depends on the Drought Code (DC) and Duff Moisture Code (DMC) - is related to drought-driven fire regimes.




Fig. S2: a) Total number of days exceeding local 95th percentile of ISI for 2023; c) Total forested area exceeding local 95th percentile of ISI (black) compared with 5th and 95th percentiles (shaded) and mean (red) values for

1991-2020; e) Mean percent daily forested area exceeding local 95th percentile of ISI during fire season (May-Sept) for each year (1940-2023); b) Total number of days exceeding local 95th percentile of BUI for 2023; d) Total forested area exceeding local 95th percentile of BUI (black) compared with 5th and 95th percentiles (shaded) and mean (red) values for 1991-2020; f) Mean percent daily forested area exceeding local 95th percentile of BUI during fire season (May-Sept) for each year (1940-2023).




Figure. S3: Number of active fires > 200 ha for 2019–2023, by reported suppression action (fire response) using archived CIFFC fire agency data.




Fig. S4: Total annual area burned in Canada as a function of number of fires > 200ha using NBAC data (1986–2023). The four years with the greatest area burned are indicated. The relationship follows an approximate power law. Large fires > 200 ha account for 95% of area burned in Canada. In 2023, 2.6 times the average number of large fires burned about 7 times the average area.

Population exposure to wildfire smoke

We combined population data with the 2023 modeled surface PM_{2.5} concentrations (see methods in main text) to determine population exposure to wildfire smoke during 2023. For regional exposure (province or territory) we used the gridded population downloaded from:

<https://sedac.ciesin.columbia.edu/data/collection/gpw-v4/sets/browse>. For urban centers we used the Census Metropolitan areas from the Government of Canada open data portal:

<https://open.canada.ca/data/en/dataset/096b2a17-2755-40fe-b750-1c489c5e6b6a>. For regional wildfire smoke exposure, we calculated a weighted mean of days of PM_{2.5} > 27µgm⁻³ using the gridded population density as the weighting factor. For census metropolitan areas we extracted days of PM_{2.5} > 27µgm⁻³ to each area. The resulting values represented the per capita number of poor air quality experienced in 2023 .

Table S4: Poor air quality days per capita for each province/territory of Canada, defined as the population weighted mean of days with PM_{2.5} greater than 27µgm⁻³.

Region	Smoky days per capita
British Columbia	3.5
Alberta	17.1
Saskatchewan	10.6
Manitoba	7.9
Ontario	8.7
Quebec	6.6
New Brunswick	0.2
Yukon	1.7
Nunavut	0.4
Newfoundland and Labrador	0.7
Nova Scotia	0.1
Northwest Territories	43.6
Prince Edward Island	0
Canada	8

Table S5: Mean poor air quality days and maximum daily PM_{2.5} for the 10 most populous census metropolitan areas.

Census metropolitan area	population	poor AQ days	max daily PM2.5
Toronto	6,202,225	14	129
Montreal	4,291,732	12	135
Vancouver	2,642,825	4	112
Ottawa - Gatineau	1,488,307	11	215
Calgary	1,481,806	18	103
Edmonton	1,418,118	21	219
Quebec	839,311	4	117

Winnipeg	834,678	10	70
Hamilton	785,184	11	127
Kitchener - Cambridge - Waterloo	575,847	7	97

Daily growth of ten largest wildfires in 2023.

We considered individual cases of the 10 largest mapped fires of 2023 by examining daily area burned, the corresponding mean daily FWI values and the daily precipitation at the location of the fire.








Fig. S5: Location of the 10 largest wildfires that burned in Canada in 2023.





NBAC ID = 346
Agency ID = QC-F218





NBAC ID = 354
Agency ID = AB-HWF-036-2023, BC-2023-G92498
Long Lake Complex, Ekwan Tower





NBAC ID = 313
Agency ID = NT-SS022-23





NBAC ID = 359
Agency ID = BC-G80280
Donnie Creek



NBAC ID = 321
Agency ID = NT-SS052-23
Hay River



NBAC ID = 349
Agency ID = SK-23LX-SMITH






Fig. S6: Interpolated day of year of burning for the 10 largest wildfires that burned in Canada in 2023. Also shown for each fire is daily area burned, Fire Weather Index values at the fire location (with extreme fire weather days defined by days exceeding the 95th percentile of the local climatology fire season FWI, shown by red points), daily precipitation and 100m wind speed (mean values were taken over each fire perimeter).

References

1. Lawrimore, J., Heim, R.R., Svoboda, M., Swail, V. and Englehart, P.J., 2002. Beginning a new era of drought monitoring across North America. *Bulletin of the American Meteorological Society*, 83(8), pp.1191-1192.
2. National Drought Mitigation Center, University of Nebraska-Lincoln. "Statistics." North American Drought Monitor website. <https://droughtmonitor.unl.edu/NADM/Statistics.aspx>. Retrieved November 1st 2023.